

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 03/051842 A2

(51) International Patent Classification⁷: **C07D 213/00**

DK-2880 Bagsværd (DK). JACOBSEN, Poul; Morænevej 2A, DK-3500 Slangerup (DK).

(21) International Application Number: PCT/DK02/00853

(22) International Filing Date:
13 December 2002 (13.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2001 01879 14 December 2001 (14.12.2001) DK
PA 2002 00645 30 April 2002 (30.04.2002) DK
PA 2002 01000 27 June 2002 (27.06.2002) DK
PA 2002 01562 11 October 2002 (11.10.2002) DK

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).

Published:

— without international search report and to be republished upon receipt of that report

(72) Inventors: EBDROP, Søren; Nymarksvej 11, DK-4000 Roskilde (DK). HANSEN, Holger, Claus; Annexgårdsparken 43, DK-3500 Værløse (DK). VEDSØ, Per; Dalgas Have 4, 2th, DK-2000 Frederiksberg (DK). CORNELIS DE JONG, Johannes; Møllemarken 16, 1th,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/051842 A2

(54) Title: COMPOUNDS AND USE THEREOF FOR DECREASING ACTIVITY OF HORMONE-SENSITIVE LIPASE

(57) Abstract: Use of compounds to inhibit hormone-sensitive lipase, pharmaceutical compositions comprising the compounds, methods of treatment employing these compounds and compositions, and novel compounds. The present compounds are inhibitors of hormone-sensitive lipase and may be useful in the treatment and/or prevention of medical disorders where a decreased activity of hormone-sensitive lipase is desirable.

USE OF COMPOUNDS FOR DECREASING ACTIVITY OF HORMONE-SENSITIVE LI- PASE.

5 FIELD OF THE INVENTION

The present invention relates to compounds, compositions containing them, and their use for treating medical disorders where it is desirable to modulate the activity of hormone-sensitive lipase.

10

BACKGROUND OF THE INVENTION

The overall energy homeostasis of a mammalian system requires a high degree of regulation to ensure the availability of the appropriate substrate at the appropriate time. Plasma glucose levels rise during the post-prandial state, to return to pre-prandial levels within 2-3 hours. During these 2-3 hours, insulin promotes glucose uptake by skeletal muscle and adipose tissue and decreases the release of free fatty acids (FFA) from adipocytes, to ensure that the two substrates do not compete with each other. When plasma glucose levels fall, an elevation in plasma FFA is necessary to switch from glucose to fat utilization by the various tissues.

In individuals with insulin resistance, FFA levels do not fall in response to insulin, as they do in normal individuals, preventing the normal utilization of glucose by skeletal muscle, adipose and liver. Furthermore, there is a negative correlation between insulin sensitivity and plasma FFA levels.

Hormone-sensitive lipase (HSL) is an enzyme, expressed primarily in adipocytes, that catalyses the conversion of triglycerides to glycerol and fatty acids. It is through the regulation of this enzyme that the levels of circulating FFA are modulated. Insulin leads to the inactivation of HSL with a subsequent fall in plasma FFA levels during the post-prandial state, followed by the activation of the enzyme when the insulin concentration falls and catecholamines rise during the post-absorptive period. The activation of HSL leads to an increase in plasma FFA, as they become the main source of energy during fasting.

35

The activation-inactivation of HSL is primarily mediated through the cAMP-protein kinase A and AMP-dependent kinase pathways. There are compounds like nicotinic acid and its derivatives, that decrease the activation of HSL via these pathways and cause a decrease in lipolysis that leads to a reduction in the FFA levels. These drugs have a beneficial effect in the utilization of glucose and in the normalization of the excess triglyceride synthesis seen in patients with elevated FFA. However, since these pathways are used by other processes in the body, these drugs have severe side effects.

We have found compounds that specifically inhibit the lipolytic activity of HSL and lead to a decrease in plasma FFA levels. These compounds can be used to treat disorders where a decreased level of plasma FFA is desired, such as insulin resistance, syndrome X, dyslipidemia, abnormalities of lipoprotein metabolism.

One object of the present invention is to provide compounds and pharmaceutical compositions that inhibit the lipolytic activity of HSL. A further object is to provide compounds which have good pharmaceutical properties such as solubility, bioavailability etc.

DEFINITIONS

The following is a detailed definition of the terms used to describe the compounds of the invention.

"Halogen" designates an atom selected from the group consisting of F, Cl, Br and I.

The term "C₁₋₆-alkyl" in the present context designates a saturated, branched or straight hydrocarbon group having from 1 to 6 carbon atoms. Representative examples include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, tert-pentyl, n-hexyl, isohexyl and the like.

The term "C₂₋₆-alkyl" in the present context designates a saturated, branched or straight hydrocarbon group having from 2 to 6 carbon atoms. Representative examples include, but are not limited to, ethyl, n-propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, tert-pentyl, n-hexyl, isohexyl and the like.

The term "C₁₋₆-alkoxy" in the present context designates a group -O-C₁₋₆-alkyl wherein C₁₋₆-alkyl is as defined above. Representative examples include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, butoxy, isobutoxy, *sec*-butoxy, *tert*-butoxy, n-pentoxy, isopentoxy, neopentoxy, *tert*-pentoxy, n-hexoxy, isohexoxy and the like.

5

The term "C₃₋₆-alkoxy" in the present context designates a group -O-C₃₋₆-alkyl wherein C₃₋₆-alkyl is a saturated, branched or straight hydrocarbon group having from 3-6 carbon atoms. Representative examples of C₃₋₆-alkoxy include, but are not limited to, n-propoxy, isopropoxy, butoxy, isobutoxy, *sec*-butoxy, *tert*-butoxy, n-pentoxy, isopentoxy, neopentoxy, *tert*-pentoxy, n-hexoxy, isohexoxy and the like.

10

The term "C₂₋₆-alkenyl" as used herein, represent an olefinically unsaturated branched or straight hydrocarbon group having from 2 to 6 carbon atoms and at least one double bond. Examples of such groups include, but are not limited to, vinyl, 1-propenyl, 2-propenyl, allyl, iso-propenyl, 1,3-butadienyl, 1-butenyl, hexenyl, pentenyl and the like.

15

The term "C₃₋₁₀-cycloalkyl" as used herein represents a saturated mono-, bi-, tri- or spirocarbocyclic group having from 3 to 10 carbon atoms. Representative examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, bicyclo[3.2.1]octyl, spiro[4.5]decyl, norpiny, norbonyl, norcaryl, adamantyl and the like.

20

The term "C₃₋₈-heterocyclyl" as used herein represents a saturated 3 to 8 membered ring containing one or more heteroatoms selected from nitrogen, oxygen and sulfur. Representative examples are pyrrolidyl, piperidyl, piperaziny, morpholinyl, thiomorpholinyl, aziridinyl, tetrahydrofuranyl and the like.

25

The term "aryl" as used herein represents a carbocyclic aromatic ring system being either monocyclic, bicyclic, or polycyclic, such as phenyl, biphenyl, naphthyl, anthracenyl, phenanthrenyl, fluorenyl, indenyl, pentalenyl, azulenyl, biphenylenyl and the like. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic aromatic systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 1,2,3,4-tetrahydronaphthyl, 1,4-dihydronaphthyl and the like.

30

The term "aryloxy" as used herein represents an aryl which is linked via an oxygen atom, e.g. phenoxy, 1-naphthyloxy, 2-naphthyloxy and the like.

35

The term "heteroaryl" as used herein represents a heterocyclic aromatic ring system containing one or more heteroatoms selected from nitrogen, oxygen and sulfur such as furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 5 pyranlyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiadiazinyl, indolyl, isoindolyl, benzofuranyl, benzothiophenyl (thianaphthenyl), indazolyl, benzimidazolyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, purinyl, quinazolinyl, quinoliziny, quinolinyl, iso- 10 quinolinyl, quinoxaliny, naphthyridinyl, pteridinyl, carbazolyl, azepinyl, diazepinyl, acridinyl and the like. Heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 2,3-dihydrobenzofuranyl, 3,4-dihydroisoquinolinyl, pyrrolinyl, pyrazolinyl, indolinyl, oxazolidinyl, oxazoliny, oxazepinyl and the like.

15

The term "perhalomethyl" as used herein designates a methyl moiety substituted with three halogen atoms. Non-limiting examples of perhalomethyl are CF_3 , CCl_3 , and CF_2Cl .

20

The term "perhalomethoxy" as used herein designates a perhalomethyl linked via an oxygen atom, e.g. $-\text{O}-\text{CF}_3$, $-\text{O}-\text{CCl}_3$, and $-\text{O}-\text{CF}_2\text{Cl}$

25

The term "ring system" as used herein includes aromatic as well as non-aromatic ring moieties, which may be monocyclic, bicyclic or polycyclic, and they encompass moieties with zero, one or more heteroatoms selected from nitrogen, oxygen and sulphur. Non-limiting examples of such ring systems are aryl, C_{3-8} -heterocyclyl and heteroaryl.

30

The term "heterocyclic system" as used herein includes aromatic as well as non-aromatic ring moieties, which may be monocyclic, bicyclic or polycyclic, and containing in their ring structure one or more heteroatoms selected from nitrogen, oxygen and sulfur. Non-limiting examples of such heterocyclic systems are C_{3-8} -heterocyclyl and heteroaryl.

Certain of the above defined terms may occur more than once in the structural formulae, and upon such occurrence each term shall be defined independently of the other.

The term "optionally substituted" as used herein means that the groups in question are either unsubstituted or substituted with one or more of the substituents specified. When the groups in question are substituted with more than one substituent the substituents may be the same or different.

5

The term "optionally covalently bound" as used herein means that the substituents in question are either not covalently bound to each other or the substituents are directly connected to each other by a covalent bond. A non-limiting example of such optionally covalently bound substituents is $-NR^1R^2$ wherein R^1 is ethyl and R^2 is propyl which provided that the substituents, ethyl and propyl, are optionally covalently bound may be ethyl-propyl-amino, 1-piperidyl, 3-methyl-1-pyrrolidyl or 2,3-dimethyl-1-azetidyl.

10

The term "hydrolysable group" as used herein means a group which can be hydrolysed at certain chemical conditions, i.e. an internal covalent bond in the group can be cleaved to give two compounds. A hydrolysable group does not have to be hydrolysed during carrying out the present invention. For instance, the hydrolysable group $-C(=X)-L$ can be hydrolysed to give the products $-C(=X)OH$ and HL , which means that the covalent bond between $-C(X)$ and L is cleaved. Examples of hydrolysable groups are carbamates, esters and amides.

15

The terms "disease", "condition" and "disorder" as used herein are used interchangeably to specify a state of a patient which is not the normal physiological state of man.

20

The term "treatment" as used herein means the management and care of a patient having developed a disease, condition or disorder, as well as the management and care of an individual at risk of developing the disease, condition or disorder prior to the clinical onset of said disease, condition or disorder. The purpose of treatment is to combat the disease, condition or disorder, as well as to combat the development of the disease, condition or disorder. Treatment includes the administration of the active compounds to prevent or delay the onset of the symptoms or complications and to eliminate or control the disease, condition or disorder as well as to alleviate the symptoms or complications associated with the disease, condition or disorder.

25

30

The term "effective amount" as used herein means a dosage which is sufficient in order for the treatment of the patient to be effective compared with no treatment.

35

The term "modulate" as used herein means to influence, i.e. to modulate a parameter means to influence that parameter in a desired way. Examples are to modulate insulin secretion from beta cells and to modulate the plasma level of free fatty acids.

- 5 The term "medicament" as used herein means a pharmaceutical composition suitable for administration of the pharmaceutically active compound to a patient.


The term "pharmaceutically acceptable" as used herein means suited for normal pharmaceutical applications, i.e. giving rise to no adverse events in patients etc.

10

DESCRIPTION OF THE INVENTION

The present invention relates to the use of a compound of the general formula I

15



20

wherein R¹ is selected from hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

25

R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl,

30

C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₆-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₆-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₆-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and


X is O or S; and

L is a group such that -C(=X)-L is a hydrolysable group; or

a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs

for inhibition of the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters.

Another aspect of the invention relates to use of a compound of the general formula I



wherein R¹ is selected from hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₆-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-

alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

20 X is O or S; and

L is a group such that -C(=X)-L is a hydrolysable group; or

25 a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs

for the preparation of a medicament for the treatment of any disorder where it is desirable to

30 modulate the plasma level of free fatty acids, glycerol, LDL-cholesterol, HDL-cholesterol, insulin and/or glucose; and/or

modulate intracellular triacylglycerol and cholesterol ester stores, intracellular level of fatty acids, fatty acid esters such as diacylglycerols, phosphatidic acids, long chain acyl-CoA's as well as citrate or malonyl-CoA; and/or

- 5 increase insulin sensitivity in adipose tissue, skeletal muscle, liver or pancreatic β cells; and/or

modulate insulin secretion from pancreatic β cells.


- 10 Another aspect of the invention is the use of a compound of the general formula I for the preparation of a pharmaceutical medicament.

In one embodiment the present invention relates to the use of a compound of the general formula (I) for the preparation of a medicament for the treatment of insulin resistance, diabetes type 1, diabetes type 2, metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

15

Another aspect of the invention is pharmaceutical compositions comprising a compound of formula I

20



wherein R^1 is selected from hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{3-10} -cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

25

R^2 is selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl,

30

heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

X is O or S; and

L is a group such that -C(=X)-L is a hydrolysable group; or


a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.


In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein X is O. In another embodiment, the invention is concerned with pharmaceutical compositions comprising a compound of formula (I) and uses of the compounds of formula (I), wherein X is O. In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the group L contains an O, via which L is bound to the C in formula (I). In another embodiment, the invention is concerned with pharmaceutical compositions comprising a compound of formula (I) and uses of the compounds of formula (I), wherein the group L contains an O, via which L is bound to the C in formula (I).


In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the group L contains a N, via which L is bound to the C in formula (I).

5 In another embodiment, the invention is concerned with pharmaceutical compositions comprising a compound of formula (I) and uses of the compounds of formula (I), wherein the group L contains a N, via which L is bound to the C in formula (I).

10 In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the group L is selected from the group consisting of







wherein Y is O or S; and

R^{a1} , R^{a2} , R^{a3} , R^{a4} , R^{a5} and R^{a6} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, or C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl,

wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein at least one of R^{a1}, R^{a2}, R^{a3}, R^{a4}, R^{a5} and R^{a6} is F.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the group L is an optionally substituted -O-phenyl, via which L is bound to the C in formula (I). In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperazine, said piperazine being bound to the C in formula (I).

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperidine, said piperidine being bound to the C in formula (I).

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R¹ is selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R^1 is methyl.

5 In yet another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R^1 is phenyl.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R^2 is a heteroaryl.

10 In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein R^1 is methyl and R^2 is phenyl.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the
15 group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 and R^2 are covalently bound to each other so that the group R^1 -N- R^2 forms a piperazine, said piperazine being bound to the C in formula (I).

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the
20 group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 and R^2 are covalently bound to each other so that the group R^1 -N- R^2 forms a piperidine, said piperidine being bound to the C in formula (I).

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein the
25 group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 is methyl and R^2 is phenyl.

In another embodiment, the invention is concerned with pharmaceutical compositions containing a compound of formula (I) and uses of the compounds of formula (I), wherein pK_a of
30 the group L is between 4 and 12, between 6 and 12, between 7 and 12, between 8 and 12, preferably between 8.5 to 11.5, and most preferable between 9.0 to 11.0.

In another embodiment, the invention is concerned with pharmaceutical compositions containing the compounds of formula (I) and uses of the compounds of formula (I), wherein administration of said compound is by oral administration

In another embodiment, the invention is concerned with pharmaceutical compositions containing the compounds of formula (I) and uses of the compounds of formula (I), the nasal, transdermal, pulmonal, or parenteral route.


- 5 Another aspect of the invention is a method of treating any disorder where it is desirable to inhibit the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters, wherein said method comprises the use as described above.

- 10 In another embodiment, the invention is concerned with a method of treating any disorder where it is desirable to modulate the plasma level of free fatty acids or to modulate the handling, storage and oxidation of intracellular fatty acid and cholesterol, wherein said method comprises the use as described above.

- In another embodiment, the invention is concerned with said method, wherein said disorder is selected from the group consisting of insulin resistance, diabetes type 1, diabetes type 2, 15 metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

The present invention also relates to compounds of the general formula II

20



wherein R¹ is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

25

- R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo,
- 30

amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and


R³ is selected from hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

X is O or S; or

a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

5

In an aspect of the invention, compounds are of the general formula III



10

wherein R^{1a} and R^{2a} are independently selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl,

15

wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano and nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋

20

₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, perhalomethyl and perhalomethoxy; and

25

wherein R^{1a} is optionally covalently bound to R^{2a} by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R^{1a} and R^{2a} are bound; and

30

R^{4a}, R^{5a}, R^{6a} and R^{7a} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted by one or more substituents selected from hydroxy, sulfanyl, sulfo, oxo, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl,

C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, perhalomethyl and perhalomethoxy; and

5

A^{1a} is N or C-R^{8a}; A^{2a} is N or C-R^{9a}; A^{3a} is N or C-R^{10a}; A^{4a} is N or C-R^{11a}; and A^{5a} is N or C-R^{12a}; and

wherein R^{8a}, R^{9a}, R^{10a}, R^{11a} and R^{12a} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, perhalomethyl and perhalomethoxy.

In one embodiment, the invention is concerned with compounds of formula (III), wherein R^{2a} is phenyl, optionally substituted by halogen or methyl.

In another embodiment, the invention is concerned with compounds of formula (III), wherein R^{1a} is selected from methyl and ethyl, optionally substituted by one or more halogen.

In another embodiment, the invention is concerned with compounds of formula (III), wherein R^{1a} and R^{2a} are covalently bound so as to form a ring system with the N-atom to which they are bound, wherein said ring system is a piperidine, piperazine, morpholine, or thiomorpholine.


In another embodiment, the invention is concerned with compounds of formula (III), wherein R^{4a}, R^{5a}, R^{6a} and R^{7a} are independently selected from hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy,

-C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of formula (III), wherein A^{1a}, A^{2a}, A^{3a}, A^{4a} and A^{5a} are independently selected from N, CH, CF, C-Cl and C-CF₃.

5

In an aspect of the invention, compounds are of the general formula IV



- 10 wherein R^{1b} and R^{2b} are independently selected from C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which may optionally be substituted with one or more substituents selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy;

- 25 wherein R^{1b} is optionally covalently bound to R^{2b} by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R^{1b} and R^{2b} are bound;

- 30 R^{5b} and R^{6b} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro,

5 sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, perhalomethyl and perhalomethoxy;

10 R^{4b} and R^{7b} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, perhalomethyl and perhalomethoxy; and

A^{1b} is N or C-R^{8b}; A^{2b} is N or C-R^{9b}; A^{3b} is N or C-R^{10b}; A^{4b} is N or C-R^{11b}; and A^{5b} is N or C-R^{12b}; and

20 wherein R^{8b}, R^{9b}, R^{10b}, R^{11b} and R^{12b} are selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, perhalomethyl and perhalomethoxy;

30 wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4b}, R^{5b}, R^{6b}, R^{7b}, R^{8b}, R^{9b}, R^{10b}, R^{11b} and R^{12b}.

35 In one embodiment, the invention is concerned with compounds of formula IV, wherein R^{2b} is phenyl, optionally substituted by halogen.


In another embodiment, the invention is concerned with compounds of the general formula IV, wherein R^{1b} is selected from methyl and ethyl, optionally substituted by one or more halogen.

In another embodiment, the invention is concerned with compounds of the general formula IV, wherein R^{1b} and R^{2b} are covalently bound so as to form a ring system with the N-atom to which they are bound, wherein the ring system is a piperidine, piperazine, morpholine, or thiomorpholine.

In another embodiment, the invention is concerned with compounds of the general formula IV, wherein R^{4b} , R^{5b} , R^{6b} and R^{7b} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula IV, wherein A^{1b} , A^{2b} , A^{3b} , A^{4b} and A^{5b} are independently selected from N, CH and CF.

In an aspect of the invention, compounds are of the general formula V



wherein R^{4c} , R^{5c} , R^{6c} , R^{7c} and R^{8c} are independently selected from hydrogen, hydroxy, sulfanyl, amino, halogen, cyano, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl,

heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

- R^{8c}, R^{10c}, R^{11c}, R^{12c} and R^{13c} are independently selected from hydrogen, sulfanyl, amino, halogen, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and
- wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4c}, R^{5c}, R^{6c}, R^{7c} and R^{8c}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{8c}, R^{10c}, R^{11c}, R^{12c} and R^{13c}; and

at least one of R^{4c}, R^{5c}, R^{6c}, R^{7c}, R^{8c}, R^{9c}, R^{10c}, R^{11c}, R^{12c} and R^{13c} are different from hydrogen.

In one embodiment, the invention is concerned with compounds of formula V, wherein R^{9c}, R^{10c}, R^{11c}, R^{12c} and R^{13c} are selected from H and F.


In another embodiment, the invention is concerned with compounds of the general formula V, wherein R^{4c} , R^{5c} , R^{6c} , R^{7c} and R^{8c} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

- 5 In another embodiment, the invention is concerned with compounds of the general formula V, where there are no covalent bonds between any of the substituents R^{9c} , R^{10c} , R^{11c} , R^{12c} and R^{13c} .

In another embodiment, the invention is concerned with compounds of the general formula V, where there are no covalent bonds between any of the substituents R^{4c} , R^{5c} , R^{6c} , R^{7c} and R^{8c} .

10

In an aspect of the invention, compounds are of the general formula VI



- 15 wherein R^{4d} and R^{5d} are independently selected from hydrogen, hydroxy, sulfanyl, amino, C_{2-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, amino, C_{2-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy: and
- 20
- 25

R^{6d} , R^{7d} , R^{8d} , R^{9d} , R^{10d} , R^{11d} , R^{12d} , R^{13d} and R^{14d} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy:

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4d} , R^{5d} , R^{6d} , R^{7d} , R^{8d} and R^{9d} ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{10d} , R^{11d} , R^{12d} , R^{13d} and R^{14d} .

In one embodiment, the invention is concerned with compounds of the general formula VI, wherein R^{4d} and R^{5d} are H.


In another embodiment, the invention is concerned with compounds of the general formula VI, wherein R^{6d} , R^{7d} , R^{8d} and R^{9d} are selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula VI, wherein R^{10d} , R^{11d} , R^{12d} , R^{13d} and R^{14d} are selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula VI, wherein there are no covalent bonds between any of the substituents R^{4d} , R^{5d} , R^{6d} , R^{7d} , R^{8d} and R^{9d} .

In another embodiment, the invention is concerned with compounds of the general formula VI, wherein there are no covalent bonds between any of the substituents R^{10d} , R^{11d} , R^{12d} , R^{13d} and R^{14d} .

In an aspect of the invention, compounds are of the general formula VII




wherein R^{4e} is selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, C_{2-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{5e} is selected from hydrogen, F, cyano, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and

C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy;


wherein R^{4e} and R^{5e} may be covalently bound to each other by a C-O bond.

In one embodiment, the invention is concerned with compounds of the general formula VII, wherein R^{4e} or R^{5e} is selected from the group consisting of



wherein R^{6e}, R^{7e}, R^{8e}, R^{9e} and R^{10e} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl.


In another embodiment, the invention is concerned with compounds of the general formula VII, wherein R^{4e} or R^{5e} is selected from the group consisting of



wherein R^{6e} , R^{7e} , R^{8e} , R^{9e} and R^{10e} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy and C_{2-6} -alkenyl.

In another embodiment, the invention is concerned with compounds of the general formula VII, wherein at least one of the substituents R^{6e} , R^{7e} , R^{8e} , R^{9e} and R^{10e} are selected from the group consisting of F and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula VII, wherein R^{4e} and R^{5e} are connected by a C-O bond to form the compound



20


wherein R^{11e} is selected from hydrogen, hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -

25

alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

R^{12e} is selected from hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula VIII



wherein R^{4f}, R^{5f}, R^{6f}, R^{7f}, R^{9f} and R^{10f} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, het-

eroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: with the proviso that R^{4f} and R^{5f} are not both methoxy; and

R^{8f} is selected from hydrogen, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy;


wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4f}, R^{5f}, R^{6f}, R^{7f} and R^{8f}; and

wherein there may optionally be a covalent bond between R^{9f} and R^{10f}.

In one embodiment, the invention is concerned with compounds of the general formula VIII, wherein R^{9f} and R^{10f} are covalently bound so as to form a ring system with the C-atom to which they are bound.

In another embodiment, the invention is concerned with compounds of the general formula VIII, wherein said ring system is a cycloalkyl, phenyl, heteroaryl, piperidine, piperazine, morpholine, or thiomorpholine.

In another embodiment, the invention is concerned with compounds of the general formula VIII, wherein R^{4f} and R^{8f} are connected by a C-O bond to form the compound




wherein R^{11f} is selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

R^{17f} is selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl; and

wherein there may optionally be a covalent bond between R^{9f} and R^{10f} .

In an aspect of the invention, compounds are of the general formula IX



wherein R^{4g} , R^{5g} , R^{6g} , R^{7g} , R^{8g} , R^{9g} , R^{10g} and R^{11g} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino,

cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo,

5 halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl,

10 heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{12g} is selected from the group consisting of -C(=O)-, -C(=O)NH-, -CH₂-, -CH₂CH₂-, -CHR^{15g}-, -CH₂CHR^{15g}-, -CHR^{15g}-CH₂-, -NH-, -NR^{15g}-, -NHC(=O)-, -NR^{15g}-C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-;

15 wherein R^{15g} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino,

20 cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4g}, R^{5g}, R^{6g}, R^{7g} and R^{15g}; and

25

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{6g}, R^{9g}, R^{10g}, R^{11g} and R^{15g}; and

30 R^{13g} is selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl,

35 C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋

6-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{14g} is selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, C₂₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of C₂₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

wherein R^{13g} and R^{14g} may optionally be covalently bound to each other.

In one embodiment, the invention is concerned with compounds of the general formula IX, wherein R^{13g} and R^{14g} are covalently bound so as to form a ring system with the N-atom to which they are bound.


In another embodiment, the invention is concerned with compounds of the general formula IX, wherein said ring system is a piperidine, piperazine, morpholine, or thiomorpholine.

In another embodiment, the invention is concerned with compounds of the general formula IX, wherein there are no covalent bonds between any of the substituents R^{4g} , R^{5g} , R^{6g} , R^{7g} and R^{15g} .

In another embodiment, the invention is concerned with compounds of the general formula IX, wherein there are no covalent bonds between any of the substituents R^{8g} , R^{9g} , R^{10g} , R^{11g} and R^{15g} .

In another embodiment, the invention is concerned with compounds of the general formula IX, wherein R^{4g} , R^{5g} , R^{6g} , R^{7g} , R^{8g} , R^{9g} , R^{10g} , R^{11g} are selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-$ OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula X



15

wherein R^{4h} and R^{5h} are independently selected from cyano, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy,

sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

5 R^{6h}, R^{7h}, R^{8h}, R^{9h}, R^{10h}, R^{12h}, R^{14h} and R^{15h} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

15 R^{11h} and R^{13h} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, perhalomethyl, perhalomethoxy, C₂₋₆-alkyl, methoxy, C₃₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfanyl, amino, C₂₋₆-alkyl, C₃₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

30 wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4h}, R^{5h}, R^{6h}, R^{7h}, R^{8h}, R^{9h}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{10h}, R^{11h}, R^{12h}, R^{13h}, R^{14h} and R^{15h}.

In one embodiment, the invention is concerned with compounds of the general formula X, wherein R^{6h} , R^{7h} , R^{8h} , R^{9h} , R^{10h} , R^{12h} and R^{14h} are selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

- 5 In another embodiment, the invention is concerned with compounds of the general formula X, wherein there are no covalent bonds between R^{4h} , R^{5h} , R^{6h} , R^{7h} , R^{8h} and R^{9h} .


In another embodiment, the invention is concerned with compounds of the general formula X, wherein there are no covalent bonds between R^{10h} , R^{11h} , R^{12h} , R^{13h} , R^{14h} and R^{15h} .

- In another embodiment, the invention is concerned with compounds of the general formula X,
10 wherein R^{15h} is hydrogen.

In another embodiment, the invention is concerned with compounds of the general formula X, wherein R^{11h} and R^{13h} are selected from the group consisting of hydrogen, F, Cl, C_{2-6} -alkyl, methoxy, C_{3-6} -alkoxy, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

15

In an aspect of the invention, compounds are of the general formula XI



- 20 R^{5i} is selected from hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-6} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-6} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl,
25 heteroaryl, C_{3-6} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-6} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy,

C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R⁶ⁱ, R⁷ⁱ, R⁸ⁱ, R⁹ⁱ, R¹⁰ⁱ, R¹⁴ⁱ and R¹⁵ⁱ are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R¹¹ⁱ and R¹³ⁱ are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, methoxy, C₃₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfanyl, sulfo, C₁₋₆-alkyl, C₃₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R¹²ⁱ is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl,

C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy.


In one embodiment, the invention is concerned with compounds of the general formula XI, wherein R⁶ⁱ, R⁷ⁱ, R⁸ⁱ and R⁹ⁱ are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XI, wherein R¹⁰ⁱ, R¹¹ⁱ, R¹²ⁱ, R¹³ⁱ and R¹⁴ⁱ are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XI, wherein R¹⁵ⁱ is hydrogen.

In another embodiment, the invention is concerned with compounds of the general formula XI, wherein R¹⁰ⁱ, R¹¹ⁱ, R¹²ⁱ, R¹³ⁱ, R¹⁴ⁱ and R¹⁵ⁱ are selected from the group consisting of H, F and methyl.

In an aspect of the invention, compounds are of the general formula XII



wherein R^{4j}, R^{5j}, R^{6j}, R^{7j}, R^{8j}, R^{9j} and R^{10j} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo,

amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{6j} and R^{7j}; and

wherein there may optionally be a covalent bond between R^{5j} and R^{6j}; and

R^{11j} and R^{12j} are independently selected from cyano, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

with the proviso that R^{9j} and R^{10j} are both hydrogen, there may optionally be a covalent bond connecting R^{11j} and R^{12j}.


In one embodiment, the invention is concerned with compounds of the general formula XII, wherein R^{4j}, R^{5j}, R^{6j} and R^{7j} are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-

alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XII, wherein at least one of the substituents R^{4j}, R^{5j}, R^{6j} and R^{7j} are different from hydrogen.

- 5 In another embodiment, the invention is concerned with compounds of the general formula XII, wherein R^{6j} is covalently bound to R^{5j}.

In an aspect of the invention, compounds are of the general formula XIII



10

- wherein R^{4k} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

- R^{5k} is selected from hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein

25

each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: with the proviso that when R^{4k} is hydrogen, then R^{6k} is not C(=O)N(Me)₂; and

R^{6k}, R^{7k}, R^{8k}, R^{9k}, R^{10k}, R^{11k}, R^{13k} and R^{14k} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{12k} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

wherein there may optionally be a covalent bond between any of the substituents selected from the group consisting of R^{4k} , R^{5k} , R^{7k} and R^{8k} ; and

- 5 wherein there may optionally be a covalent bond between R^{5k} and any one of the substituents R^{7k} and R^{8k} ; and

wherein there may optionally be one or more covalent bonds between R^{10k} , R^{11k} , R^{12k} , R^{13k} and R^{14k} .

10

In one embodiment, the invention is concerned with compounds of the general formula XIII, wherein the substituents R^{4k} and R^{5k} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XIII, wherein the substituents R^{5k} and R^{8k} are connected by a covalent bond.


- 15 In another embodiment, the invention is concerned with compounds of the general formula XIII, wherein the substituents R^{10k} and R^{11k} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XIII, wherein the substituents R^{12k} and R^{13k} are connected by a covalent bond.

- In another embodiment, the invention is concerned with compounds of the general formula
20 XIII, wherein R^{6k} , R^{7k} , R^{8k} and R^{9k} are selected from the group consisting of hydrogen, F, Cl, hydroxy, amino, methyl, methoxy, ethoxy and perhalomethyl.

- In another embodiment, the invention is concerned with compounds of the general formula XIII, wherein R^{10k} , R^{11k} , R^{12k} , R^{13k} and R^{14k} are selected from the group consisting of hydro-
25 gen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XIV



30

wherein R¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl or C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

- 5 R² is C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, or C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with
 10 one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, perhalomethyl and perhalomethoxy; with the proviso that when R¹ and R² are identical they are not methyl or benzyl; and

R² is optionally covalently bound to R¹ by an ether, thioether or C-C bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

- 20 R⁵ⁱ, R⁶ⁱ, R⁸ⁱ, R⁹ⁱ, R¹⁰ⁱ and R¹¹ⁱ are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally
 25 be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

- R⁴ⁱ and R⁷ⁱ are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, methoxy, C₃₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₃₋₆-

- alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and
- wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4l}, R^{5l}, R^{6l}, R^{7l}, R^{8l}, R^{9l}, R^{10l} and R^{11l}.

In one embodiment, the invention is concerned with compounds of the general formula XIV, wherein R¹ and R² are covalently bound so as to form a ring system with the N-atom to which they are bound.

In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein said ring system is a piperidine, piperazine, morpholine, or thiomorpholine.

In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein there are no covalent bonds between R¹ and R².


- In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein there are no covalent bonds between R^{8l} and any of the substituents selected from the group consisting of R^{9l}, R^{10l}, R^{11l}.

- In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein there are no covalent bonds between R^{8l} and any of the substituents selected from the group consisting of R^{4l}, R^{5l}, R^{6l} and R^{7l}.

In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein there are no covalent bonds between any of the substituents R^{4l}, R^{5l}, R^{6l}, R^{7l}, R^{8l}, R^{9l}, R^{10l} and R^{11l}.

- In another embodiment, the invention is concerned with compounds of the general formula XIV, wherein R^{4l}, R^{5l}, R^{6l} and R^{7l} are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XV



wherein R^{4m} is selected from hydrogen, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{5m} , R^{6m} , R^{7m} , R^{8m} , R^{9m} , R^{10m} , R^{11m} , R^{12m} , R^{13m} , R^{14m} , R^{15m} and R^{16m} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents

independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4m}, R^{5m}, R^{6m}, R^{7m} and R^{8m}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9m}, R^{10m}, R^{11m}, R^{12m}, R^{13m}, R^{14m}, R^{15m} and R^{16m}.

In one embodiment, the invention is concerned with compounds of the general formula XV, wherein there are no covalent bonds between any of the substituents selected from the group consisting of R^{5m}, R^{6m}, R^{7m} and R^{8m}.

In another embodiment, the invention is concerned with compounds of the general formula XV, wherein there are no covalent bonds between any of the substituents R^{9m}, R^{10m}, R^{11m}, R^{12m} and R^{13m}.


In another embodiment, the invention is concerned with compounds of the general formula XV, wherein there are no covalent bonds between any of the substituents R^{14m}, R^{15m} and R^{16m}.

In another embodiment, the invention is concerned with compounds of the general formula XV, wherein R^{4m} is covalently bound to R^{6m}.

In another embodiment, the invention is concerned with compounds of the general formula XV, wherein R^{5m}, R^{6m}, R^{7m}, R^{8m}, R^{9m}, R^{10m}, R^{11m}, R^{12m}, R^{13m} and R^{14m} are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XV, wherein R^{14m}, R^{15m} and R^{16m} are selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XVI



wherein R^{4n} is selected from hydrogen, sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

Z is selected from S, S(=O) and S(=O)₂; and

R^{5n} , R^{6n} , R^{7n} , R^{8n} , R^{9n} , R^{10n} , R^{11n} , R^{12n} , R^{13n} and R^{14n} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and

C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R⁴ⁿ, R⁵ⁿ, R⁶ⁿ, R⁷ⁿ and R⁸ⁿ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R¹⁰ⁿ, R¹¹ⁿ, R¹²ⁿ and R¹⁴ⁿ; and

at least one of the substituents R⁵ⁿ, R⁶ⁿ, R⁷ⁿ, R⁸ⁿ, R⁹ⁿ, R¹⁰ⁿ, R¹¹ⁿ, R¹²ⁿ, R¹³ⁿ and R¹⁴ⁿ are different from hydrogen.

In one embodiment, the invention is concerned with compounds of the general formula XVI,

wherein there are no covalent bonds between any of the substituents R⁵ⁿ, R⁶ⁿ, R⁷ⁿ and R⁸ⁿ.

In another embodiment, the invention is concerned with compounds of the general formula XVI, wherein there are no covalent bonds between any of the substituents R¹⁰ⁿ, R¹¹ⁿ, R¹²ⁿ.


In another embodiment, the invention is concerned with compounds of the general formula XVI, wherein R¹⁴ⁿ is not covalently bound to any other substituent selected from the group consisting of R⁴ⁿ, R⁵ⁿ, R⁶ⁿ, R⁷ⁿ, R⁸ⁿ, R¹⁰ⁿ, R¹¹ⁿ, R¹²ⁿ.

In another embodiment, the invention is concerned with compounds of the general formula XVI, wherein R⁴ⁿ is covalently bound to R⁶ⁿ.

In another embodiment, the invention is concerned with compounds of the general formula XVI, wherein R⁵ⁿ, R⁶ⁿ, R⁷ⁿ and R⁸ⁿ are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XVI, wherein R¹⁴ⁿ is selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XVII



5

wherein R^{20} is selected from sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{20} is not methyl; and

25 R^{40} , R^{50} , R^{60} and R^{70} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl,

30

C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

A^{1o} is N or C-R^{8o}; A^{2o} is N or C-R^{8o}; A^{3o} is N or C-R^{10o}; A^{4o} is N or C-R^{11o}; and A^{5o} is N or C-R^{12o}; and

wherein R^{8o}, R^{9o}, R^{10o}, R^{11o} and R^{12o} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, perhalomethyl and perhalomethoxy; with the proviso that

when A^{1o}, A^{2o}, A^{3o}, A^{4o} and A^{5o} are all CH, and R^{4o}, R^{5o}, R^{6o} and R^{7o} are all hydrogen, then R^{2o} is not phenyl; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4o}, R^{5o}, R^{6o}, R^{7o}, R^{8o}, R^{9o}, R^{10o}, R^{11o} and R^{12o}

30


In one embodiment, the invention is concerned with compounds of the general formula XVII, wherein there are no covalent bonds between any of the substituents R^{4o}, R^{5o}, R^{6o} and R^{7o}. In another embodiment, the invention is concerned with compounds of the general formula XVII, wherein there are no covalent bonds between any of the substituents A^{1o}, A^{2o}, A^{3o}, A^{4o} and A^{5o}.

35

In another embodiment, the invention is concerned with compounds of the general formula XVII, wherein R^{2o} is selected from the group consisting of cycloalkyl, phenyl, piperidine, piperazine, morpholine, thiomorpholine and heteroaryl.

In another embodiment, the invention is concerned with compounds of the general formula XVII, wherein R^{4o} , R^{5o} , R^{6o} and R^{7o} are independently selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XVIII



wherein R^{4p} is selected from hydrogen, sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl,

oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{5p}, R^{6p}, R^{7p} and R^{8p} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, F, amino, cyano, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

R^{9p}, R^{10p}, R^{11p}, R^{12p}, R^{13p}, R^{14p}, R^{15p} and R^{16p} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{5p}, R^{6p} and R^{7p}; and

wherein there may optionally be a covalent bond between R^{4p} and R^{5p} so as to form a chromen ring system; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9p} , R^{10p} , R^{11p} , R^{12p} , R^{13p} , R^{14p} , R^{15p} and R^{16p} ; and


E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{17p}- and -CR^{17p}R^{18p}-;
5 ; and

wherein R^{17p} and R^{18p} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally
10 be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may
15 optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independ-
20 lected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independ-
25 ently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy.

In one embodiment, the invention is concerned with compounds of the general formula XVIII, wherein E is selected from the group consisting of -O- and -CR^{17p}R^{18p}-.


30 In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{4p} and R^{5p} are connected by a covalent bond so as to form said chromen ring system.

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{4p} and R^{5p} are connected by a covalent bond so as to form the chromen ring
35 system



wherein R^{19p} is selected from the group consisting of hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy.

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{4p} is selected from the group consisting of



wherein R^{20p} , R^{21p} , R^{22p} , R^{23p} and R^{24p} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, perhalomethyl and perhalomethoxy.

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein there are no covalent bonds between any of the substituents R^{4p} , R^{5p} , R^{6p} , R^{7p} and R^{8p} .

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein there are no covalent bonds between any of the substituents R^{9p} , R^{10p} , R^{11p} , R^{12p} , R^{13p} , R^{14p} , R^{15p} and R^{16p} .


- 5 In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{10p} and R^{11p} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{9p} , R^{10p} , R^{11p} , R^{12p} , R^{13p} , R^{14p} , R^{15p} and R^{16p} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2NH_2$, $-NH-S(=O)_2-$
 10 OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XVIII, wherein R^{5p} , R^{6p} , R^{7p} and R^{8p} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

15

In an aspect of the invention, compounds are of the general formula XIX



- 20 wherein R^{4q} , R^{6q} , R^{7q} , R^{9q} , R^{10q} , R^{11q} , R^{12q} , R^{13q} , R^{14q} , R^{15q} , R^{16q} , R^{17q} , and R^{18q} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen,
 25 amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -
 30

heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and

R^{5q} and R^{8q} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, F, Br, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy: and


wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4q}, R^{5q}, R^{6q}, R^{7q} and R^{8q}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{8q}, R^{10q}, R^{11q}, R^{12q}, R^{13q}, R^{14q}, R^{15q}, R^{16q}, R^{17q} and R^{18q}.

In one embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{4q} and R^{6q} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{4q} is selected from the group consisting of substituted heteroaryl and substituted C_{3-8} -heterocyclyl.

- 5 In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{4q} is selected from the group consisting of



- wherein R^{19q} , R^{20q} , R^{21q} , R^{22q} and R^{23q} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy.

25

In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein there are no covalent bonds between any of the substituents R^{4q} , R^{5q} , R^{6q} , R^{7q} and R^{8q} .

- 30 In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein there are no covalent bonds between any of the substituents R^{9q} , R^{10q} , R^{11q} , R^{12q} , R^{13q} , R^{14q} , R^{15q} , R^{16q} , R^{17q} and R^{18q} .

In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{10p} and R^{11p} are connected by a covalent bond.


In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{11p} and R^{13p} are connected by a covalent bond.

5 In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{9q} , R^{10q} , R^{11q} , R^{12q} , R^{13q} , R^{14q} , R^{15q} , R^{16q} , R^{17q} and R^{18q} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{9q} , R^{10q} , R^{11q} , R^{12q} , R^{14q} , R^{15q} , R^{16q} , R^{17q} and R^{18q} are all hydrogen or F.

In another embodiment, the invention is concerned with compounds of the general formula XIX, wherein R^{5q} , R^{6q} , R^{7q} and R^{8q} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

15 In an aspect of the invention, compounds are of the general formula XX



20 wherein R^{4r} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, and C_{3-10} -cycloalkyl.

10-cycloalkyl, perhalomethyl and perhalomethoxy: provided that R^{4r} is not methyl or phenyl;
and

R^{5r} , R^{6r} , R^{7r} , R^{8r} , R^{9r} , R^{10r} , R^{11r} , R^{12r} , R^{13r} , R^{14r} , R^{15r} , R^{16r} , R^{17r} and R^{18r} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, F, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy: and


wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{5r} , R^{6r} , R^{7r} and R^{8r} ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9r} , R^{10r} , R^{11r} , R^{12r} , R^{13r} , R^{14r} , R^{15r} , R^{16r} , R^{17r} and R^{18r} .

In one embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{4r} and R^{5r} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{4r} is selected from the group consisting of substituted heteroaryl and substituted C_{3-8} -heterocyclyl.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{4r} is selected from the group consisting of



wherein R^{18r} , R^{20r} , R^{21r} , R^{22r} and R^{23r} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein there are no covalent bonds between any of the substituents R^{5r} , R^{6r} , R^{7r} and R^{8r} .

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein there are no covalent bonds between any of the substituents R^{9r} , R^{10r} , R^{11r} , R^{12r} , R^{13r} , R^{14r} , R^{15r} and R^{16r} .

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{10r} and R^{11r} are connected by a covalent bond.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{11r} and R^{13r} are connected by a covalent bond.


In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{9r} , R^{10r} , R^{11r} , R^{12r} , R^{13r} , R^{14r} , R^{15r} , R^{16r} , R^{17r} and R^{18r} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{9r} , R^{10r} , R^{11r} , R^{12r} , R^{14r} , R^{15r} , R^{16r} , R^{17r} and R^{18r} are all hydrogen or F.

In another embodiment, the invention is concerned with compounds of the general formula XX, wherein R^{5r} , R^{6r} , R^{7r} and R^{8r} are independently selected from hydrogen, F, Cl, C_{1-6} -alkyl,

C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In an aspect of the invention, compounds are of the general formula XXI



wherein A^{1s} is N or C-R^{17s}; A^{2s} is N or C-R^{18s}; A^{3s} is N or C-R^{19s}; A^{4s} is N or C-R^{20s}; and A^{5s} is N or C-R^{21s}; and

10

wherein R^{17s}, R^{18s}, R^{19s}, R^{20s} and R^{21s} are independently selected from hydrogen, hydroxy, sulfanyl, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

15

20

25

62

R^{5s}, R^{6s}, R^{7s} and R^{8s} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, F, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

R^{9s} , R^{10s} , R^{11s} , R^{12s} , R^{13s} , R^{14s} , R^{15s} and R^{16s} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, F, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{22s}- and -CR^{22s}R^{23s}-; and

wherein R^{22s} and R^{23s} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋

₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

5

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{5s}, R^{6s}, R^{7s} and R^{8s}; and

10

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9s}, R^{10s}, R^{11s}, R^{12s}, R^{13s}, R^{14s}, R^{15s}, R^{16s}, R^{18s}, R^{19s}, R^{20s}, R^{21s}, R^{22s} and R^{23s}.

In one embodiment, the invention is concerned with compounds of the general formula XXI, wherein E is selected from the group consisting of -O- and -CR^{22s}R^{23s}-.

15

In another embodiment, the invention is concerned with compounds of the general formula XXI, wherein there are no covalent bonds between any of the substituents R^{5s}, R^{6s}, R^{7s} and R^{8s}.

20

In another embodiment, the invention is concerned with compounds of the general formula XXI, wherein R^{5s}, R^{6s}, R^{7s} and R^{8s} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXI, wherein there is a covalent bond between R^{10s} and R^{11s}.


25

In another embodiment, the invention is concerned with compounds of the general formula XXI, wherein there is a covalent bond between R^{13s} and R^{22s}.

In another embodiment, the invention is concerned with compounds of the general formula XXI, wherein R^{9s}, R^{10s}, R^{11s}, R^{12s}, R^{13s}, R^{14s}, R^{15s} and R^{16s} are all hydrogen or F.

In an aspect of the invention, compounds are of the general formula XXII

30



wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{18t}- and -CR^{18t}R^{19t}-; and

- 5
- R^{4t}, R^{5t}, R^{6t}, R^{7t}, R^{8t}, R^{9t}, R^{10t}, R^{11t}, R^{12t}, R^{13t}, R^{14t}, R^{15t}, R^{16t}, R^{17t}, R^{18t} and R^{19t} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{4t} and R^{5t} are not both methyl; and
- 20
- 25

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9t}, R^{10t}, R^{11t}, R^{12t}, R^{13t}, R^{14t}, R^{15t}, R^{16t}, R^{17t}, R^{18t} and R^{19t}.

In one embodiment, the invention is concerned with compounds of the general formula XXII, wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂- and -CR^{18t}R^{19t}.

In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein R^{18t} is hydrogen or F.

- 5 In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein R^{10t}, R^{12t}, R^{4t} and R^{8t} are all hydrogen.

In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein R^{6t}, R^{7t}, R^{8t} and R^{9t} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-


- 10 OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein there is a covalent bond between R^{11t} and R^{13t}.

In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein there is a covalent bond between R^{13t} and R^{19t}.

- 15 In another embodiment, the invention is concerned with compounds of the general formula XXII, wherein R^{10t}, R^{11t}, R^{12t}, R^{13t}, R^{14t}, R^{15t}, R^{16t} and R^{17t} are all hydrogen or F.

In an aspect of the invention, compounds are of the general formula XXIII



20

wherein R^{4u}, R^{5u}, R^{6u}, R^{7u}, R^{8u}, R^{9u}, R^{10u}, R^{11u}, R^{12u}, R^{13u}, R^{14u} and R^{15u} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of

- 25 hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more
- 30

- substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted
- 5 with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{4u}, R^{5u} and R^{6u} are not all hydrogen; and
- 10 E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{16u}- and -CR^{16u}R^{17u}-;

- wherein R^{16u} and R^{17u} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and
- 15
- 20
- 25

- 30 wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4u}, R^{5u} and R^{6u}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{8u}, R^{9u}, R^{10u} and R^{11u}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{12u} , R^{13u} , R^{14u} , R^{15u} , R^{16u} and R^{17u} .

In one embodiment, the invention is concerned with compounds of the general formula XXIII, wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂- and -CR^{18u}R^{17u}- . In another embodiment, the invention is concerned with compounds of the general formula XXIII, wherein R^{17u} is hydrogen or F.


In another embodiment, the invention is concerned with compounds of the general formula XXIII, wherein there are no covalent bonds between any of the substituents R^{8u}, R^{9u}, R^{10u} and R^{11u}.

In another embodiment, the invention is concerned with compounds of the general formula XXIII, wherein R^{8u}, R^{9u}, R^{10u} and R^{11u} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXIII, wherein there is a covalent bond between R^{12u} and R^{13u}.

In another embodiment, the invention is concerned with compounds of the general formula XXIII, wherein R^{12u}, R^{13u}, R^{14u} and R^{15u} are all selected from the group consisting of hydrogen, F, methyl and C₁₋₆-alkyl.

In an aspect of the invention, compounds are of the general formula XXIVa-b



wherein R^{4v}, R^{5v}, R^{6v}, R^{7v}, R^{8v}, R^{9v}, R^{10v}, R^{11v}, R^{12v}, R^{13v}, R^{14v}, R^{15v}, R^{16v}, R^{17v} and R^{18v} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro,

C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that none of the substituents R^{4v}, R^{5v} and R^{6v} are benzothiazolyl or benzooxazolyl; and

E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{19v}- and -CR^{18v}R^{20v}-; and

wherein R^{19v} and R^{20v} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{7v} and R^{8v} or between the substituents R^{9v} and R^{10v}; and

wherein there may optionally be a covalent bond between R^{6v} and a substituent selected from R^{8v} and R^{9v} ; and

wherein there may be a covalent bond between any of the substituents selected from the
5 group consisting of R^{4v} , R^{5v} and R^{6v} ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{11v} , R^{12v} , R^{13v} , R^{14v} , R^{15v} , R^{16v} , R^{17v} , R^{18v} , R^{19v} and R^{20v} .

10 In one embodiment, the invention is concerned with compounds of the general formula XXIV, wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂- and -CR^{19v}R^{20v}-.

In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein R^{20v} is hydrogen or F.

15 In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein there are no covalent bonds between any of the substituents R^{4v} , R^{5v} and R^{6v} .
In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein there are no covalent bonds between any of the substituents R^{7v} , R^{8v} , R^{9v} and R^{10v} .


20 In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein there are no covalent bonds between a substituent selected from the group consisting of R^{4v} , R^{5v} and R^{6v} and a substituent selected from the group consisting of R^{7v} , R^{8v} , R^{9v} and R^{10v} .

In another embodiment, the invention is concerned with compounds of the general formula
25 XXIV, wherein R^{7v} , R^{8v} , R^{9v} and R^{10v} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein there is a covalent bond between R^{12v} and R^{13v} .

30 In another embodiment, the invention is concerned with compounds of the general formula XXIV, wherein R^{11v} , R^{12v} , R^{13v} , R^{14v} , R^{15v} , R^{16v} , R^{17v} and R^{18v} are all hydrogen or F.

In an aspect of the invention, compounds are of the general formula XXV



wherein R^{4w} is selected from hydrogen, hydroxy, amino, sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, amino, sulfo, C_{2-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy: with the proviso that R^{4w} is not methyl, morpholine or a 2-chromen derivative; and

Z is selected from S, $S(=O)$ and $S(=O)_2$; and

R^{5w} , R^{6w} , R^{7w} , R^{8w} , R^{9w} , R^{10w} , R^{11w} , R^{12w} , R^{13w} , R^{14w} , R^{15w} and R^{16w} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents

independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

- 5 E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{17w}- and -CR^{17w}R^{18w}-; and

wherein R^{17w} and R^{18w} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

- 20 wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{4w}, R^{5w}, R^{6w}, R^{7w} and R^{8w}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9w}, R^{10w}, R^{11w}, R^{12w}, R^{13w}, R^{14w}, R^{15w}, R^{16w}, R^{17w} and R^{18w}.

In one embodiment, the invention is concerned with compounds of the general formula XXV, wherein the only substituents which are covalently bound are R^{4w} and R^{5w}.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein there are no covalent bonds between any of the substituents R^{5w}, R^{6w}, R^{7w} and R^{8w}.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein there are no covalent bonds between any of the substituents R^{9w}, R^{10w}, R^{11w}, R^{12w}, R^{13w}, R^{14w}, R^{15w} and R^{16w}.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂- and -CR^{17w}R^{18w}-.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein R^{18w} is hydrogen or F.


In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein R^{5w}, R^{6w}, R^{7w} and R^{8w} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein there is a covalent bond between R^{10w} and R^{11w}.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein there is a covalent bond between R^{13w} and R^{17w}.

In another embodiment, the invention is concerned with compounds of the general formula XXV, wherein R^{9w}, R^{10w}, R^{11w}, R^{12w}, R^{13w}, R^{14w}, R^{15w} and R^{16w} are all hydrogen or F.

In an aspect of the invention, compounds are of the general formula XXVI



20

wherein R^{4x} is selected from imino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of imino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein

30

- each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein
- 5 each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and
- 10 R^{5x}, R^{6x}, R^{7x}, R^{8x}, R^{9x}, R^{10x}, R^{11x}, R^{12x} and R^{13x} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and
- 25

E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{14x}- and -CR^{15x}R^{16x}-;

- 30 wherein R^{14x} is selected from hydrogen, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl,
- 35 wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl,

- heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and
- 10 R^{16x} and R^{16x} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and
- 15 wherein there may optionally be a covalent bond between the substituents R^{4x} and R^{5x}; and
- wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{6x}, R^{7x}, R^{8x} and R^{9x}; and
- 25 wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{10x}, R^{11x}, R^{12x}, R^{13x}, R^{14x}, R^{15x} and R^{16x}.
- 30 In one embodiment, the invention is concerned with compounds of the general formula XXVI, wherein R^{6x}, R^{7x}, R^{8x} and R^{9x} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.
- In another embodiment, the invention is concerned with compounds of the general formula XXVI, wherein R^{10x}, R^{11x}, R^{12x} and R^{13x} are independently selected from the group consisting
- 35

of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.


In another embodiment, the invention is concerned with compounds of the general formula XXVI, wherein there is a covalent bond between the substituents R^{6x} and R^{7x}.

- 5 In another embodiment, the invention is concerned with compounds of the general formula XXVI, wherein there is a covalent bond between the substituents R^{10x} and R^{11x}.

In another embodiment, the invention is concerned with compounds of the general formula XXVI, wherein there is a covalent bond between R^{10x} and a substituent selected from R^{14x} and R^{15x}.

10

In an aspect of the invention, compounds are of the general formula XXVII



- 15 wherein R^{4y} and R^{11y} are independently selected from imino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of imino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen,
- 20
- 25
- 30

amino, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

R^{6y}, R^{6y}, R^{7y}, R^{8y}, R^{8y} and R^{10y} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{4y} and R^{5y}; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{6y}, R^{7y}, R^{8y} and R^{9y}; and

wherein there may optionally be a covalent bond between the substituents R^{10y} and R^{11y}.


In one embodiment, the invention is concerned with compounds of the general formula XXVII, wherein R^{6y}, R^{7y}, R^{8y} and R^{9y} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXVII, wherein there are no covalent bonds between any of the substituents R^{6y} , R^{7y} , R^{8y} and R^{9y} .

In another embodiment, the invention is concerned with compounds of the general formula XXVII, wherein there is a covalent bond between the substituents R^{4y} and R^{5y} .

In another embodiment, the invention is concerned with compounds of the general formula XXVII, wherein there is a covalent bond between the substituents R^{10y} and R^{11y} .

In an aspect of the invention, compounds are of the general formula XXVIII



wherein R^{4z} , R^{8z} , R^{9z} , R^{10z} , R^{11z} , R^{12z} , R^{13z} , R^{14z} , R^{15z} , R^{16z} and R^{17z} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

- R^{5z} is a carbon bound substituent selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and
- R^{6z} and R^{7z} are independently selected from hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂-, -NR^{18z}- and -CR^{19z}R^{20z}-;

wherein R^{18z} is selected from hydrogen, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

R^{19z} and R^{20z} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{4z} and R^{5z} ; and wherein there may optionally be a covalent bond between any of the substituents R^{7z} , R^{8z} , and R^{9z} ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{10z} , R^{11z} , R^{12z} , R^{13z} , R^{14z} , R^{15z} , R^{16z} , R^{17z} , R^{18z} , R^{19z} , and R^{20z} .

In one embodiment, the invention is concerned with compounds of the general formula XXVIII, wherein there is a covalent bond between the substituents R^{4z} and R^{5z} .


In another embodiment, the invention is concerned with compounds of the general formula

- 5 XXVIII, wherein R^{6z} , R^{7z} , R^{8z} and R^{9z} are independently selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXVIII, wherein there is a covalent bond between R^{13z} and a substituent selected from R^{18z}

- 10 and R^{19z} .

In an aspect of the invention, compounds are of the general formula XXIX



15

wherein R^{4aa} is selected from hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

30

R^{5aa} , R^{6aa} , R^{7aa} , R^{8aa} , R^{9aa} , R^{10aa} , R^{11aa} , R^{12aa} and R^{13aa} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{5aa} and R^{6aa} or between the substituent R^{7aa} and R^{8aa} ; and

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9aa} , R^{10aa} , R^{11aa} and R^{12aa} ; and


E is selected from the group consisting of -O-, -S-, -S(=O)- and -S(=O)₂-.

In one embodiment, the invention is concerned with compounds of the general formula XXIX, wherein there is a covalent bond between the substituents R^{9aa} and R^{10aa} .

In another embodiment, the invention is concerned with compounds of the general formula XXIX, wherein R^{5aa} , R^{6aa} , R^{7aa} and R^{8aa} are independently selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXIX, wherein at least one of the substituents R^{5aa} , R^{6aa} , R^{7aa} and R^{8aa} are different from hydrogen.

In an aspect of the invention, compounds are of the general formula XXX



- 5 wherein R^{4ab} is selected from sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

E is selected from the group consisting of -NR^{13ab}- and -CR^{14ab}R^{15ab}-; and

- 20 R^{6ab}, R^{6ab}, R^{7ab}, R^{8ab}, R^{9ab}, R^{10ab}, R^{11ab}, R^{12ab}, R^{13ab}, R^{14ab} and R^{15ab} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-

alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{5ab} and R^{6ab} or between the substituent R^{7ab} and R^{8ab}; and

10

wherein there may optionally be one or more covalent bonds between any of the substituents selected from the group consisting of R^{9ab}, R^{10ab}, R^{11ab}, R^{12ab}, R^{13ab}, R^{14ab} and R^{15ab}.

In one embodiment, the invention is concerned with compounds of the general formula XXX, wherein there is a covalent bond between the substituents R^{9ab} and R^{10ab}.

15

In another embodiment, the invention is concerned with compounds of the general formula XXX, wherein there is a covalent bond between R^{10ab} and a substituent selected from R^{13ab} and R^{14ab}.

In another embodiment, the invention is concerned with compounds of the general formula XXX, wherein R^{5ab}, R^{6ab}, R^{7ab} and R^{8ab} are independently selected from the group consisting of hydrogen, F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.


20

In another embodiment, the invention is concerned with compounds of the general formula XXX, wherein at least one of the substituents R^{5ab}, R^{6ab}, R^{7ab} and R^{8ab} are different from hydrogen.

25

In another embodiment, the invention is concerned with compounds of the general formula XXX, wherein at least one of the substituents R^{9ab}, R^{10ab}, R^{11ab}, R^{12ab}, R^{13ab}, R^{14ab} and R^{15ab} are different from hydrogen.

30 In an aspect of the invention, compounds are of the general formula XXXI



wherein R^{4ac} is a carbon bound substituent selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

A^{1ac} is N or $C-R^{9ac}$; A^{3ac} is N or $C-R^{10ac}$; and A^{5ac} is N or $C-R^{11ac}$; and

R^{5ac} , R^{6ac} , R^{7ac} , R^{8ac} , R^{9ac} , R^{10ac} and R^{11ac} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -

alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that A^{3ac} is not C-C(OH)(CF₃)₂; and

A^{2ac} is N or C-R^{12ac}; and A^{4ac} is N or C-R^{13ac}; wherein

10

R^{12ac} and R^{13ac} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between R^{4ac} and a substituent selected from the group consisting of R^{5ac} and R^{6ac}; with the proviso that A^{1ac}, A^{2ac}, A^{3ac}, A^{4ac} and A^{5ac} are not all CH; and

30

wherein there may optionally be a covalent bond between any of the substituents R^{6ac}, R^{7ac} and R^{8ac}; and

wherein there may optionally be one or more covalent bonds between any of the substituents in A^{1ac}, A^{2ac}, A^{3ac}, A^{4ac} and A^{5ac}.

35

In one embodiment, the invention is concerned with compounds of the general formula XXXI, wherein there is a covalent bond between the substituents R^{4ac} and R^{6ac} .

In another embodiment, the invention is concerned with compounds of the general formula

5 XXXI, wherein there is a covalent bond between A^{3ac} and A^{4ac} .

In another embodiment, the invention is concerned with compounds of the general formula

XXXI, wherein there is a covalent bond between A^{3ac} and A^{2ac} .

In another embodiment, the invention is concerned with compounds of the general formula


10 XXXI, wherein R^{5ac} , R^{6ac} , R^{7ac} and R^{8ac} are independently selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula

XXXI, wherein at least one of the substituents A^{1ac} , A^{2ac} , A^{3ac} , A^{4ac} and A^{5ac} are different from CH.

15

In an aspect of the invention, compounds are of the general formula XXXIIa-b



20 wherein R^{4ad} is selected from hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, F, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{1-6} -alkoxy, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently se-

25

30

lected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

- 5 R^{5ad}, R^{6ad}, R^{7ad}, R^{8ad} and R^{9ad} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, 10 oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, 15 heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and 20

- R^{10ad} are selected from imino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein 25 each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₁₋₆-alkoxy, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and 30

wherein there may optionally be a covalent bond between R^{4ad} and a substituent selected from R^{6ad} and R^{7ad} ; and

- 5 wherein there may optionally be a covalent bond between the substituents R^{9ad} and R^{10ad} so as to form a 5-membered ring system.

In one embodiment, the invention is concerned with compounds of the general formula XXXIIa-b, wherein the substituents R^{9ad} and R^{10ad} are covalently bound so as to form a 5-membered ring system.

- 10 In another embodiment, the invention is concerned with compounds of the general formula XXXIIa-b, wherein said 5-membered ring system comprises at least one nitrogen atom in the ring.


In another embodiment, the invention is concerned with compounds of the general formula XXXIIa-b, wherein said 5-membered ring system contains 5 carbons in the ring.

- 15 In another embodiment, the invention is concerned with compounds of the general formula XXXIIa-b, wherein there is a covalent bond between the substituents R^{4ad} and R^{7ad} .

In another embodiment, the invention is concerned with compounds of the general formula XXXIIa-b, wherein R^{5ad} , R^{6ad} , R^{7ad} and R^{8ad} are independently selected from the group consisting of hydrogen, F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2NH_2$, -

- 20 $NH-S(=O)_2OH$, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXIIIa-b



25

wherein E is selected from the group consisting of $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NR^{16ae}-$ and $-CR^{17ae}R^{18ae}-$; and

R^{4ae} , R^{5ae} , R^{6ae} , R^{7ae} , R^{8ae} , R^{9ae} , R^{10ae} , R^{11ae} , R^{12ae} , R^{13ae} , R^{14ae} , R^{15ae} , R^{16ae} , R^{17ae} and R^{18ae} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between any of the substituents selected from R^{4ae} , R^{5ae} , R^{6ae} and R^{7ae} ; with the proviso that when R^{5ae} and R^{6ae} are covalently bound they do not form a isoquinolin-3-yl or a substituted isoquinolin-3-yl together with the pyridine which they are both bound to; and

wherein there may optionally be a covalent bond between any of the substituents selected from R^{8ae} , R^{9ae} , R^{10ae} , R^{11ae} , R^{12ae} , R^{13ae} , R^{14ae} , R^{15ae} , R^{16ae} , R^{17ae} and R^{18ae} .

In one embodiment, the invention is concerned with compounds of the general formula XXXIIIa-b, wherein there is a covalent bond between R^{6ae} and R^{7ae} .


In another embodiment, the invention is concerned with compounds of the general formula XXXIIIa-b, wherein there is a covalent bond between R^{5ae} and R^{6ae} .

In another embodiment, the invention is concerned with compounds of the general formula XXXIIIa-b, wherein at least one of R^{8ae} , R^{9ae} , R^{10ae} , R^{11ae} , R^{12ae} , R^{13ae} , R^{14ae} and R^{15ae} is se-

lected from F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXIV

5



wherein R^{4af}, R^{5af}, R^{6af}, R^{7af}, R^{8af}, R^{9af}, R^{10af}, R^{11af}, R^{12af}, R^{13af}, R^{14af} and R^{15af} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl; and

wherein there may optionally be a covalent bond between R^{6af} and any of the substituents selected from R^{5af} and R^{7af}; and

wherein there may optionally be a covalent bond between the substituents R^{9af} and R^{10af} ; and

R^{16af} is selected from hydrogen, hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{16af} is not methyl; and


wherein there may optionally be a covalent bond between the substituents R^{12af} and R^{16af} .

In one embodiment, the invention is concerned with compounds of the general formula XXXIV, wherein there is a covalent bond between the substituents R^{6af} and R^{6af} .

In another embodiment, the invention is concerned with compounds of the general formula XXXIV, wherein there is a covalent bond between the substituents R^{9af} and R^{10af} .

In another embodiment, the invention is concerned with compounds of the general formula XXXIV, wherein at least one of R^{8af} , R^{9af} , R^{10af} , R^{11af} , R^{12af} , R^{13af} , R^{14af} and R^{15af} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXV



wherein E is selected from the group consisting of $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$ and $-CR^{16ag}R^{17ag}-$; and

5

R^{4ag} , R^{5ag} , R^{6ag} , R^{10ag} , R^{11ag} , R^{12ag} , R^{13ag} , R^{16ag} and R^{17ag} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{4ag} is not $-CH_2-N(CH_3)_2$; and

25

wherein there may optionally be a covalent bond between the substituents R^{5ag} and R^{6ag} ; and

R^{7ag} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sul-

fanyl, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy;

R^{8ag}, R^{9ag}, R^{14ag} and R^{15ag} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-

alkyl, C₂₋₆-alkenyl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{9ag} and R^{10ag};

5 and


wherein there may optionally be a covalent bond between the substituents R^{12ag} and R^{16ag}.

In one embodiment, the invention is concerned with compounds of the general formula XXXV, wherein there is a covalent bond between the substituents R^{5ag} and R^{6ag}.

In another embodiment, the invention is concerned with compounds of the general formula XXXV, wherein there is a covalent bond between the substituents R^{9ag} and R^{10ag}.

In another embodiment, the invention is concerned with compounds of the general formula XXXV, wherein at least one of R^{8ag}, R^{9ag}, R^{10ag}, R^{11ag}, R^{12ag}, R^{13ag}, R^{14ag} and R^{15ag} are selected from F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXVIa-b



20

wherein R^{4ah}, R^{5ah}, R^{6ah}, R^{7ah}, R^{8ah} and R^{9ah} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted

25

with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{5ah} and R^{6ah} are not both CF₃; and

wherein there may optionally be a covalent bond between any of the substituents selected from R^{4ah}, R^{5ah}, R^{6ah} and R^{7ah}; and

wherein there may optionally be a covalent bond between the substituents R^{8ah} and R^{9ah}; with the proviso that when R^{8ah} and R^{9ah} together with the carbon to which they are bound forms phenyl then R^{5ah}, R^{6ah}, R^{7ah} and R^{8ah} are not all hydrogen.


In one embodiment, the invention is concerned with compounds of the general formula XXXVIa-b, wherein there is a covalent bond between R^{8ah} and R^{9ah}.

In another embodiment, the invention is concerned with compounds of the general formula XXXVIa-b, wherein there is a covalent bond between R^{6ah} and R^{7ah}.

In another embodiment, the invention is concerned with compounds of the general formula XXXVIa-b, wherein there is a covalent bond between R^{5ah} and R^{6ah}.

In another embodiment, the invention is concerned with compounds of the general formula XXXVIa-b, wherein at least one of R^{5ah}, R^{6ah}, R^{7ah} and R^{8ah} are selected from F, Cl, C₁₋₆-alkyl, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NH-S(=O)₂-OH, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXVII



R^{4ai} , R^{5ai} , R^{6ai} , R^{7ai} , R^{8ai} and R^{9ai} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{4ai} is not $CH_2-N(Me)_2$; and

wherein there may optionally be a covalent bond between any of the substituents selected from R^{4ai} , R^{5ai} , R^{6ai} and R^{7ai} , and

wherein there may optionally be a covalent bond between the substituents R^{8ai} and R^{9ai} , with the proviso that R^{8ai} and R^{9ai} together with the carbon to which they are bound do not form 4-methoxy-phenyl, 4-chloro-phenyl or 4-nitro-phenyl.


In one embodiment, the invention is concerned with compounds of the general formula XXXVII, wherein there is a covalent bond between R^{8ai} and R^{9ai} .

In another embodiment, the invention is concerned with compounds of the general formula XXXVII, wherein there is a covalent bond between R^{6ai} and R^{7ai} .

- 5 In another embodiment, the invention is concerned with compounds of the general formula XXXVII, wherein there is a covalent bond between R^{5ai} and R^{6ai} .

In another embodiment, the invention is concerned with compounds of the general formula XXXVII, wherein at least one of R^{5ai} , R^{6ai} , R^{7ai} and R^{8ai} are selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and per-halomethyl.

In another aspect of the invention, compounds are of the general formula XXXVIII



15

wherein R^{4aj} , R^{5aj} , R^{6aj} and R^{7aj} are independently selected from amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino,

20

25

30

cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{4aj} and R^{5aj}; and

5

wherein there may optionally be a covalent bond between the substituents R^{6aj} and R^{7aj}.


In one embodiment, the invention is concerned with compounds of the general formula XXXVIII, wherein there is a covalent bond between R^{6aj} and R^{7aj}.

- 10 In another embodiment, the invention is concerned with compounds of the general formula XXXVIII, wherein R^{6aj}, R^{7aj} and said covalent bond are selected so as to form a ring system selected from an optionally substituted piperidine, piperazine, morpholine and thiomorpholine.

- In another embodiment, the invention is concerned with compounds of the general formula
15 XXXVIII, wherein said ring system is selected from piperidine, piperazine, morpholine and thiomorpholine

In another embodiment, the invention is concerned with compounds of the general formula XXXVIII, wherein there is a covalent bond between R^{4aj} and R^{5aj}.

- 20 In another aspect of the invention, compounds are of the general formula XXXIX



- wherein R^{4ak}, R^{5ak}, R^{6ak} and R^{7ak} are independently selected from amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo,
- 25
- 30

halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that none of R^{4ak} and R^{5ak} are -C(=O)-O-isopropyl; and

wherein there may optionally be a covalent bond between the substituents R^{4ak} and R^{5ak}, with the proviso that R^{4ak}, R^{5ak} and the nitrogen to which they are bound do not form a substituted tetrazolyl moiety; and

wherein there may optionally be a covalent bond between the substituents R^{6ak} and R^{7ak};


In one embodiment, the invention is concerned with compounds of the general formula XXXIX, wherein there is a covalent bond between R^{6ak} and R^{7ak}.

In another embodiment, the invention is concerned with compounds of the general formula XXXIX, wherein R^{6ak}, R^{7ak} and said covalent bond are selected so as to form a ring system selected from an optionally substituted piperidine, piperazine, morpholine and thiomorpholine.

In another embodiment, the invention is concerned with compounds of the general formula XXXIX, wherein said ring system is selected from piperidine, piperazine, morpholine and thiomorpholine

In another embodiment, the invention is concerned with compounds of the general formula XXXIX, wherein there is a covalent bond between R^{4ak} and R^{5ak}.

In another aspect of the invention, compounds are of the general formula XXXX



wherein R^{4al} and R^{5al} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, C_{3-10} -cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between R^{4al} and R^{5al} ; with the proviso that R^{4al} and R^{5al} together with the nitrogen to which they are bound do not form 2,5-pyrrolidinedione or an annealed ring system comprising 3 or more rings; and

wherein E is selected from the group consisting of -O-, -S-, -S(=O)-, -S(=O)₂- and -NR^{14al}-, -CR^{15al}R^{16al}-; and

R^{6al} , R^{7al} , R^{8al} , R^{9al} , R^{10al} , R^{11al} , R^{12al} , R^{13al} , R^{14al} , R^{15al} and R^{16al} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl,

aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, het-

5 eroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; and

wherein there may optionally be a covalent bond between the substituents R^{7al} and R^{8al}; and

25 wherein there may optionally be a covalent bond between the substituents R^{10al} and a substituent selected from R^{14al} and R^{15al} .

In one embodiment, the invention is concerned with compounds of the general formula XXXX, wherein there is a covalent bond between the substituents R^{4al} and R^{5al}.

30 In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein there is a covalent bond between the substituents R^{7al} and R^{8al}.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein there is a covalent bond between R^{10al} and a substituent selected from R^{14al} or R^{15al}.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein E is $-NR^{14al}$.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein R^{14al} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.


In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein E is $-CR^{15al}R^{16al}$.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein R^{16al} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein R^{15al} is hydrogen.

In another embodiment, the invention is concerned with compounds of the general formula XXXX, wherein at least one of R^{6al} , R^{7al} , R^{8al} , R^{9al} , R^{10al} , R^{11al} , R^{12al} and R^{13al} are selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another aspect of the invention, compounds are of the general formula XXXXI



wherein E is selected from the group consisting of $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$ and $-NR^{14am}$, $-CR^{15am}R^{16am}$; and

wherein A^{1am} is N or $C-R^{17am}$; A^{2am} is N or $C-R^{18am}$; A^{3am} is N or $C-R^{19am}$; A^{4am} is N or $C-R^{20am}$; and

R^{6am} , R^{7am} , R^{8am} , R^{9am} , R^{10am} , R^{11am} , R^{12am} , R^{13am} , R^{14am} , R^{15am} , R^{16am} , R^{17am} , R^{18am} , R^{19am} and R^{20am} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl,

wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the
5 hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-
10 cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hy-
15 droxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy,
20 sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{17am} and R^{20am} are not both hydroxy; and

wherein there may optionally be a covalent bond between R^{18am} and a substituent selected from R^{17am} and R^{19am}; and

wherein there may optionally be a covalent bond between the substituents R^{10am} and a substituent selected from R^{14am} and R^{15am}; and

wherein there may optionally be a covalent bond between the substituents R^{7am} and R^{8am} .

In one embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein there is a covalent bond between R^{18am} and a substituents selected from R^{17am} and R^{19am}.

In another embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein there is a covalent bond between R^{10am} and a substituent selected from R^{14am} or R^{15am} .

In another embodiment, the invention is concerned with compounds of the general formula
5 XXXXI, wherein E is $-NR^{14am}-$.

In another embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein R^{14am} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula
10 XXXXI, wherein E is $-CR^{15am}R^{16am}-$.


In another embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein R^{16am} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula
15 XXXXI, wherein R^{15am} is hydrogen.

In another embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein there is a covalent bond between R^{15am} and R^{16am} .

In another embodiment, the invention is concerned with compounds of the general formula XXXXI, wherein at least one of R^{6am} , R^{7am} , R^{8am} , R^{9am} , R^{10am} , R^{11am} , R^{12am} and R^{13am} are selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.
20

In another aspect of the invention, compounds are of the general formula XXXXII



25

wherein E is selected from the group consisting of $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$ and $-NR^{17an}-$, $-CR^{18an}R^{19an}-$; and

R^{4an}, R^{5an}, R^{6an}, R^{7an}, R^{8an}, R^{9an}, R^{10an}, R^{11an}, R^{12an}, R^{13an}, R^{14an}, R^{15an}, R^{16an}, R^{17an}, R^{18an} and
R^{19an} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino,
cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl,
wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-
heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents
independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-
alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of the
hydroxy, sulfanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and
C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently
selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-
alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sul-
fanyl, amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-
cycloalkyl may optionally be substituted with one or more substituents independently se-
lected from hydroxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl,
aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl,
amino, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl
may optionally be substituted with one or more substituents independently selected from hy-
droxy, sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, het-
eroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of the hydroxy, sulfanyl, amino,
sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl may op-
tionally be substituted with one or more substituents independently selected from hydroxy,
sulfanyl, oxo, halogen, amino, cyano, nitro, sulfo, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-
heterocyclyl, C₃₋₁₀-cycloalkyl, perhalomethyl and perhalomethoxy; with the proviso that R^{4an} is
not methyl or phenyl; and

25

wherein there may optionally be a covalent bond between R^{13an} and a substituent selected from R^{17an} and R^{18an}; and

wherein there may optionally be a covalent bond between the substituents R^{7an} and R^{8an} .

30

In one embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein R^{4an} is a substituted heteroaryl.

In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein said substituted heteroaryl is a substituted pyridine.

In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein there is a covalent bond between the substituents R^{7an} and R^{8an} .

In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein there is a covalent bond between R^{13an} and a substituents selected from

5 R^{17an} and R^{18an} .

In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein E is $-NR^{17an}-$.


In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein E is $-CR^{18an}R^{19an}-$.

10 In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein R^{19an} is selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein R^{18an} is hydrogen.

15 In another embodiment, the invention is concerned with compounds of the general formula XXXXII, wherein at least one of R^{5an} , R^{6an} , R^{7an} , R^{8an} , R^{9an} , R^{10an} , R^{11an} , R^{12an} , R^{13an} , R^{14an} , R^{15an} and R^{16an} are selected from F, Cl, C_{1-6} -alkyl, C_{1-6} -alkoxy, $-C(=O)NH_2$, $-NHC(=O)-OH$, $-S(=O)_2-NH_2$, $-NH-S(=O)_2-OH$, hydroxy, amino and perhalomethyl.

20 In another aspect of the invention, compounds are of the general formula XXXXIII



wherein R^{4ao} is selected from hydrogen, hydroxy, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of

25 hydroxy, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl
30 may optionally be substituted with one or more substituents independently selected from hy-

- droxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy,
- 5 sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋
- 10 ₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and
- 15 wherein R^{5ao}, R^{6ao}, R^{7ao}, R^{8ao}, R^{9ao}, R^{10ao}, R^{11ao} and R^{12ao} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy,
- 20 sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋
- 25 ₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl,
- 30 wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋
- ₁₀-cycloalkyl may optionally be substituted with one or more substituents independently se-

lected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy;

with the proviso that any one of R^{6ao} and R^{8ao} is not methyl; and with the further proviso that R^{4ao}, R^{5ao}, R^{6ao}, R^{7ao} and R^{8ao} are not all hydrogen; and

5

wherein there may optionally be a covalent bond between R^{4ao} and R^{6ao}; and

wherein there may optionally be a covalent bond between R^{7ao} and R^{8ao}; and


wherein there may optionally be a covalent bond between R^{9ao} and R^{10ao}.

- 10 In one embodiment the invention is concerned with compounds of the general formula XXXXIII, wherein R^{5ao}, R^{6ao}, R^{7ao} and R^{8ao} are independently selected from hydrogen and fluor.

In another embodiment the invention is concerned with compounds of the general formula XXXXIII, wherein R^{8ao}, R^{10ao}, R^{11ao} and R^{12ao} are independently selected from hydrogen and

15 fluor.

In another aspect of the invention, compounds are of the general formulae XXXXIVa-b



(XXXXIVa-b)

20

wherein R^{1ap} and R^{2ap} are independently selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy,

25

sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; with the proviso that if R^{1ap} and R^{2ap} are identical then they are not methyl or ethyl; and

wherein there may optionally be a covalent bond between the substituents R^{1ap} and R^{2ap} ; and

wherein R^{5ap} , R^{6ap} and R^{7ap} are independently selected from hydrogen and F; and



wherein R^{4ap} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl.

heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy;

5 with the proviso that R^{4ap} is not methyl.

In one embodiment the invention is concerned with compounds of the general formulae XXXXIVa-b, wherein R^{1ap} and R^{2ap} are not identical.

10 In another aspect of the invention, compounds are of the general formulae XXXXVa-b



wherein R^{1aq} and R^{2aq} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl

$_{10}$ -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, perhalomethyl and perhalomethoxy; with the proviso that R^{1aq} and R^{2aq} are not both methyl; and

wherein there may optionally be a covalent bond between R^{1aq} and R^{2aq}; and

wherein R^{3aq}, R^{4aq}, R^{5aq}, R^{6aq}, R^{7aq}, R^{8aq}, R^{9aq}, R^{10aq} and R^{11ao} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl; and

wherein there may optionally be a covalent bond between R^{4aq} and R^{5aq}; and

wherein R^{3aq} is selected from the group consisting of $-C(=O)-$, $-C(=O)NH-$, $-CH_2-$, $-CH_2CH_2-$, $-CHF-$, $-CH_2CHF-$, $-CHFCH_2-$, $-NH-$, $-S(=O)_2NH-$, $-NH-S(=O)_2-$, $-NHC(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$; and

- 5 wherein there may optionally be a covalent bond between two substituents selected from R^{8aq} , R^{9aq} , R^{10aq} and R^{11aq} .


In one embodiment the invention is concerned with compounds of the general formulae XXXVa-b, wherein there is one covalent bond between the substituents R^{1aq} and R^{2aq} .

- 10 In another embodiment the invention is concerned with compounds of the general formulae XXXVa-b, wherein R^{4aq} , R^{5aq} , R^{6aq} and R^{7aq} are independently selected from hydrogen and fluor.

In another embodiment the invention is concerned with compounds of the general formula XXXVa-b, wherein R^{3aq} is selected from $-O-$, $-S-$, $-CH_2-$, $-CH_2CH_2-$, $-NH-$, $-NH-C(=O)-$, $-$

- 15 $C(=O)-NH-$, $-S(=O)_2NH-$ and $-NH-S(=O)_2-$.

In another aspect of the invention, compounds are of the general formula XXXVI



20

wherein A^{1ar} is N or $C-R^{10ar}$; and A^{2ar} is N or $C-R^{11ar}$; and A^{3ar} is N or $C-R^{12ar}$; and

wherein R^{1ar} , R^{2ar} , R^{3ar} , R^{4ar} , R^{5ar} , R^{6ar} , R^{7ar} , R^{8ar} , R^{9ar} , R^{10ar} , R^{11ar} and R^{12ar} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl,

25

sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo,

amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; with the proviso that if R^{1ar} and R^{2ar} are identical then they are not methyl or ethyl; and

wherein there may optionally be a covalent bond between R^{1ar} and R^{2ar} .


In one embodiment the invention is concerned with compounds of the general formula XXXXVI, wherein R^{9ar} is S(=O)₂-R^{13ar} or C(=O)-R^{13ar}; and wherein R^{13ar} is selected from hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl.

may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy

In another embodiment the invention is concerned with compounds of the general formula XXXXVI, wherein R^{1ar} and R^{2ar} are not identical.

In another embodiment the invention is concerned with compounds of the general formula XXXXVI, wherein there is a covalent bond between the substituents R^{1ar} and R^{2ar}.

In another aspect of the invention, compounds are of the general formula XXXXVII



wherein R^{1as}, R^{2as}, R^{3as}, R^{4as}, R^{5as}, R^{6as}, R^{7as}, R^{8as} and R^{9as} are independently selected from hydrogen and fluor; and

wherein R^{10as} is selected from aryl and heteroaryl, which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo,

amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

5

wherein R^{12as} is selected from hydrogen, hydroxy, sulfanyl, sulfo, amino, cyano, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and


wherein R^{13as}, R^{14as}, R^{15as} and R^{16as} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or

more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl may optionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; with the proviso that R^{12as}, R^{13as}, R^{14as}, R^{15as} and R^{16as} are not all hydrogen; and

wherein there may optionally be a covalent bond between the substituents R^{13as} and R^{14as}.

In one embodiment the invention is concerned with compounds of the general formula XXXXVII, wherein R^{13as}, R^{14as}, R^{15as} and R^{16as} are independently selected from hydrogen and fluor.

In another aspect of the invention, compounds are of the general formula XXXXVIII



(XXXXVIII)

25

wherein R^{5at}, R^{6at}, R^{7at}, R^{8at}, R^{9at}, R^{10at}, R^{11at}, R^{12at}, R^{13at}, R^{14at}, R^{15at}, R^{16at}, R^{17at} and R^{18at} are independently selected from hydrogen and fluor, and

wherein R^{4at} is selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl, and C₃₋₁₀-cycloalkyl, wherein each

30

[illegible]

tionally be substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, fluor, iodine, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl; with the proviso that R^{19at} is not methoxy.

- 5 In one embodiment the invention is concerned with compounds of the general formula XXXXVIII, wherein R^{5at}, R^{6at}, R^{7at}, R^{8at}, R^{9at}, R^{10at}, R^{11at}, R^{12at}, R^{13at}, R^{14at}, R^{15at}, R^{16at}, R^{17at} and R^{18at} are all hydrogen.

In another embodiment the invention is concerned with compounds of the general formula XXXXVIII, wherein R^{4at} is an optionally substituted aryl or heteroaryl.

10

In another embodiment, the invention is concerned with compounds of the general formulae I-XXXXVIII, wherein said compound comprises only one F.


In another embodiment, the invention is concerned with compounds of the general formulae I-XXXXVIII, wherein said compound comprises two F and preferably three F.

- 15 In another embodiment, the invention is concerned with compounds of the general formulae I-XXXXVIII, wherein said compound comprises a CF₃ moiety.

In another embodiment, the invention is concerned with compounds of the general formulae I-XXXXVIII, wherein said compound comprises a hydrophilic substituent selected from the group consisting of hydroxy, amino, C₁₋₆-alkoxy, -C(=O)NH₂, -NHC(=O)-OH, -S(=O)₂-NH₂, -NHS(=O)₂-OH, -NHC(=O)-R^{1am}, -NHS(=O)₂-R^{1am}, -N(R^{1am})C(=O)-R^{2am}, -N(R^{1am})S(=O)₂-R^{2am},
20 wherein R^{1am} and R^{2am} are independently selected from C₁₋₆-alkyl.

In another embodiment, the invention is concerned with compounds of the general formulae I-XXXXVIII, wherein said compound comprises a moiety selected from the group consisting of

25



Examples of specific compounds of the invention are :

- Methyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 3-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-ylamino)-phenyl ester,
5 Methyl-phenyl-carbamic acid 4-(3,5-dichloro-pyridin-4-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-cyano-5-trifluoromethyl-pyridine-3-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 2-benzenesulfonyl-4-(3-chloro-5-trifluoromethyl-pyridine-2-yloxy)-phenyl ester,
10 Methyl-phenyl-carbamic acid 4-tert-butoxy-phenyl ester,
Methyl-phenyl-carbamic acid 3-(4-fluorobenzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 4-phenoxy-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-chlorobenzoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-yloxy)-phenyl ester,
15 Methyl-phenyl-carbamic acid 4-[4-(4-chloro-phenyl)-thiazol-2-yl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-pyrrol-1-yl-phenyl ester,
Methyl-phenyl-carbamic acid 4-imidazol-1-yl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-ylmethyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-trifluoromethylsulfanyl-phenyl ester,
20 Methyl-phenyl-carbamic acid 4-pentafluoroethoxy-phenyl ester,
Methyl-phenyl-carbamic acid 4-benzyloxy-phenyl ester,
Methyl-phenyl-carbamic acid 4-benzyl-phenyl ester,
Methyl-phenyl-carbamic acid 4'-cyano-biphenyl-4-yl-ester,
Methyl-phenyl-carbamic acid 4'-bromo-biphenyl-4-yl-ester,
25 Methyl-phenyl-carbamic acid biphenyl-4-yl-ester,
Methyl-phenyl-carbamic acid 4-[3-(4-chlorophenyl)-ureido]-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-nitro-phenoxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-heptylsulfanyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-butoxy-phenyl ester,
30 Methyl-phenyl-carbamic acid 4-(4-chloro-benzenesulfonyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-chloromethyl-thiazol-2-yl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
cis-Methyl-phenyl-carbamic acid 4-(1,3-dioxo-octahydro-isoidol-2-yl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(cyclohexanecarbonyl-amino)-phenyl ester,
35 Methyl-phenyl-carbamic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,

- cis/trans-Methyl-phenyl-carbamic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
cis-Methyl-phenyl-carbamic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
trans-Methyl-phenyl-carbamic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
5 Methyl-phenyl-carbamic acid 4-(3,3-dimethyl-butyrylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 3-benzyl-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
10 Methyl-phenyl-carbamic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propyl-2-oxo-2H-chromen-7-yl ester,
15 Methyl-phenyl-carbamic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
20 Methyl-phenyl-carbamic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Methyl-phenyl-carbamic acid 4-pyrrolidine-1-yl-phenyl ester,
Methyl-phenyl-carbamic acid 4-piperidine-1-yl-phenyl ester,
Methyl-phenyl-carbamic acid 4-morpholine-1-yl-phenyl ester,
25 Methyl-phenyl-carbamic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[(pyridine-2-carbonyl)-amino]-phenyl ester,
4-Chlor-phenyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Chlor-phenyl-methyl-carbamic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,
30 (4-Chloro-phenyl)-methyl-carbamic acid 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenyl ester,
Ethyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Ethyl-phenyl-carbamic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,
Methyl-phenyl-carbamic acid pyrazol-1-yl ester,
35 Methyl-phenyl-carbamic acid 4-bromo-pyrazol-1-yl ester,

- Methyl-phenyl-carbamic acid 3,4,5-tribromo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid imidazol-1-yl ester,
Methyl-phenyl-carbamic acid [1,2,3]triazol-1-yl ester,
5 Methyl-phenyl-carbamic acid 3-bromo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-bromo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-chloro-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-chloro-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-chloro-pyrazol-1-yl ester,
10 Methyl-phenyl-carbamic acid 4-iodo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-iodo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-methyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-methyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-methyl-pyrazol-1-yl ester,
15 Methyl-phenyl-carbamic acid 4-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(2-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(2-nitro-phenyl)-pyrazol-1-yl ester,
20 Methyl-phenyl-carbamic acid 3-pyridin-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-pyridin-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-(2-fluoro-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-(2-fluoro-phenyl)-pyrazol-1-yl ester,
25 Methyl-phenyl-carbamic acid 3-phenylsulfanyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-phenylsulfanyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-phenylsulfanyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-thiophen-3-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-thiophen-2-yl-pyrazol-1-yl ester,
30 Methyl-phenyl-carbamic acid 4-thiophen-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-thiophen-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-chloro-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-bromo-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-iodo-imidazol-1-yl ester,
35 Methyl-phenyl-carbamic acid 2-methyl-imidazol-1-yl ester,

- Methyl-phenyl-carbamic acid 2-phenylsulfanyl-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-(4-fluoro-phenyl)-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-thiophen-2-yl-imidazol-1-yl ester,
5 Methyl-phenyl-carbamic acid 2-pyridin-2-yl-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2,5-dichloro-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-bromo-2,5-dichloro-imidazol-1-yl ester,
4-(Methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester,
Methyl-phenyl-carbamic acid 4-(1,3,5-trimethyl-1H-pyrazol-4-ylmethyl)-phenyl ester,
10 Methyl-phenyl-carbamic acid 4-(2-cyano-ethyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-([1,2,3,4]thiaziazol-5-ylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-pentyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-methoxy-ethyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-acetyl-phenyl ester,
15 Methyl-phenyl-carbamic acid pyridin-4-yl ester,
Methyl-phenyl-carbamic acid pyridin-3-yl ester,
Methyl-phenyl-carbamic acid 6-methyl-pyridin-3-yl ester,
Methyl-phenyl-carbamic acid isoquinolin-1-yl ester,
Methyl-phenyl-carbamic acid 3-phenoxy-phenyl ester,
20 Methyl-phenyl-carbamic acid 3-acetyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-acetyl-2-carbamoyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-acetyl-3-methyl-phenyl ester,
Methyl-phenyl-carbamic acid 1-oxo-indan-4-yl ester,
Methyl-phenyl-carbamic acid benzothiazol-2-yl ester,
25 Methyl-phenyl-carbamic acid 5-oxo-5,6,7,8-tetrahydro-naphthalen-2-yl ester,
Methyl-phenyl-carbamic acid benzo[d]isoxazol-3-yl ester,
Methyl-phenyl-carbamic acid pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 1-(methyl-phenyl-carbamoyl)-1H-benzimidazol-2-yl ester,
Methyl-phenyl-carbamic acid 4-[(pyridine-3-carbonyl)-amino]-phenyl ester,
30 Methyl-phenyl-carbamic acid 4-(3-pyridin-3-yl-acryloyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-[3-(3,4,5-trimethoxy-phenyl)-acryloyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-diethylcarbamoyl-2-methoxy-phenyl ester,
Methyl-phenyl-carbamic acid 3-phenylcarbamoyl-phenyl ester,
Methyl-phenyl-carbamic acid quinolin-7-yl ester,
35 Methyl-phenyl-carbamic acid 4-(4-methyl-piperazine-1-carbonyl)-phenyl ester,

- Methyl-phenyl-carbamic acid 3-acetylamino-phenyl ester,
Methyl-phenyl-carbamic acid 4-benzoyl-phenyl ester,
Methyl-phenyl-carbamic acid biphenyl-3-yl ester,
Methyl-phenyl-carbamic acid 1H-indol-4-yl ester,
5 Methyl-phenyl-carbamic acid 5,6,7,8-tetrahydro-naphthalen-1-yl ester,
Methyl-phenyl-carbamic acid 5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yl ester,
Methyl-phenyl-carbamic acid 1,3-dioxo-1,3-dihydro-isobenzofuran-4-yl ester,
Methyl-phenyl-carbamic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester,
Methyl-phenyl-carbamic acid 4-morpholin-4-yl-phenyl ester,
10 Methyl-phenyl-carbamic acid 4-(5,6-dichloro-1,3-dioxo-1,3-dihydro-isoindol-2-yl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-phenoxy-acetylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(4-chloro-phenyl)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[(pyridine-2-carbonyl)-amino]-phenyl ester,
15 Methyl-phenyl-carbamic acid 4-[methyl-(thiophene-2-carbonyl)-amino]-phenyl ester,
Methyl-phenyl-carbamic acid 4-butyrylamino-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4,6-dimethyl-pyrimidin-2-ylsulfanyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-methanesulfonyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(3-oxo-1,2,3,4-tetrahydro-quinoxalin-2-yl)-acetylamino]-
20 phenyl ester,
Methyl-phenyl-carbamic acid 4-phenylacetyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-[[4-(methyl-phenyl-carbamoyloxy)-2-oxo-1,2-dihydro-quinoline-3-carbonyl]-amino]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[(4-hydroxy-2-oxo-1,2-dihydro-quinoline-3-carbonyl)-amino]-
25 phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-hydroxy-benzyl)-phenyl ester
Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-benzylcarbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(butyl-methyl-carbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(methyl-phenethyl-carbamoyl)-phenyl ester,
30 Methyl-phenyl-carbamic acid 4-[(pyridin-2-ylmethyl)-carbamoyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-pyridin-2-yl-ethylcarbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-phenylamino-ethylcarbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(3-methyl-butylcarbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester,
35 Methyl-phenyl-carbamic acid 4-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-phenyl ester,

- Methyl-phenyl-carbamic acid 4-cyclohexylcarbamoyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-cyclopropylcarbamoyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(cyclohexylmethyl-carbamoyl)-phenyl ester,
Methyl-phenyl-carbamic acid 5-nitro-pyridin-2-yl ester,
5 Methyl-phenyl-carbamic acid pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 7-chloro-quinolin-4-yl ester,
Methyl-phenyl-carbamic acid quinolin-4-yl ester,
Methyl-phenyl-carbamic acid 5-methyl-isoxazol-3-yl ester,
Methyl-phenyl-carbamic acid quinoxalin-2-yl ester,
10 Methyl-phenyl-carbamic acid 4-methyl-quinolin-2-yl ester,
Methyl-phenyl-carbamic acid 3-methyl-quinoxalin-2-yl ester,
Methyl-phenyl-carbamic acid 4,6-dimethyl-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid isoquinolin-6-yl ester,
Methyl-phenyl-carbamic acid quinolin-2-yl ester,
15 Methyl-phenyl-carbamic acid isoquinolin-3-yl ester,
Methyl-phenyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 3-nitro-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-chloro-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(2-nitro-phenyl)-pyrimidin-2-yl ester,
20 Methyl-phenyl-carbamic acid 5-trifluoromethyl-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 3-chloro-5-trifluoromethyl-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-nitro-3-trifluoromethyl-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 4,5-dichloro-pyridazin-3-yl ester,
Methyl-phenyl-carbamic acid 5-benzoylamino-pyridin-2-yl ester,
25 Methyl-phenyl-carbamic acid 5-(cyclohexanecarbonyl-amino)-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl ester,
Methyl-phenyl-carbamic acid 5-(2,2-dimethyl-propionylamino)-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(2-cyclohexyl-acetylamino)-pyridin-2-yl ester,
30 Methyl-phenyl-carbamic acid 5-(4-methoxy-phenoxy)-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(3,4-dichloro-phenoxy)-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 6-pyridin-2-ylmethyl-pyridazin-3-yl ester,
Methyl-phenyl-carbamic acid 6-(4-methoxy-benzyl)-pyridazin-3-yl ester,
Methyl-phenyl-carbamic acid 6-(2,4-dichloro-benzyl)-pyridazin-3-yl ester,
35 Methyl-phenyl-carbamic acid 4-iodo-pyrazol-1-yl ester,

- Methyl-phenyl-carbamic acid benzotriazol-1-yl ester,
Methyl-phenyl-carbamic acid [1,2,3]triazolo[4,5-b]pyridin-3-yl ester,
Methyl-phenyl-carbamic acid 3-(2-nitro-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester,
5 Methyl-phenyl-carbamic acid 3-pyridin-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-thiophen-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 3-bromo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-iodo-pyrazol-1-yl ester,
10 Methyl-phenyl-carbamic acid 2-chloro-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-benzoyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-(4-dimethylamino-phenyl)-pyrazol-1-yl ester,
15 Methyl-phenyl-carbamic acid 4,5-diiodo-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 5-thiophen-2-yl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-methylsulfanyl-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 3,5-bis-(4-methoxy-phenyl)-pyrazol-1-yl ester,
20 Methyl-phenyl-carbamic acid 4-(4-fluoro-phenyl)-5-(4-methoxy-phenyl)-3-(4-methylphenyl)-
pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-benzyl-5-(4-methoxy-phenyl)-3-(methylphenyl)-pyrazol-1-yl
ester,
Methyl-phenyl-carbamic acid 4-acetyl-pyrazol-1-yl ester,
25 Methyl-phenyl-carbamic acid 2-(4-nitro-phenyl)-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-chloro-5-(4-methylphenyl)-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-formyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-hydroxymethyl-pyrazol-1-yl ester,
Methyl-phenyl-carbamic acid 4-phenylethynyl-pyrazol-1-yl ester,
30 Methyl-phenyl-carbamic acid 2-bromo-imidazol-1-yl ester,
Methyl-phenyl-carbamic acid 2-phenylsulfanyl-imidazol-1-yl ester,
Methyl-o-tolyl-carbamic acid 4-(trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Methyl-m-tolyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(3-Chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
35 Methyl-p-tolyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

- (3-Fluoro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(3-Chloro-phenyl)-methyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester,
Methyl-m-tolyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-butyrylamino)-pyridin-2-yl ester,
5 Methyl-phenyl-carbamic acid 5-[(pyridine-2-carbonyl)-amino]-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 2-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-pyrimidin-5-yl ester,
Methyl-phenyl-carbamic acid 5-bromo-pyrimidin-2-yl ester,
Methyl-phenyl-carbamic acid 5-[(6-chloro-pyridine-3-carbonyl)-amino]-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(2,2-dimethyl-propylcarbonyl)-pyridin-2-yl ester,
10 Methyl-phenyl-carbamic acid 6-(3,4-dichloro-phenoxy)-pyridazin-3-yl ester,
Methyl-phenyl-carbamic acid 2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl ester,
Methyl-phenyl-carbamic acid 5-(2,5-dioxo-pyrrolidin-1-yl)-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 5-(4-trifluoromethyl-benzoylamino)-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid quinolin-6-yl ester,
15 Methyl-phenyl-carbamic acid 5-(4-chloro-benzoylamino)-pyridin-2-yl ester,
4 Methyl-phenyl-carbamic acid 5-(4-methoxy-benzoylamino)-pyridin-2-yl ester,
Methyl-phenyl-carbamic acid 4,4-dimethyl-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl ester,
Methyl-phenyl-carbamic acid 2-methyl-quinolin-6-yl ester,
{2-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-ethyl}-carbamic acid tert-butyl ester,
20 Methyl-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(5-dimethylamino-naphthalene-1-sulfonylamino)-ethyl]-
phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(3,4-difluoro-benzenesulfonylamino)-ethyl]-phenyl ester,
25 2-[2-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-ethylsulfamoyl]-benzoic acid methyl ester,
Methyl-phenyl-carbamic acid 4-[2-(2,5-dichloro-thiophene-3-sulfonylamino)-ethyl]-phenyl es-
ter,
Methyl-phenyl-carbamic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-phenyl
ester,
30 Methyl-phenyl-carbamic acid 4-[2-(1-methyl-1H-imidazole-4-sulfonylamino)-ethyl]-phenyl es-
ter,
Methyl-phenyl-carbamic acid 4-[2-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonylamino)-ethyl]-
phenyl,
Methyl-phenyl-carbamic acid 4-[2-(4-nitro-benzenesulfonylamino)-ethyl]-phenyl ester,
35 Methyl-phenyl-carbamic acid 4-[2-(6-chloro-imidazo[2,1-b]thiazole-5-sulfonylamino)-ethyl]-

- phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-(2-trifluoromethoxy-benzenesulfonylamino)-ethyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-(2-dimethylaminosulfonylamino-ethyl)-phenyl ester,,
 5 Methyl-phenyl-carbamic acid 4-(2-methanesulfonylamino-ethyl)-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-(6-morpholin-4-yl-pyridine-3-sulfonylamino)-ethyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-(6-phenoxy-pyridine-3-sulfonylamino)-ethyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonylamino]-ethyl]-phenyl ester,
 10 Methyl-phenyl-carbamic acid 4-[2-(4-dimethylamino-benzenesulfonylamino)-ethyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-[4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfonylamino]-ethyl]-phenyl ester,
 15 Methyl-phenyl-carbamic acid 4-(2-cyclohexyl-ethylsulfamoyl)-phenyl ester,
 Methyl-phenyl-carbamic acid 4-(3-methyl-butylsulfamoyl)-phenyl ester,
 Methyl-phenyl-carbamic acid 4-(1,1,3,3-tetramethyl-butylcarbamoyl)-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[(2-dimethylamino-ethyl)-methyl-carbamoyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-(cyclopropylmethyl-carbamoyl)-phenyl ester,
 20 Methyl-phenyl-carbamic acid 4-(methyl-pyridin-3-ylmethyl-carbamoyl)-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[(1H-benzimidazol-2-ylmethyl)-carbamoyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-phenyl ester,
 Methyl-phenyl-carbamic acid 5-amino-pyridin-2-yl ester,
 Methyl-phenyl-carbamic acid 5-benzenesulfonylamino-pyridin-2-yl ester,
 25 3,3-Dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbamoyl]-butyric acid,
 2,2-Dimethyl-N-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-yl]-succinamic acid,
 Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-2,5-dioxo-pyrrolidin-1-yl)-pyridin-2-yl ester,
 Methyl-phenyl-carbamic acid 5-[3,3-dimethyl-5-(4-methyl-piperazin-1-yl)-5-oxo-pentanoylamino]-pyridin-2-yl ester,
 30 Methyl-phenyl-carbamic acid 5-[3,3-dimethyl-4-(pyridin-3-ylcarbamoyl)-butyrylamino]-pyridin-2-yl ester,
 Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-5-morpholin-4-yl-5-oxo-pentanoylamino)-pyridin-2-yl ester,
 Methyl-phenyl-carbamic acid 5-[4-(2-dimethylamino-ethylcarbamoyl)-3,3-dimethyl-
 35 butyrylamino]-pyridin-2-yl ester,

- Methyl-phenyl-carbamic acid 4-iodo-phenyl ester,
Methyl-phenyl-carbamic acid 4'-trifluoromethyl-biphenyl-4-yl ester,
Methyl-phenyl-carbamic acid 4'-trifluoromethoxy-biphenyl-4-yl ester,
Methyl-phenyl-carbamic acid 4-pyridin-3-yl-phenyl ester,
5 Methyl-phenyl-carbamic acid 4-(5-chloro-thiophen-2-yl)-phenyl ester,
Methyl-phenyl-carbamic acid 4'-benzylsulfamoyl-biphenyl-4-yl ester,
Methyl-phenyl-carbamic acid 4-styryl-phenyl ester,
Methyl-phenyl-carbamic acid 4-phenylethynyl-phenyl ester,
3-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-acrylic acid methyl ester,
10 Methyl-phenyl-carbamic acid 4-(toluene-4-sulfonylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(1-methyl-1H-imidazole-4-sulfonylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2,5-dichloro-thiophene-3-sulfonylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonylamino)-phenyl
15 ester,
Methyl-phenyl-carbamic acid 4-(5-dimethylamino-naphthalene-1-sulfonylamino)-phenyl ester,
2-[4-(Methyl-phenyl-carbamoyloxy)-phenylsulfamoyl]-benzoic acid methyl ester,
Methyl-phenyl-carbamic acid 4-(3,4-difluoro-benzenesulfonylamino)-phenyl ester,
Methyl-phenyl-carbamic acid 4-pyridin-2-ylmethyl-phenyl ester,
20 Methyl-phenyl-carbamic acid 4-pyridin-3-ylmethyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-benzyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-thiophen-3-ylmethyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-thiophen-2-ylmethyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(4-amino-benzenesulfonylamino)-ethyl]-phenyl ester,
25 Methyl-phenyl-carbamic acid 4-[2-[(pyridine-3-carbonyl)-amino]-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(2-dimethylamino-acetyl)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 2-(toluene-4-sulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl ester,
Methyl-phenyl-carbamic acid 4-[4-(2-pyrrolidin-1-yl-ethoxy)-benzyl]-phenyl ester,
Methyl-phenyl-carbamic acid 2-[4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfonyl]-1,2,3,4-
30 tetrahydro-isoquinolin-7-yl ester,
Methyl-phenyl-carbamic acid 4-[2-[(1-methyl-piperidine-4-carbonyl)-amino]-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 2-(3,4-difluoro-benzenesulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl ester,
35 Methyl-phenyl-carbamic acid 1-methyl-2-(toluene-4-sulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-

yl ester,

Methyl-phenyl-carbamic acid 2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydro-isoquinolin,

Methyl-phenyl-carbamic acid 1-methyl-2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-

5 1,2,3,4-tetrahydro-isoquinolin-7-yl ester,

3,3-Dimethyl-4-{2-[4-(methyl-phenyl-carbamoyloxy)-phenyl]-ethylcarbamoyl}-butyric acid,

Methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-yl)-benzoylamino]-ethyl}-phenyl ester,

Methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-ylmethyl)-benzoylamino]-ethyl}-phenyl ester,

10

Methyl-phenyl-carbamic acid 4-[2-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-ethyl]-phenyl ester,

3,3-Dimethyl-4-{2-[4-(methyl-phenyl-carbamoyloxy)-phenyl]-ethylcarbamoyl}-butyric acid ethyl ester,

Methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester,

15 Methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-(4-dimethylamino-pyridin-2-ylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-(4-imidazol-1-yl-phenoxy-methyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-[4-(2-dimethylamino-ethyl)-phenoxy-methyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-(pyrazol-1-yloxy-methyl)-phenyl ester,

20 Methyl-phenyl-carbamic acid 4-(imidazol-1-yloxy-methyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-(2-oxo-2H-pyridin-1-ylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(pyridin-2-yloxy)-ethyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(4-imidazol-1-yl-phenoxy)-ethyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-{2-[4-(2-dimethylamino-ethyl)-phenoxy]-ethyl}-phenyl ester,

25 Methyl-phenyl-carbamic acid 4-[2-(pyrazol-1-yloxy)-ethyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(imidazol-1-yloxy)-ethyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-(4-oxo-4H-pyridin-1-ylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(pyridin-3-yloxy)-ethyl]-phenyl ester,

30 Methyl-phenyl-carbamic acid 4-(2-oxo-2H-pyridin-1-ylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-(pyridin-3-yloxy-methyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(2,5-dioxo-pyrrolidin-1-yl)-ethyl]-phenyl ester,

Methyl-phenyl-carbamic acid 4-[2-(1,3-dioxo-1,3-dihydro-pyrrolo[3,4-]pyridin-2-yl)-ethyl]-phenyl ester,

35 Methyl-phenyl-carbamic acid 4-(1-methyl-1H-imidazol-2-ylsulfanylmethyl)-phenyl ester,

Methyl-phenyl-carbamic acid 4-tetrazol-1-ylmethyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2,5-dioxo-pyrrolidin-1-ylmethyl)-phenyl ester, Methyl-phenyl-carbamic acid 4-[2-(2-thioxo-2H-pyridin-1-yl)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-(1,3-dioxo-1,3-dihydro-pyrrolo[3,4]pyridin-2-ylmethyl)-phenyl ester,
5 Methyl-phenyl-carbamic acid 4-[1,2,4]triazol-1-ylmethyl-phenyl ester,
Methyl-phenyl-carbamic acid 4-(2-thioxo-2H-pyridin-1-ylmethyl)-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-phenyl ester,
Ethyl-phenyl-carbamic acid 4-(2-tetrazol-1-yl-ethyl)-phenyl ester,
10 Methyl-phenyl-carbamic acid 4-[2-(pyrimidin-2-yloxy)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(pyridin-4-ylsulfanyl)-ethyl]-phenyl ester,
Methyl-phenyl-carbamic acid 4-[2-(1-pyridin-3-yl-1H-imidazol-2-ylsulfanyl)-ethyl]-phenyl ester,
and
Methyl-phenyl-carbamic acid 4-[2-(1,3-dioxo-1,3-dihydro-isindol-2-yl)-ethyl]-phenyl ester.

15

Further examples of specific compounds of the invention are :

Benzyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester, and
Benzyl-methyl-carbamic acid 4-(3,5-dichloro pyridin-2-yloxy)-phenyl ester.

20 Further examples of specific compounds of the invention are :

Isopropyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Cyclohexyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.
Dimethyl-carbamic acid 4-(3,5-dichloro pyridin-2-yloxy)-phenyl ester,
Methyl-pyridin-2-yl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
25 (2-Dimethylamino-ethyl)methylcarbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(6-Methoxy-pyridin-2-yl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(4-Methoxy-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
30 (2-Methoxy-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(2-Carbamoyl-4-chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
(2-Carbamoyl-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
35 (2-Chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

(2,4-Difluoro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
and

Methyl-(2-trifluoromethoxy-phenyl)-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester.

5

Further examples of specific compounds of the invention are :

Pyrrolidine-1-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,

2,3-Dihydro-indole-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester. and

1,3-Dihydro-isoindole-2-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

10

Further examples of specific compounds of the invention are :

Piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 3-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,

15

Piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-ylamino)-phenyl ester,

Piperidine-1-carboxylic acid 4-(3,5-dichloro-pyridin-4-yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,

Piperidine-1-carboxylic acid 4-(2-cyano-5-trifluoromethyl-pyridine-3-yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 2-benzenesulfonyl-4-(3-chloro-5-trifluoromethyl-pyridine-2-

20

yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 4-tert-butoxy-phenyl ester,

Piperidine-1-carboxylic acid 3-(4-fluorobenzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

Piperidine-1-carboxylic acid 4-phenoxy-phenyl ester,

Piperidine-1-carboxylic acid 4-(4-chlorobenzoyl)-phenyl ester,

25

Piperidine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-yloxy)-phenyl ester,

Piperidine-1-carboxylic acid 4-[4-(4-chloro-phenyl)-thiazol-2-yl]-phenyl ester,

Piperidine-1-carboxylic acid 4-pyrrol-1-yl-phenyl ester,

Piperidine-1-carboxylic acid 4-imidazol-1-yl-phenyl ester,

Piperidine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-ylmethyl)-phenyl ester,

30

Piperidine-1-carboxylic acid 4-trifluoromethylsulfanyl-phenyl ester,

Piperidine-1-carboxylic acid 4-pentafluoromethyloxy-phenyl ester,

Piperidine-1-carboxylic acid 4-benzyloxy-phenyl ester,

Piperidine-1-carboxylic acid 4-benzyl-phenyl ester,

Piperidine-1-carboxylic acid 4'-cyano-biphenyl-4-yl-ester,

35

Piperidine-1-carboxylic acid 4'-bromo-biphenyl-4-yl-ester,

- Piperidine-1-carboxylic acid biphenyl-4-yl-ester,
Piperidine-1-carboxylic acid 4-[3-(4-chlorophenyl)-ureido]-phenyl ester,
Piperidine-1-carboxylic acid 4-(4-nitro-phenoxy)-phenyl ester,
Piperidine-1-carboxylic acid 4-heptylsulfanyl-phenyl ester,
5 Piperidine-1-carboxylic acid 4-butoxy-phenyl ester,
Piperidine-1-carboxylic acid 4-(4-chloro-benzenesulfonyl)-phenyl ester,
Piperidine-1-carboxylic acid 4-(4-chloromethyl-thiazol-2-yl)-phenyl ester,
Piperidine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
cis-Piperidine-1-carboxylic acid 4-(1,3-dioxo-octahydro-isoindol-2-yl)-phenyl ester,
10 Piperidine-1-carboxylic acid 4-(cyclohexanecarbonyl-amino)-phenyl ester,
Piperidine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,
cis/trans-Piperidine-1-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl
ester,
cis- Piperidine-1-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
15 trans- Piperidine-1-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
Piperidine-1-carboxylic acid 4-(3,3-dimethyl-butyrylamino)-phenyl ester,
Piperidine-1-carboxylic acid 3-benzyl-4-methyl-2-oxo-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl
20 ester,
Piperidine-1-carboxylic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl
ester,
Piperidine-1-carboxylic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propyl-2-oxo-2H-
25 chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Piperidine-1-carboxylic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
30 Piperidine-1-carboxylic acid 4-pyrrolidine-1-yl-phenyl ester,
Piperidine-1-carboxylic acid 4-piperidine-1-yl-phenyl ester,
Piperidine-1-carboxylic acid 4-morpholine-1-yl-phenyl ester,
Piperidine-1-carboxylic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
Piperidine-1-carboxylic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
35 Piperidine-1-carboxylic acid 4-[(pyridine-2-carbonyl)-amino]-phenyl ester,

- Piperidine-1-carboxylic acid pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-bromo-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 4-bromo-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 5-bromo-pyrazol-1-yl ester,
5 Piperidine-1-carboxylic acid 3,4,5-tribromo-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 4-chloro-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-(2-methoxy-phenyl)-pyrazol-1-yl ester,
10 Piperidine-1-carboxylic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-pyridin-2-yl-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 4-phenylsulfanyl-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid 3-thiophen-2-yl-pyrazol-1-yl ester,
15 Piperidine-1-carboxylic acid 5-thiophen-2-yl-pyrazol-1-yl ester,
Piperidine-1-carboxylic acid imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-chloro-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-bromo-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-iodo-imidazol-1-yl ester,
20 Piperidine-1-carboxylic acid 2-methyl-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-phenylsulfanyl-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-(4-fluoro-phenyl)-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2-thiophen-2-yl-imidazol-1-yl ester,
25 Piperidine-1-carboxylic acid 2-pyridin-2-yl-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 2,5-dichloro-imidazol-1-yl ester,
Piperidine-1-carboxylic acid 4-bromo-2,5-dichloro-imidazol-1-yl ester,
2-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
3-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
30 4-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Benzyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
1,4-Dioxo-8-aza-spiro[4.5]decane-8-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
3-Hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
35 3,4-Dihydro-1H-isoquinoline-2-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl es-

ter,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenyl ester,

5 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-benzyl-4-methyl-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

10 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl ester,

15 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(4-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propy-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

20 3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,

25 7-Trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-Oxo-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

30 4-[5-(4-Dimethylamino-phenyl)-1H-pyrazol-3-yl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-(5-Furan-2-yl-1H-pyrazol-3-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-Benzylamino-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

35 4-(3,4-Dihydro-1H-isoquinolin-2-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-

- pyridin-2-yloxy)-phenyl ester,
3-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
3-Hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
5 4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Pyrrolidin-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
10 4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester,
1,4-Dioxo-8-aza-spiro[4.5]decane-8-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Benzoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
[1,4']Bipiperidinyl-1'-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
15 4-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
3-Diethylcarbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Carbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
20 3-Carbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(tert-Butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-trifluoromethyl-pyridin-2-yl ester,
25 4-Hydroxy-piperidine-1-carboxylic acid 5-(4-chloro-benzoylamino)-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(3-methoxy-benzoylamino)-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(4-methoxy-benzoylamino)-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(2,4-dichloro-benzoylamino)-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(4-trifluoromethyl-benzoylamino)-pyridin-2-yl ester,
30 4-Hydroxy-piperidine-1-carboxylic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(5-trifluoromethyl-pyridin-2-yloxy)-pyridin-2-yl ester,
4-Hydroxy-piperidine-1-carboxylic acid 5-(3,5-dichloro-pyridin-2-yloxy)-pyridin-2-yl ester,
4-Aminomethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
35 4-Benzimidazol-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl

ester,

4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,

4-(4-Amino-phenyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

5 4-(Methyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-(2-Oxo-pyrrolidin-1-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

10 4-(Methyl-phenethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-[(Benzyl-ethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-[Methyl-phenethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

15 4-[(Cyclohexyl-methyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-[(Ethyl-pyridin-4-ylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

20 4-[(Benzyl-methyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-[(Methyl-pyridin-3-ylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-(1,3-Dihydro-isoindol-2-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

25 4-Benzotriazol-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-[(Cyclopropylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

30 4-[Methyl-(2-pyridin-2-yl-ethyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-(Cyclohexyl-methyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

4-(Isopropyl-methyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

35 4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbonyl)-phenyl ester

ester,

Piperidine-1-carboxylic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3]bipyridinyl-6'-yl,
4-[Methyl-(2-pyridin-4-yl-ethyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-
pyridin-2-yloxy)-phenyl ester,

- 5 4-(Cyclopropyl-pyridin-4-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-
pyridin-2-yloxy)-phenyl ester,
4-[Cyclopropyl-(2-fluoro-benzyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-
pyridin-2-yloxy)-phenyl ester,
4-(Cyclopropyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-
10 pyridin-2-yloxy)-phenyl ester,
4-(Cyclopropylmethyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-
trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(Cyclopropylmethyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-
trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
15 4-(4-Hydroxy-piperidin-1-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-{3-[1-(2-Hydroxy-ethyl)-piperidin-4-yl]-propyl}-piperidine-1-carboxylic acid 4-(5-
trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(2-Pyrrolidin-1-yl-ethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
20 phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-
phenyl,
4-Hydroxy-piperidine-1-carboxylic acid 4-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-phenyl
25 ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-pyridin-2-ylmethyl-phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-pyridin-3-ylmethyl-phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-(4-trifluoromethyl-benzyl)-phenyl ester,
4-Hydroxy-piperidine-1-carboxylic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester,
30 4-(3-Amino-phenyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl es-
ter, 4-Phenyl-piperidine-1-carboxylic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester, and
4-(4-Methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid 4-(5-methyl-pyridin-2-
ylmethyl)-phenyl ester.

35 Further examples of specific compounds of the invention are :

- 4-Methyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(2-Hydroxyethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
5 4-(Pyridin-2-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(Pyrrolidinocarbonylmethyl)- piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-Phenyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(Isopropylaminocarbonylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
10 yloxy)-phenyl ester,
4-Ethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Propyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Butyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(4-Chlorobenzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
15 ester,
4-(4-Chlorophenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(Diphenylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
20 4-(3-Hydroxypropyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,

4-(3-Trifluoromethylphenyl)- piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
25 4-(3-Chlorophenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(2-Chlorophenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(3,4-Dichlorophenyl)- piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
30 phenyl ester,
4-(4-Fluorophenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(4-Methoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
35 4-(3-Methoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl

- ester,
4-(2-Methoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(2,4-Dimethoxyphenyl)- piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
5 phenyl ester,
4-(3,4,5-Trimethoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-[3-(Trifluoromethyl)pyridin-2-yl]-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
10 4-(3,4-Methylenedioxy-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-(3,4-Methylenedioxy-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-(Pyridin-4-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
15 4-Cyclopentyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(2-Pyrimidinyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl es-
ter,
4-(4-Acetylphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
20 4-(2-(2-Hydroxyethoxy)ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-Benzyl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3-bromo-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-bromo-pyrazol-1-yl ester,
25 4-Benzyl-piperazine-1-carboxylic acid 5-bromo-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3,4,5-tribromo-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-chloro-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester,
30 4-Benzyl-piperazine-1-carboxylic acid 3-(2-methoxy-phenyl)-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 3-pyridin-2-yl-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-phenylsulfanyl-pyrazol-1-yl ester,
35 4-Benzyl-piperazine-1-carboxylic acid 3-thiophen-2-yl-pyrazol-1-yl ester,

- 4-Benzyl-piperazine-1-carboxylic acid 5-thiophen-2-yl-pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-chloro-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-bromo-imidazol-1-yl ester,
5 4-Benzyl-piperazine-1-carboxylic acid 2-iodo-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-methyl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-phenylsulfanyl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-(4-fluoro-phenyl)-imidazol-1-yl ester,
10 4-Benzyl-piperazine-1-carboxylic acid 2-thiophen-2-yl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2-pyridin-2-yl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 2,5-dichloro-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-bromo-2,5-dichloro-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-bromo-2-chloro-imidazol-1-yl ester,
15 4-Benzyl-piperazine-1-carboxylic acid 5-(4-methoxy-phenyl)-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 5-(4-fluoro-phenyl)-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 5-thiophen-2-yl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 5-pyridin-2-yl-imidazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester,
20 4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester,
4-Pyridin-2-yl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
- 4-(1,3-Benzodioxol-5-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
25 phenyl ester,
4-[2-(2-Hydroxyethoxy)ethyl]-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-(Diphenylmethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
30 4-(4-tert-Butylbenzyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl
ester,
4-(4-Fluorobenzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(2-Thienylethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl es-
35 ter,

- 4-(1-Phenylethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
- 4-Octylpiperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
- 4-(3-Dimethylamino-propyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
- 5 phenyl ester,
- 4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 10 4-Phenethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 4-Pyridin-2-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 15 4-(3-Phenylpropyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 4-(4-Phenylbutyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 4-(3,4-Dichlorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl
- 20 ester,
- 4-(4-Fluorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
- 4-(2-Chlorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
- 25 4-Methylpiperazine-1-carboxylic acid 4-chlorophenyl ester,
- 4-(4-Phenylbutyl)piperazine-1-carboxylic acid 4-chlorophenyl ester,
- 4-[2-(2-Hydroxyethoxy)ethyl]piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester,
- 4-(1-Ethylpropyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester,
- 30 4-Cycloheptylpiperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)phenyl ester,
- 4-Cyclohexylpiperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)phenyl ester,
- 4-(4-Chlorobenzyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester,
- 4-(4-Methylbenzyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester,
- 4-(4-Methoxybenzyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
- 35 ester,

- 4-(2-Chloro-6-fluoro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(3-Methoxyphenyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
5 4-Methyl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Cyclopentyl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Phenyl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Pyridin-2-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
10 4-Pyrimidin-2-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Benzo[1,3]dioxol-5-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester,
4-Benzyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Cyclopentyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-(4-Fluoro-benzyl)-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
15 4-Phenyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Pyridin-2-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Benzo[1,3]dioxol-5-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
4-Methyl-1,4-diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester,
20 4-Benzyl-1,4-diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-(Tetrahydrofuran-2-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
25 4-(Tetrahydro-furan-2-ylmethyl)-piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)-phenyl ester,
4-Cyclohexylmethyl-piperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,
4-Cyclohexylmethyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
30 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,
4-(Tetrahydrofuran-2-ylmethyl)-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
4-Naphthalen-1-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
35 4-(2-Cyclohexyl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-

- phenyl ester,
4-(3-Methoxy-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-
5 phenyl ester,
4-(Tetrahydro-furan-2-ylmethyl)-piperazine-1-carboxylic acid 4-[2-(4-chloro-phenyl)-
ethylcarbamoyl]-phenyl ester,
4-(3,4-Dichloro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
10 4-Cyclopropylmethyl-[1,4]diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-(2-Pyridin-2-yl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-(Pyrazin-2-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl es-
15 ter,
4-(Benzo-isothiazol-3-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-(5-Chloro-thiophen-2-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
20 4-(3-Trifluoromethyl-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester,
4-(5-Chloro-2-methyl-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl ester,
4-(1-Methyl-piperidin-4-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
25 yloxy)-phenyl ester,
4-Biphenyl-4-ylmethyl-[1,4]diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester
4-(5-Dimethylamino-naphthalene-1-sulfonyl)-piperazine-1-carboxylic acid 4-(5-
trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
30 4-(3-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(3-Fluoro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl
ester,
4-(3-Trifluoromethyl-pyridin-2-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-
35 yloxy)-phenyl ester,

- 4-(3-Fluorobenzyl)-piperazine-1-carboxylic acid 4-(4,6-dimethyl-pyrimidin-2-ylsulfanyl)-phenyl ester,
- 5-(4-Trifluoromethoxybenzyl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl ester,
- 5 4-(2,4-Dimethoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 5-Benzyl-2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl ester,
- 4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester,
- 10 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
- 4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
- 4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
- 15 4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,
- 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,
- 4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,
- 20 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester,
- 4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester,
- 4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester,
- 25 4-(2-Pyridin-2-yl-acetyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- Piperazine-1,4-dicarboxylic acid tert-butyl ester 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
- 30 Piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester hydrochloride,
- 4-(2-Pyridin-2-yl-acetyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester, and
- 4-(2-Pyridin-4-yl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

Further examples of specific compounds of the invention are :

- Morpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 3-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester,
5 Morpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-ylamino)-phenyl ester,
Morpholine-4-carboxylic acid 4-(3,5-dichloro-pyridin-4-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-(2-cyano-5-trifluoromethyl-pyridine-3-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 2-benzenesulfonyl-4-(3-chloro-5-trifluoromethyl-pyridine-2-
10 yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-tert-butoxy-phenyl ester,
Morpholine-4-carboxylic acid 3-(4-fluorobenzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 4-phenoxy-phenyl ester,
Morpholine-4-carboxylic acid 4-(4-chlorobenzoyl)-phenyl ester,
15 Morpholine-4-carboxylic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-[4-(4-chloro-phenyl)-thiazol-2-yl]-phenyl ester,
Morpholine-4-carboxylic acid 4-pyrrol-1-yl-phenyl ester,
Morpholine-4-carboxylic acid 4-imidazol-1-yl-phenyl ester,
Morpholine-4-carboxylic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-ylmethyl)-phenyl ester,
20 Morpholine-4-carboxylic acid 4-trifluoromethylsulfanyl-phenyl ester,
Morpholine-4-carboxylic acid 4-pentafluoromethyloxy-phenyl ester,
Morpholine-4-carboxylic acid 4-benzyloxy-phenyl ester,
Morpholine-4-carboxylic acid 4-benzyl-phenyl ester,
Morpholine-4-carboxylic acid 4'-cyano-biphenyl-4-yl-ester,
25 Morpholine-4-carboxylic acid 4'-bromo-biphenyl-4-yl-ester,
Morpholine-4-carboxylic acid biphenyl-4-yl-ester,
Morpholine-4-carboxylic acid 4-[3-(4-chlorophenyl)-ureido]-phenyl ester,
Morpholine-4-carboxylic acid 4-(4-nitro-phenoxy)-phenyl ester,
Morpholine-4-carboxylic acid 4-heptylsulfanyl-phenyl ester,
30 Morpholine-4-carboxylic acid 4-butoxy-phenyl ester,
Morpholine-4-carboxylic acid 4-(4-chloro-benzenesulfonyl)-phenyl ester,
Morpholine-4-carboxylic acid 4-(4-chloromethyl-thiazol-2-yl)-phenyl ester
Morpholine-4-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester,
cis-Morpholine-4-carboxylic acid 4-(1,3-dioxo-octahydro-isoindol-2-yl)-phenyl ester,
35 Morpholine-4-carboxylic acid 4-(cyclohexanecarbonyl-amino)-phenyl ester,

- Morpholine-4-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester,
cis/trans-Morpholine-4-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
cis-Morpholine-4-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
5 trans-Morpholine-4-carboxylic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester,
Morpholine-4-carboxylic acid 4-(3,3-dimethyl-butyrylamino)-phenyl ester,
Morpholine-4-carboxylic acid 3-benzyl-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
10 Morpholine-4-carboxylic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl ester,
15 Morpholine-4-carboxylic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propy-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
20 ter,
Morpholine-4-carboxylic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester,
Morpholine-4-carboxylic acid 4-(5,7-bis-trifluoromethyl-[1,8]naphthypyridin-2-yloxy)-phenyl ester,
25 Morpholine-4-carboxylic acid 4-pyrrolidine-1-yl-phenyl ester,
Morpholine-4-carboxylic acid 4-piperidine-1-yl-phenyl ester,
Morpholine-4-carboxylic acid 4-morpholine-1-yl-phenyl ester,
Morpholine-4-carboxylic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
Morpholine-4-carboxylic acid 4-[(6-chloro-pyridine-3-carbonyl)-amino]-phenyl ester,
30 Morpholine-4-carboxylic acid 4-[(pyridine-2-carbonyl)-amino]-phenyl ester,
2,6-dimethyl-morpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,
Morpholine-4-carboxylic acid pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 3-bromo-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 4-bromo-pyrazol-1-yl ester,
35 Morpholine-4-carboxylic acid 5-bromo-pyrazol-1-yl ester,

- Morpholine-4-carboxylic acid 3,4,5-tribromo-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 4-chloro-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester,
5 Morpholine-4-carboxylic acid 3-(2-methoxy-phenyl)-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 3-pyridin-2-yl-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 4-phenylsulfanyl-pyrazol-1-yl ester,
10 Morpholine-4-carboxylic acid 3-thiophen-2-yl-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid 5-thiophen-2-yl-pyrazol-1-yl ester,
Morpholine-4-carboxylic acid imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-chloro-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-bromo-imidazol-1-yl ester,
15 Morpholine-4-carboxylic acid 2-iodo-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-methyl-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-phenylsulfanyl-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-(4-fluoro-phenyl)-imidazol-1-yl ester,
20 Morpholine-4-carboxylic acid 2-thiophen-2-yl-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-pyridin-2-yl-imidazol-1-yl ester
Morpholine-4-carboxylic acid 2,5-dichloro-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 4-bromo-2,5-dichloro-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 4-bromo-2-chloro-imidazol-1-yl ester,
25 Morpholine-4-carboxylic acid 5-(4-methoxy-phenyl)-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 5-(4-fluoro-phenyl)-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 5-thiophen-2-yl-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 5-pyridin-2-yl-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 4-trifluoromethyl-pyrimidin-2-yl ester,
30 Morpholine-4-carboxylic acid 4-trifluoromethyl-pyrimidin-2-yl ester,
Morpholine-4-carboxylic acid imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-bromo-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-chloro-imidazol-1-yl ester,
Morpholine-4-carboxylic acid 2-phenylsulfanyl-imidazol-1-yl ester,
35 Morpholine-4-carboxylic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester,

- Morpholine-4-carboxylic acid 4-bromo-pyrazol-1-yl ester,
 Morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 Morpholine-4-carboxylic acid 3,4,5-tribromo-pyrazol-1-yl ester,
 Morpholine-4-carboxylic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester,
 5 Morpholine-4-carboxylic acid 3-thiophen-2-yl-pyrazol-1-yl ester,
 Morpholine-4-carboxylic acid pyrazol-1-yl ester, and
 1-Oxo-1*A*⁴-thiomorpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

Further examples of specific compounds of the invention are :

- 10 Methyl-o-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester,
 Methyl-m-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester, Methyl-p-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester, (3-Chloro-phenyl)-methyl-carbamic acid 4-iodo-pyrazol-1-yl ester, (3-Fluoro-phenyl)-methyl-carbamic acid 4-iodo-pyrazol-1-yl ester, 4-(3-Trifluoromethyl-pyridin-2-yl)-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 15 2,6-Dimethyl-morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 Thiomorpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 3,5-Dimethyl-morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 Piperidine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester,
 Methyl-o-tolyl-carbamic acid 2-chloro-imidazol-1-yl ester,
 20 (3-Fluoro-phenyl)-methyl-carbamic acid 2-chloro-imidazol-1-yl ester, and
 Methyl-phenyl-carbamic acid 5-phenylsulfanyl-pyrazol-1-yl ester.

Further examples of compounds of the invention are :

- N-Methyl-N-phenyl-5-hexylsulfanyl-3-p-tolyl-[1,2,4]triazole-1-carboxamide,
 25 N-Methyl-N-phenethyl-5-ethyl-3-(4-chlorophenyl)-[1,2,4]triazole-1-carboxamide,
 [3-(4-Chlorophenyl)-5-methylsulfanyl-[1,2,4]triazol-1-yl]-morpholin-4-yl-methanone,
 N,N-Dimethyl-5-methylsulfanyl-3-naphthalen-2-yl-[1,2,4]triazole-1-carboxamide,
 N,N-Dimethyl-3-(4-chloro-phenyl)-5-ethylsulfanyl-[1,2,4]triazole-1-carboxamide, and
 N,N-Dimethyl-3-biphenyl-4-yl-5-methylsulfanyl-[1,2,4]triazole-1-carboxamide.

30

The present invention also encompasses compounds of formulae I-XXXXVIII, which possess a range of pharmaceutically desirable properties.

In one embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have a solubility in water of at least 0.5 mg/L, preferably at least 2 mg/L, more prefer-

able at least 10 mg/L, more preferable at least 50 mg/L and most preferable at least 200 mg/L as determined at 25 °C and pH 7.0.

In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have a solubility in water of at least 0.5 mg/L, preferably at least 2 mg/L, more preferable at least 10 mg/L, more preferable at least 50 mg/L and most preferable at least 200 mg/L as determined at 25 °C and pH 2.0.

In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have an IC_{50} value of no greater than 5 μ M as determined by the assay 3190.2 or 3180.1 disclosed herein.

In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have an IC_{50} value of less than 1 μ M, preferably less than 500 nM, preferably less than 100 nM, preferably less than 50 nM, more preferable less than 25 nM, more preferable less than 10nM and even more preferable less than 5 nM as determined by the assay 3190.2 or 3180.1 disclosed herein.


In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which are ionized at pH 7.0.

In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have a pK_a in the range from 8 to 12, preferable from 9 to 12, more preferable from 10 to 12.

In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have a molar weight of no greater than 1000 D.


In another embodiment, the invention is concerned with compounds of formulae I-XXXXVIII, which have a molar weight of less than 750 D, preferably less than 500 D, more preferable less than 400 D, more preferable less than 300D and even more preferably less than 250 D.

In another aspect the invention is concerned with a process for the preparation of a compound of formulae I-XXXXVIII or their pharmaceutically acceptable salts, which process comprises reacting the appropriate alcohol, $Rz-OH$, with the appropriate carbamoylating reagent, $Lv-C(=O)-NRxRy$, in a solvent according to the reaction scheme P_1




and isolating the disubstituted carbamate product.

In one embodiment, the invention is concerned with process P₁, wherein said carbamoylating reagent



is selected from the group consisting of




5

In another embodiment, the invention is concerned with the process of scheme P₁, wherein said solvent is selected from the group consisting of tetrahydrofurane, dimethylformamide and N-methylpyrrolidone.

In another embodiment, the invention is concerned with the process of scheme P₁, wherein said base is selected from the group consisting of triethylamine, N,N-diisopropyl-N-ethylamine and DABCO.

15

In another aspect the invention is concerned with a process for the preparation of a compound according to the general formula



wherein R¹ is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

20

R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is op-

25

tionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

R³ is selected from hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and


X is O or S; or

a compound according to any one of formulae III-XXXVIII; or

a pharmaceutically acceptable salt thereof;

5

said process comprising the treatment of the appropriate amine, R^1-NH-R^2 , with the appropriate acylating reagent, $Y-C(=X)-R^3$, in a solvent and in the presence of a base according to the reaction scheme P_2



10

In one embodiment, the invention is concerned with the process of scheme P_2 , wherein Y is Cl.

In another embodiment, the invention is concerned with the process of scheme P_2 , wherein R^3 is an aryloxy group.

15 In another embodiment, the invention is concerned with the process of scheme P_2 , wherein said solvent is selected from the group consisting of diethyl ether, tetrahydrofuran and dichloromethane.

In another embodiment, the invention is concerned with the process of scheme P_2 , wherein said base is selected from the group consisting of trimethylamine, triethylamine, ethyl-

20 diisopropyl-amine and 1,4-diazabicyclo[2.2.2]octane.

In another embodiment, the invention is concerned with the process of scheme P_2 , wherein said base is present as a functionality in one or both of the substituents R^1 and R^2 , thus forming a salt with the acid H-Y.

25 In another aspect the invention is concerned with a pharmaceutical composition comprising a compound of formulae I-XXXVIII or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

In one embodiment the invention is concerned with a pharmaceutical composition, wherein said composition is in unit dosage form, comprising from about 0.05 to about 2000 mg,

30 preferably from about 0.1 to about 500 mg and even more preferable from about 1.0 to about 100 mg of a compound of formulae I-XXXVIII or a pharmaceutically acceptable salt thereof.

In another embodiment the invention is concerned with a pharmaceutical composition for use as a medicament for inhibiting the lipolytic activity of hormone-sensitive lipase against

triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters, said composition comprising a compound of any one of formulae I-XXXXVIII or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

- 5 In another embodiment the invention is concerned with a pharmaceutical composition for oral, nasal, transdermal, pulmonal, or parenteral administration.

In another aspect the invention is concerned with use of a compound according to any one of formulae I-XXXXVIII for preparation of a medicament for inhibition of the lipolytic activity of
10 hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters.

In one embodiment the invention is concerned with said use, wherein a further antidiabetic, antiobesity, antihypertensive or appetite regulating drug is used.

- 15 In another aspect the invention is concerned with use of a compound according to any one of formulae I-XXXXVIII for the preparation of a medicament for the treatment of any disorder where it is desirable to

modulate the plasma level of free fatty acids, glycerol, LDL-cholesterol, HDL-cholesterol, in-
20 sulin and/or glucose; and/or

modulate intracellular triacylglycerol and cholesterol ester stores, intracellular level of fatty acids, fatty acid esters such as diacylglycerols, phosphatidic acids, long chain acyl-CoA's as well as citrate or malonyl-CoA; and/or

25 increase insulin sensitivity in adipose tissue, skeletal muscle, liver or pancreatic β cells; and/or

modulate insulin secretion from pancreatic β cells.

- 30 In one embodiment the invention is concerned with said use, wherein said disorder is selected from the group consisting of insulin resistance, diabetes type 1 and 2, metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

In another aspect the invention is concerned with use of a compound according to any one of formulae I-XXXXVIII or a pharmaceutically acceptable salt thereof for the preparation of a medicament.

- 5 In another aspect the invention is concerned with a method of treating a disorder of a patient where modulation of the activity of hormone-sensitive lipase is desired, the method comprising administering to said patient an effective amount of a compound according to any one of formulae I-XXXXVIII or a pharmaceutically acceptable salt thereof.

10 In one embodiment the invention is concerned with said method, wherein said administration is carried out by the oral, nasal, transdermal, pulmonal, or parenteral route.

In another embodiment the invention is concerned with said method, wherein said disorder is selected from the group consisting of insulin resistance, diabetes type 1 and 2, metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

- 15 In another embodiment the invention is concerned with said method, wherein a further antidiabetic, antiobesity, antihypertensive or appetite regulating drug is administered to the patient.

The present invention also encompasses pharmaceutically acceptable salts of the present
20 compounds. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable base addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, nitric acids and the like. Representative exam-
25 ples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, sul-
30 phates, nitrates, phosphates, perchlorates, borates, acetates, benzoates, hydroxynaphthoates, glycerophosphates, ketoglutarates and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977, 66, 2, which is incorporated herein by reference. Examples of metal salts include lithium, sodium, potassium, magnesium, zinc, calcium salts and
35 the like. Examples of amines and organic amines include ammonium, methylamine, di-

methylamine, trimethylamine, ethylamine, diethylamine, propylamine, butylamine, tetramethylamine, ethanolamine, diethanolamine, triethanolamine, meglumine, ethylenediamine, choline, N,N'-dibenzylethylenediamine, N-benzylphenylethylamine, N-methyl-D-glucamine, guanidine and the like. Examples of cationic amino acids include lysine, arginine, histidine and the like.

The pharmaceutically acceptable salts are prepared by reacting the compound of formulae I-XXXXVIII with 1 to 4 equivalents of a base such as sodium hydroxide, sodium methoxide, sodium hydride, potassium t-butoxide, calcium hydroxide, magnesium hydroxide and the like, in solvents like ether, THF, methanol, t-butanol, dioxane, isopropanol, ethanol etc. Mixture of solvents may be used. Organic bases like lysine, arginine, diethanolamine, choline, guanidine and their derivatives etc. may also be used. Alternatively, acid addition salts wherever applicable are prepared by treatment with acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, p-toluenesulphonic acid, methanesulfonic acid, acetic acid, citric acid, maleic acid, salicylic acid, hydroxynaphthoic acid, ascorbic acid, palmitic acid, succinic acid, benzoic acid, benzenesulfonic acid, tartaric acid and the like in solvents like ethyl acetate, ether, alcohols, acetone, THF, dioxane etc. Mixture of solvents may also be used.

The stereoisomers of the compounds forming part of this invention may be prepared by using reactants in their single enantiomeric form in the process wherever possible or by conducting the reaction in the presence of reagents or catalysts in their single enantiomer form or by resolving the mixture of stereoisomers by conventional methods. Some of the preferred methods include use of microbial resolution, enzymatic resolution, resolving the diastereomeric salts formed with chiral acids such as mandelic acid, camphorsulfonic acid, tartaric acid, lactic acid, and the like wherever applicable or chiral bases such as brucine, (R)- or (S)-phenylethylamine, cinchona alkaloids and their derivatives and the like. Commonly used methods are compiled by Jaques et al in "Enantiomers, Racemates and Resolution" (Wiley Interscience, 1981). More specifically the compound of formula I may be converted to a 1:1 mixture of diastereomeric amides by treating with chiral amines, aminoacids, aminoalcohols derived from aminoacids; conventional reaction conditions may be employed to convert acid into an amide; the diastereomers may be separated either by fractional crystallization or chromatography and the stereoisomers of compound of formulae I-XXXXVIII may be prepared by hydrolysing the pure diastereomeric amide.

Various polymorphs of compound of general formulae I-XXXXVIII forming part of this invention may be prepared by crystallization of compound of formulae I-XXXXVIII under different conditions. For example, using different solvents commonly used or their mixtures for recrystallization.

tallization; crystallizations at different temperatures; various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Polymorphs may also be obtained by heating or melting the compound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffraction or such other techniques.

The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming active pharmacological substances. In general, such prodrugs will be functional derivatives of the present compounds, which are readily convertible in vivo into the required compound of the formulae I-XXXXVIII. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

The invention also encompasses active metabolites of the present compounds.

The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound of the formula I or any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof together with one or more pharmaceutically acceptable carriers or diluents.

Furthermore, the invention relates to the use of compounds of the general formulae I-XXXXVIII or their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts or pharmaceutically acceptable solvates thereof for the preparation of a pharmaceutical composition for the treatment and/or prevention of disorders where a decreased level of plasma FFA is desirable, such as the conditions mentioned above.

In another aspect, the present invention relates to a method of treating and/or preventing type 2 diabetes, insulin resistance, metabolic syndrome X, impaired glucose tolerance, dyslipidemia and abnormalities of lipoprotein metabolism.

In a still further aspect, the present invention relates to the use of one or more compounds of the general formulae I-XXXXVIII, or pharmaceutically acceptable salts thereof, for the preparation of a pharmaceutical composition for the treatment and/or prevention of type 2 diabetes, insulin resistance, metabolic syndrome X, impaired glucose tolerance, dyslipidemia and abnormalities of lipoprotein metabolism.

In a still further aspect, the present compounds are useful for the delaying or prevention of the progression from impaired glucose tolerance to type 2 diabetes.

In a still further aspect, the present compounds are useful for the delaying or prevention of the progression from non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes.

In another aspect, the present compounds reduce triglyceride levels and are accordingly useful for the treatment and/or prevention of ailments and disorders such as diabetes and/or obesity.

In still another aspect, the compounds of general formulae I-XXXXVIII are useful for the
5 treatment of hyperglycemia, elevated HbA_{1c} level, hyperinsulinemia, type 1.5 diabetes, latent autoimmune diabetes in adults, maturity onset diabetes, beta-cell apoptosis, hemochromatosis induced diabetes, impaired glucose tolerance, impaired fasting glucose, metabolic syndrome X, insulin resistance, impaired lipid tolerance, cystic fibrosis related diabetes, polycystic ovarian syndrome, and gestational diabetes.

10

In still another aspect, the compounds of general formulae I-XXXXVIII are useful for the treatment of obesity, dyslipidemia, diabetic dyslipidemia, hyperlipidemia, hypertriglyceridemia, hyperlipoproteinemia, hypercholesterolemia, hypertension, essential hypertension, acute hypertensive emergency, arteriosclerosis, atherosclerosis, restenosis, intermittent
15 claudication (atherosclerosis obliterans), cardiovascular disease, cardiomyopathy, cardiac hypertrophy, left ventricular hypertrophy, coronary artery disease, early coronary artery disease, heart insufficiency, exercise tolerance, chronic heart failure, mild chronic heart failure, arrhythmia, cardiac dysrhythmia, syncope, heart attack, myocardial infarction, Q-wave myocardial infarction, stroke, acute coronary syndrome, angina pectoris, unstable angina, cardiac
20 bypass reocclusion, diastolic dysfunction, systolic dysfunction, non-Q-wave cardiac necrosis, catabolic changes after surgery, acute pancreatitis, and irritable bowel syndrome

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of diabetic retinopathy, background retinopathy, preproliferative retinopathy, proliferative retinopathy, macular edema, cataracts, nephropathy, nephrotic syndrome, diabetic
25 nephropathy, microalbuminuria, macroalbuminuria, neuropathy, diabetic neuropathy, distal symmetrical sensorimotor polyneuropathy, and diabetic autonomic neuropathy.

In still another aspect, the compounds of general formulae I-XXXXVIII are useful for increasing the number of beta-cells in a patient, increasing the size of beta-cells in a patient or stimulating beta-cell proliferation, modulating beta-cell function and insulin secretion in a patient in need thereof, which method comprises administration of an effective amount of a
30 compound of formulae I-XXXXVIII to a patient in need thereof.

The compounds of the invention are also believed to be useful for reducing body weight in a patient in need thereof.

The compounds of the invention are also believed to be useful Use for weight neutral treatment of above mentioned diseases.

- 5 The compounds of the invention are also believed to be useful for redistributing fat in a patient in need thereof.

The compounds of the invention are also believed to be useful for redistributing central fat in a patient in need thereof.

- 10 The compounds of the invention are also believed to be useful for reducing or preventing central obesity.

The compounds of the invention are also believed to be useful for reducing postprandial serum lipid excursions.

The compounds of the invention are also believed to be useful for the treatment of fatty acid oxidation disorders such as MCAD.

15

In still another aspect, the compounds of general formulae I-XXXXVIII are believed to be useful for the treatment of a disease, condition or disorder wherein cholesterol is a precursor. Such diseases, conditions or disorders may relate to testosterone, e.g. male contraception, excessive testosterone levels, PCOS and prostate cancer. They may also relate to cortisol or corticotropin, e.g. Cushing disease.

20

The compounds of the invention are also believed to be useful for the treatment of cancer. Thus, the compounds of the general formulae I-XXXXVIII may be useful for the treatment of insulinoma (pancreatic islet cell tumors), e.g. malignant insulinomas and multiple insulinomas, adipose cell carcinomas, e.g. liposarcoma.

25

The compounds of the invention are also believed to be useful for the treatment of pheochromocytoma and other diseases with increased catecholamine secretion.

The compounds of the invention are also believed to be useful for the treatment of prostate cancer, e.g. adenocarcinoma.

30

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of hepatic steatosis.

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of cirrhosis.

35

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of AIDS or an AIDS related diseases, condition or disorders

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of lipodystrophy

In still another aspect, the compounds of general formulae I-XXXXVIII may be useful for the treatment of lactic acidosis.

5 In yet another aspect, the compounds of the present invention are expected to be useful for the treatment of CNS diseases, conditions or disorders.

Thus, the compound of the present invention may be used for the treatment of Parkinsons disease, Alzheimers disease, ADHD (Attention Deficit Hyperactivity Disorder), feeding disorders such as bulimia and anorexia, depression, anxiety, cognitive memory disorders, age
10 related cognitive decline, mild cognitive impairment and schizophrenia.

In yet another aspect, the compounds of the present invention may be useful for the treatment of inflammatory disorders, e.g. rheumatoid arthritis, psoriasis, systemic inflammatory response syndrome, sepsis and the like.

15

The present compounds may also be administered in combination with one or more further pharmacologically active substances eg., selected from antiobesity agents, antidiabetics, antihypertensive agents, agents for the treatment and/or prevention of complications resulting
20 from or associated with diabetes and agents for the treatment and/or prevention of complications and disorders resulting from or associated with obesity.

Thus, in a further aspect of the invention the present compounds may be administered in combination with one or more antiobesity agents or appetite regulating agents.

Such agents may be selected from the group consisting of CART (cocaine amphetamine
25 regulated transcript) agonists, NPY (neuropeptide Y) antagonists, MC4 (melanocortin 4) agonists, orexin antagonists, TNF (tumor necrosis factor) agonists, CRF (corticotropin releasing factor) agonists, CRF BP (corticotropin releasing factor binding protein) antagonists, urocortin agonists, $\beta 3$ agonists, MSH (melanocyte-stimulating hormone) agonists, MCH (melanocyte-concentrating hormone) antagonists, CCK (cholecystokinin) agonists, serotonin
30 re-uptake inhibitors, serotonin and noradrenaline re-uptake inhibitors, mixed serotonin and noradrenergic compounds, 5HT (serotonin) agonists, bombesin agonists, galanin antagonists, growth hormone, growth hormone releasing compounds, TRH (thyreotropin releasing hormone) agonists, UCP 2 or 3 (uncoupling protein 2 or 3) modulators, leptin agonists, DA
agonists (bromocriptin, dorexin), lipase/amylase inhibitors, RXR (retinoid X receptor) modu-
35 lators or TR β agonists.

In one embodiment of the invention the antiobesity agent is leptin.

In another embodiment the antiobesity agent is dexamphetamine or amphetamine.

In another embodiment the antiobesity agent is fenfluramine or dexfenfluramine.

In still another embodiment the antiobesity agent is sibutramine.

5 In a further embodiment the antiobesity agent is orlistat.

In another embodiment the antiobesity agent is mazindol or phentermine.

Suitable antidiabetics comprise insulin, exendin-4, GLP-1 (glucagon like peptide-1) derivatives such as those disclosed in WO 98/08871 to Novo Nordisk A/S, which is incorporated herein by reference as well as orally active hypoglycaemic agents.

10 The orally active hypoglycaemic agents preferably comprise sulphonylureas, biguanides, meglitinides, glucosidase inhibitors, glucagon antagonists such as those disclosed in WO 99/01423 to Novo Nordisk A/S and Agouron Pharmaceuticals, Inc., GLP-1 agonists, potassium channel openers such as those disclosed in WO 97/26265 and WO 99/03861 to Novo Nordisk A/S which are incorporated herein by reference, DPP-IV (dipeptidyl peptidase-IV)
15 inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, compounds modifying the lipid metabolism such as antihyperlipidemic agents and antilipidemic agents as HMG CoA inhibitors (statins), compounds lowering food intake, RXR agonists and agents acting on the ATP-dependent potassium channel of the β -cells.

20 In one embodiment of the invention the present compounds are administered in combination with insulin.

In a further embodiment the present compounds are administered in combination with a sulphonylurea eg. tolbutamide, glibenclamide, glipizide or glicazide.

In another embodiment the present compounds are administered in combination with a
25 biguanide eg. metformin.

In yet another embodiment the present compounds are administered in combination with a meglitinide eg. repaglinide or senaglinide.

In a further embodiment the present compounds are administered in combination with an α -glucosidase inhibitor eg. miglitol or acarbose.

30 In another embodiment the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the β -cells eg. tolbutamide, glibenclamide, glipizide, glicazide or repaglinide.

Furthermore, the present compounds may be administered in combination with nateglinide.

In still another embodiment the present compounds are administered in combination with an antihyperlipidemic agent or antilipidemic agent eg. cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

In a further embodiment the present compounds are administered in combination with more than one of the above-mentioned compounds eg. in combination with a sulphonylurea and metformin, a sulphonylurea and acarbose, repaglinide and metformin, insulin and a sulphonylurea, insulin and metformin, insulin, insulin and lovastatin, etc.

Furthermore, the present compounds may be administered in combination with one or more antihypertensive agents. Examples of antihypertensive agents are β -blockers such as alprenolol, atenolol, timolol, pindolol, propranolol and metoprolol, ACE (angiotensin converting enzyme) inhibitors such as benazepril, captopril, alatriopril, enalapril, fosinopril, lisinopril, quinapril and ramipril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and α -blockers such as doxazosin, urapidil, prazosin and terazosin. Further reference can be made to Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

It should be understood that any suitable combination of the compounds according to the invention with one or more of the above-mentioned compounds and optionally one or more further pharmacologically active substances are considered to be within the scope of the present invention.

The present invention also relates to processes according to reaction schemes P₁ and P₂ for the preparation of the above said novel compounds, their derivatives, their analogs, their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts or pharmaceutically acceptable solvates.

Pharmaceutical compositions.

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995. The compositions may appear in conventional forms, for example capsules, tablets, aerosols, solutions, suspensions or topical applications.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well-known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous and non-aqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

The therapeutic dose of the compound will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art. The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. In one embodiment the composition in unit dosage form, comprises from about 0.05 to about 2000 mg, preferably from about 0.1 to about 500 mg of the compound of formula I pharmaceutically acceptable salt thereof.

In a still further embodiment the pharmaceutical composition is for oral, nasal, transdermal, pulmonal, or parenteral administration.

- 5 For parenteral routes, such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having
10 the utility of a free base. When a compound of the invention contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of the compound with a chemical equivalent of a pharmaceutically acceptable acid, for example, inorganic and organic acids. Representative examples are mentioned above. Physiologically acceptable salts of a compound with a hydroxy group include the anion of said compound in combination with a suitable cation such as sodium or ammonium ion.
15

For parenteral administration, solutions of the present compounds in sterile aqueous solution, aqueous propylene glycol or sesame or peanut oil may be employed. Such aqueous solutions should be suitable buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.
20

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution
25 and various organic solvents. Examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, peanut oil, olive oil, gelatine, lactose, terra alba, sucrose, cyclodextrin, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, silicic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, polyoxyethylene, hydroxymethylcellulose and polyvinylpyrrolidone. Similarly, the carrier or diluent may include
30 any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavouring agents.

The pharmaceutical compositions formed by combining the compounds of the invention and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

5

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. These formulations may be in the form of powder or granules, as a solution or suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion.

10

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g.

15

If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

A typical tablet which may be prepared by conventional tableting techniques may contain:

Core:

	Active compound (as free compound or salt thereof)	5 mg
	Colloidal silicon dioxide (Aerosil)	1.5 mg
	Cellulose, microcryst. (Avicel)	70 mg
5	Modified cellulose gum (Ac-Di-Sol)	7.5 mg
	Magnesium stearate	q.s.

Coating:

	HPMC approx.	9 mg
10	*Mywacett 9-40 T approx.	0.9 mg

*Acylated monoglyceride used as plasticizer for film coating.

15 The compounds of the invention may be administered to a patient which is a mammal, especially a human in need thereof. Such mammals include also animals, both domestic animals, e.g. household pets, and non-domestic animals such as wildlife.

20 In a further aspect of the invention the present compounds may be administered in combination with further pharmacologically active substances e.g. an antidiabetic or other pharmacologically active material, including other compounds for the treatment and/or prevention of insulin resistance and diseases, wherein insulin resistance is the pathophysiological mechanism.

25 Furthermore, the compounds according to the invention may be administered in combination with antiobesity agents or appetite regulating agents.

EXAMPLES.**General Methods**

5 All reactions involving air-sensitive reagents were performed under nitrogen using syringe-septum cap techniques. The glassware were dried by heating with a heat-gun. MgSO_4 were used to dry solutions. Solvents were removed *in vacuo* by rotary evaporation. Melting points were recorded on a Büchi 535, Bruker AMX 400 and Bruker DRX 300 instruments were used
10 to record ^1H NMR spectra at 400 and 300 MHz respectively with tetramethylsilane (TMS) as internal standard. Coupling constants (*J*) are given in Hz.

Materials

Test compounds were synthesized or when commercially available they were purchased
15 from Specs, Maybridge, Comgenex, Peakdale or Bionet. For the synthesized compounds the procedure for synthesis and measured characteristics of the compound are stated in the example. All compounds for which no synthesis procedure is stated in the examples are commercially available and have been purchased, or were prepared by standard methods described in the literature.

20 *N*-methyl-phenethylcarbamoyl chloride was prepared from *N*-methyl-phenethylamine and phosgene using triethylamine as a base in dichloromethane. 1-Methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide, 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide and 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide
25 was prepared as described by Batey, R.A., Tetrahedron Lett. 39, 1998, 6267.

1-Hydroxypyrazole was prepared as described in Begtrup, Vedsø, J. Chem. Soc. Perkin Trans 1, 1995, 243. 1-hydroxy-4-bromopyrazole was prepared as described in Balle et al., J. Org. Chem. 64, 1999, 5366. 1-hydroxy-3-(4-methoxyphenyl)pyrazole was prepared as described in Eskildsen et al., J. Org. Chem. 2001 (in press). 1-hydroxyimidazole was prepared
30 as described in Eriksen et al., J. Org. Chem. 63, 1998, 12. 1-hydroxy-1,2,3-triazole was prepared as described in Uhlmann et al., J. Org. Chem. 62, 1997, 9177.

2-Piperidin-4-ylmethyl-1,2,3,4-tetrahydro-isoquinoline, cyclohexyl-methyl-piperidin-4-ylmethyl-amine, methyl-phenethyl-piperidine-4-ylmethyl-amine, ethyl-piperidin-4-ylmethyl-
35

piperidin-4-ylmethylamine, benzyl-methyl-piperidin-4-ylmethyl-amine, benzyl-ethyl-piperidin-4-ylmethyl-amine, methyl-piperidin-4-ylmethyl-piperidin-3-ylmethyl-amine, 1-piperidin-4-ylmethyl-piperidin-4-ol, 2-piperidin-4-ylmethyl-2,3-dihydro-1H-isoindole, cyclopropylmethyl-piperidin-4-ylmethyl-amine was prepared from 4-formylpiperidine-1-carboxylic acid *tert*-butyl ester prepared as described by Ting, P. C. (Bioorg, Med. Chem. Lett, 11, 4, 491, 2001) and an appropriate amine by a reductive amination (general procedure 19).

Benzylpiperidine-4-yl-amine, methyl-piperidin-4-yl-(2-pyridin-2-yl-ethyl)-amine, cyclohexyl-methyl-piperidin-4-yl-amine, Isopropyl-methyl-piperidin-4-yl-amine, methyl-phenethyl-piperidin-4-yl-amine, methyl-piperidin-4-yl-pyridin-3-ylmethyl-amine, was prepared from 4-oxopiperidine-1-carboxylic acid *tert*-butyl ester by a standard reductive amination procedure as described by Mattson R. J. (J. Org. Chem. 55, 2552, 1990).

Cyclopropyl-piperidin-4-yl-pyridin-4-ylmethyl-amine was prepared from 4-(cyclopropyl-pyridin-4-ylmethyl-amino)-piperidine-1-carboxylic acid *tert*-butyl ester by a classical N-deprotection reaction (HCl (g) in diethyl ether or ethanol). 4-(Cyclopropyl-pyridin-4-ylmethyl-amino)-piperidine-1-carboxylic acid *tert*-butyl ester was prepared from 4-cyclopropylamino-piperidine-1-carboxylic acid *tert*-butyl ester and pyridine-4-yl-acetaldehyde by a classical reductive amination procedure as described by Mattson R. J. 4-Cyclopropylamino-piperidine-1-carboxylic acid *tert*-butyl ester was prepared from cyclopropylamine and 4-oxopiperidine-1-carboxylic acid *tert*-butyl ester by a standard reductive amination procedure as described by Mattson R. J. Cyclopropyl-(2-fluoro-benzyl)-piperidin-4-yl-amine, cyclopropyl-piperidin-4-yl-pyridin-3-ylmethyl-amine, cyclopropylmethyl-piperidin-4-yl-pyridin-3-ylmethyl-amine and cyclopropylmethyl-piperidin-4-yl-pyridin-4-ylmethyl-amine was prepared by a procedure similar to the one described for cyclopropyl-piperidin-4-yl-pyridin-4-ylmethyl-amine.

25

Chloroformates were synthesized from the appropriate phenols and phosgene or a phosgene substitute like e.g. trichloromethyl chloroformate as described in Konakahara, Ozaki, Sato, Gold, Synthesis 1993 (1) 103-106, except that the crude product was separated from the diisopropylethylamine hydrochloride by extraction with diethyl ether rather than with THF.

30

Non-commercial N-monosubstituted piperazines were prepared by alkylation (alkylation procedure as described in e.g. Masaguer, Ravina, Tetrahedron Lett. 1996, 37 (29) 5171-5174) of 1-Boc-piperazine, and subsequent removal of the Boc group under acidic conditions, e.g.

by heating in a mixture of hydrochloric acid and ethanol. N-Monosubstituted homopiprazines and N-monosubstituted 2,5-diazabicyclo[2.2.1]heptanes were prepared in a similar manner.

Thin layer chromatography was performed on Merck DC-Alufolien, silica gel 60 F₂₅₄ and
5 components were visualized by UV₂₅₄. Flash chromatography was performed using silica gel Merck 60 size 0.04-0-063 mm and a Quad 12/25 flash system.

Preparative HPLC (Method A).

The system consists of two Gilson 322 pumps equipped with 30 ml pump heads. A Gilson
10 215 combined autoinjector and fraction collector performs injection and fraction collection. Detection is performed with a Gilson Diode array detector. Separation is performed on Waters Xterra columns 19.8 mm * 100 mm, flow rate 25 ml/min. The most widely used gradient starts at 10 % acetonitrile in water and ends after 11 min on 100 % acetonitrile, the system is buffered by 0.01% TFA. In special cases the gradient is al-
15 tered to fit the separation need.

Preparative HPLC (Method B)

HPLC Purification:

The following instrumentation is used:

Gilson 306 Pump
20 Gilson 806 Manometric module
Gilson 811C Dynamic mixer
Gilson UV/VIS-155
Gilson 202 Fraction collector

25 The instrument is controlled by Gilson Unipoint software.
The HPLC pump is connected to two eluent reservoirs containing:
A: 0.01% TFA in water
B: 0.01% TFA in acetonitrile

30 The purification is performed at room temperature by injecting an appropriate volume of the sample (preferably 2 ml) onto the column, which is eluted with a gradient of acetonitrile.

The HPLC conditions and detector settings used are as follows :

Column: Waters Xterra MS C-18 X 19 x 100 mm
Gradient: 50% - 60% acetonitrile linearly during 12 min at 20 ml/min
Detection: 210 and 270 nm

5 Preparative HPLC (method C)

The system consists of two Gilson 322 pumps equipped with 30 ml pump heads. Gilson manometric module 805. A Gilson 215 combined autoinjector and fraction collector performs injection and fraction collection. Detection is performed with a Gilson Diode array detector 170. A sample contain 25-100 mg of material dissolved in 0.5-2.0 ml of solvent (minimum water concentration: 10%).

Separation is performed on Waters Xterra, RP₁₈ 7 μ m, columns 19 mm x 150 mm, flow rate 15 ml/min (sample added with a flow rate of 5 ml/min for about 1 min). The most widely used gradient starts at 5 % acetonitrile in water and ends after 14 min on 95 % acetonitrile. This concentration is maintained for 6 min. The system is buffered with 0.05% TFA. In special cases the gradient is altered to fit the separation need. The pooled fractions are evaporated to dryness in vacuo.

HPLC-MS.

The following instrumentation was used:

- Hewlett Packard series 1100 G1312A Bin Pump
- Hewlett Packard series 1100 Column compartment
- Hewlett Packard series 1100 G13 15A DAD diode array detector
- Hewlett Packard series 1100 MSD

The instrument was controlled by HP Chemstation software.

The HPLC pump was connected to two eluent reservoirs containing:

A: 0.01% TFA in water
B: 0.01% TFA in acetonitrile

The analysis was performed at 40 °C by injecting an appropriate volume of the sample (preferably 1 μ l) onto the column, which is eluted with a gradient of acetonitrile.

The HPLC conditions, detector settings and mass spectrometer settings which were used are as follows :

Column	Waters Xterra MS C-18 X 3 mm id
Gradient	10% - 100% acetonitrile linear during 7.5 min at 1.0ml/min
Detection	210 nm (analogue output from DAD)
MS	Ionisation mode API-ES, Scan 100-1000 amu step 0.1 amu

5

General procedure 1

The phenol (1.0 mmol) was dissolved in tetrahydrofuran (15 ml) in a glass screw cap vessel, 1,4-diazabicyclo[2.2.2]octane (DABCO) (2.0 mmol) was added together with the respective carbamoyl chloride (2.0 mmol) at room temperature. The reaction mixture was shaken for 16
10 hours and poured into ethyl acetate (20 ml) and aqueous citric acid (5%; 20 ml). The organic phase was dried and evaporated to give the crude product.

General procedure 2

The phenol (1.0 mmol) was dissolved in acetonitrile (15 ml) in a glass screw cap vessel. Triethylamine (1.0 mmol) was added together with the respective 1-methyl-3*H*-imidazol-1-
15 ium iodide (1.0 mmol) at room temperature. The reaction mixture was shaken for 16-48 hours at 80 °C, cooled to room temperature and evaporated. The evaporated reaction mixture was dissolved in dichloromethane (20 ml) and extracted with aqueous hydrogen chloride (0.1 M; 20 ml). The aqueous phase was extracted with dichloromethane (3 × 20 ml). The combined organic phases were dried and evaporated to give the crude product.

20 General procedure 3

The respective phenol (1.0 mmol), 3*H*-imidazol-1-ium iodide (1.0 mmol) and triethylamine (1.0 mmol) in acetonitrile (3 ml) was heated at 50 °C overnight in a closed vial. The crude product was purified by flash column chromatography (SiO₂, ethyl acetate/ heptane) yielding the respective carbamate.

25 General procedure 4

Carbonyldiimidazole (3.6 mmol) was suspended in THF (10ml) and the appropriate secondary amine (3.0 mmol) was added. The reaction mixture was refluxed for 24 to 72 hours until no traces of amine could be detected. The reaction mixture was cooled to room temperature

and the organic phase evaporated to give the crude product of high purity. The crude product was used without further purification.

General procedure 5

5 The crude imidazole carboxamide (3.0 mmol) was dissolved in acetonitrile (10 ml) and methyl iodide (12 mmol.) was added at room temperature. The reaction mixture was stirred for 24 to 48 hours before the organic phase was evaporated to give the crude product, which was used without further purification.

General procedure 6

10 The respective 1,2,4-(1H)-triazoles were prepared as described by Blaine (US 3308131).

The 1,2,3-(1H)-triazoles were carbamoylated using the following method:

15 The respective 1,2,4-(1H)-triazole (2.0 mmol) was dissolved in dimethylformamide (10 ml) in a glass screw cap vessel, 1,4-diazabicyclo[2.2.2]octane (DABCO) (5.0 mmol) was added together with the respective carbamoyl chloride (5.0 mmol) at room temperature. The reaction mixture was stirred for 16 hours, evaporated to dryness and ethyl acetate (20 ml) and aqueous citric acid (5%; 20 ml) was added. The phases were separated and the aqueous phase extracted with ethyl acetate (20 ml). The combined organic phases were dried and evaporated to give the crude product.

20 General procedure 7

The aryl chloroformate was prepared from the corresponding phenol, trichloromethyl chloroformate and ethyldiisopropylamine in dichloromethane according to the procedure described by T. Konakahara, T. Ozaki, K. Sato and B. Gold, Synthesis, 1993 (1) 103 - 106, except that the crude reaction mixture was used without removal of ethyldiisopropylamine hydrochloride.

25 To a stirred, freshly prepared solution of the aryl chloroformate in dichloromethane (1 mmol in 3 ml) at -15 °C was added a solution of the substituted piperazine (1 mmol) in dichloromethane (1 mL). The mixture was stirred at 0 °C for 2 - 6 h. The solvent was removed in vacuo and the solid residue was triturated with diethyl ether (3 x 5 ml), then with a minute amount of water (½ - 2 ml) to remove the ethyldiisopropylamine hydrochloride; filtered and

30 dried to give the hydrochloride of the respective piperazine-1-carboxylic acid aryl ester.

General procedure 8

To a solution of the *N*-hydroxyazole (1.0 mmol) and ethyldiisopropylamine (1.5 mmol) in CH₂Cl₂ (3 mL) was added the respective carbamoyl chloride (1.5 mmol) at room temperature. The reaction mixture was stirred for 16 hours, added CH₂Cl₂ (20 mL) and washed with aqueous citric acid (5%; 3x20 mL). The organic phase was separated, dried (MgSO₄) and evaporated to give the crude product.

General procedure 9

A solution of the substituted piperazine in diethyl ether is added to a stirred solution of an equimolar amount of the aryl chloroformate (prepared from the corresponding phenol by conventional methods) in the same solvent at 0 °C. After completion of the addition, the mixture is stirred at 0 °C for 1 hour, then for 1 more hour at room temperature. The mixture is filtered, the filter cake rinsed with diethyl ether and dried to give the hydrochloride of the respective piperazine-1-carboxylic acid aryl ester.

General procedure 10

A disubstituted amine (1.0 eq) and diisopropylethylamine (1.5 eq) was added to a dried reaction flask under nitrogen. Dichloromethane or tetrahydrofuran was added to give a 0.5 mM concentration of the amine. The appropriate aryl chloroformate (1.0 eq) (prepared from the corresponding phenol by conventional methods) was dissolved in a minimum amount of dichloromethane or tetrahydrofuran and added drop by drop at room temperature. The reaction mixture was stirred overnight, citric acid (5%) was added, and the two phases separated. The aqueous phase was extracted twice with dichloromethane, the combined organic phases were dried with MgSO₄, filtered and evaporated to give the crude product.

General procedure 11

An appropriate amine (1.0 eq.) was dissolved in dichloromethane (0.5 mM concentration of the amine) in a dried reaction flask under nitrogen. The appropriate aryl chloroformate (1.0 eq.) (prepared from the corresponding phenol by conventional methods) was dissolved in a minimum amount of dichloromethane and added drop by drop at room temperature. Heptane was added to give a 20% solution in dichloromethane and the crude product was isolated by filtration. The crude product was washed with a mixture of dichloromethane/heptane (5:1) and dried in vacuum.

General procedure 12

An appropriate amine (1.0 eq.) and diisopropylethylamine (1.0 eq.) was dissolved in tetrahydrofuran (0.5 mM concentration of the amine) in a dried reaction flask under nitrogen. The appropriate aryl chloroformate (1.0 eq.) (prepared from the corresponding phenol by conventional methods) was dissolved in a minimum amount of tetrahydrofuran and added drop by drop at room temperature. Acetic acid was added to the reaction mixture (pH 3-5) and the reaction mixture filtered. The organic phase was evaporated and the crude product subjected to preparative HPLC.

General procedure 13

4-(Methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester (1 eq.), diisopropylethylamine (1.5 eq.) was dissolved in tetrahydrofuran (50 mM of phenol). The reaction mixture was added to a mono or disubstituted amine. The reaction mixture was stirred at 50 °C for 16 hours. Citric acid (5%) and *tert*-butyl-methylether was added and the two phases separated. The organic phase was evaporated to give the crude product.

General procedure 14

The phenol (1.0 mmol) was dissolved in tetrahydrofuran (15 ml) in a glass screw cap vessel, 1,4-diazabicyclo[2.2.2]octane (DABCO) (2.0 mmol) was added together with the respective carbamoyl chloride (2.0 mmol) at room temperature. The reaction mixture was shaken for 16 hours. Acetic acid was added to the reaction mixture (pH 3-5) and the reaction mixture filtered. The organic phase was evaporated and the crude product subjected to preparative HPLC.

General procedure 15

A solution of the substituted piperazine in diethyl ether was added to a stirred solution of an equimolar amount of the aryl chloroformate in the same solvent at 0 °C. After completion of the addition, the mixture was stirred at ambient temperature for 1 - 2 hours. Stirring was discontinued and as much as possible of the solvent was removed by decantation. The residue was rinsed twice with ether by stirring and subsequent decantation and finally dried on a rotary evaporator to give the hydrochloride of the respective piperazine-1-carboxylic acid aryl ester.

If necessary, further purification was achieved by treating the crude product with a mixture of ethyl acetate and a slight excess of sodium bicarbonate (approx. 1.1 eqv.) in water, extracting the aqueous phase twice with ethyl acetate, drying the combined extracts, filtering and evaporating to give the piperazine-1-carboxylic acid aryl ester as a free base.

General procedure 16

To a solution of the *N*-hydroxyazole (1.0 mmol) and ethyldiisopropylamine (1.0 mmol) in
5 CHCl₃ (1 mL) at -30 °C was added trichloromethyl chloroformate (1.1 mmol). The solution
was stirred at -30 °C for 10 min and at room temperature for 1 h. The solution was evapo-
rated to dryness at room temperature and redissolved in CHCl₃ (2 mL) and cooled to -30 °C
before addition of the appropriate piperazine (3 mmol). The solution was allowed to warm to
room temperature over 30 min and evaporated to give the crude product.

10

General procedure 17

To a suspension of *N*-methyl-*N*-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its
TFA salt (0.5 mmol) and an aryl sulfonyl chloride (0.75 mmol) in CH₂Cl₂ (2 mL) was added
DIPEA (1.25 mmol). The reaction mixture was stirred at rt for 2-16 h, and evaporated to dry-
15 ness and redissolved in MeCN and purified by preparative HPLC (Gilson).

General procedure 18

A solution of the substituted piperazine in diethyl ether was added to a stirred solution of an
equimolar amount of the aryl chloroformate in the same solvent at 0 °C. Then an equimolar
20 amount of diisopropylethylamine (DIPEA) in diethyl ether solution was added and the mix-
ture was stirred at ambient temperature for 1 - 2 hours. The solvent was removed on a ro-
tary evaporator and the residue was partitioned between ethyl acetate and water. The aque-
ous phase was extracted twice with ethyl acetate and the combined organic phases were
dried and filtered. Removal of the solvent gave the piperazine-1-carboxylic acid aryl ester.

25

General procedure 19

4-Formyl-piperidine-1-carboxylic acid *tert*-butyl ester (1.5 g, 7.03 mmol) prepared as de-
scribed by Ting, P. C. was added to a dried screw cap vessel under nitrogen. The appropri-
ate amine (7.03 mmol) methanol (10 ml) and acetic acid 100 (μl) was added and the reaction
30 mixture was stirred for 2 h. at room temperature. Sodium cyanoborohydride (1.0 M sol. In
THF, 4.7 ml) was added during 1 minute and the mixture was stirred for 16 h. at room tem-
perature. The reaction mixture was evaporated to dryness and extracted with dichloro-
methane (3 × 75 ml) from a 10 % aqueous sodium hydrogen carbonate solution (100 ml).
The organic phases were pooled, dried, and evaporated to dryness to give the crude inter-

mediate, which was subjected to flash chromatography (ethyl acetate/heptane/methanol, 1:2:0 → 4:0:1).

A 3M solution of hydrogen chloride (50 ml) was added to the intermediate and the reaction mixture was stirred for 16 h. The reaction mixture was evaporated to dryness to give the crude product, which was dried in vacuum. The crude product was used without further purification.

General procedure 20

The arylboronic acid (1.2 mmol), KF (3.3 mmol), Pd₂(dba)₃ (0.03 mmol) and Pd(P(*t*-Bu)₃)₂ (0.06 mmol) were added to a Schlenk tube under nitrogen. The Schlenk tube was evacuated and refilled with nitrogen five times. Next the aryl halide (1.0 mmol) in THF (2 mL) was added. The reaction mixture was stirred at rt for 16 h.

General procedure 21

To a suspension of methyl-phenyl-carbamic acid 4-amino-phenyl ester (0.5 mmol), (see preparation below) and an aryl sulfonyl chloride (0.75 mmol) in CH₂Cl₂ (2 mL) was added DIPEA (1.25 mmol). The reaction mixture was stirred at rt for 2-16 h, and evaporated to dryness and redissolved in MeCN and purified by preparative HPLC (Gilson).

General procedure 22

A solution of 1-benzyloxy-4-iodobenzene (4.1 mmol) in dry THF (20 mL) was cooled to -78 °C. *n*-Butyllithium (1.57 M in hexanes, 4.1 mmol) was added during 2 min. After the mixture was stirred for another 5 min, an aryl aldehyde (4.1 mmol) was added. The mixture was allowed to warm to rt during 20 min and quenched with aqueous NaHCO₃. Extraction with CH₂Cl₂, drying (MgSO₄), filtration and evaporation provided the crude diarylmethanols which were recrystallised from EtOAc-heptane. A solution of the diarylmethanol product (2 mmol), NaI (14 mmol) in dry MeCN (20 mL) was added trimethylsilyl chloride (14 mmol) and stirred at 80 °C for 19 h. The purple reaction mixture was evaporated to dryness and treated with an aqueous solution of Na₂SO₃. The 4-arylmethylphenols were isolated by filtration or after extraction with CH₂Cl₂ and subsequent purification by flash chromatography (Quad flash 12, EtOAc-heptane).

General procedure 23

A solution of the sulfonamide (0.2 mmol), 37% aqueous formaldehyde (0.5 mL), and TFA (2 mL) was heated in a closed vessel in a Smith Creator microwave oven for 300 s at 150 °C. The crude product was evaporated to dryness and purified by preparative HPLC (Gilson).

5

General procedure 24

A suspension of the phenol (1.0 mmol), 1,4-diazabicyclo[2.2.2]octane (DABCO) (1.5 mmol) and 3-[4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium; iodide (1.5 mmol) in CH₂Cl₂ (1 mL) was stirred at room temperature for 16 hours. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane). The purified tert-butyldimethylsilyl ether was desilylated by stirring with a 3.2 M solution of HCl in Et₂O (20 mL) for 3 h at rt.

10

General procedure 25

A solution of an alcohol (0.4 mmol), a phenol (0.4 mmol), diisopropylethylamin (0.44 mmol) and tributylphosphine (0.5 mmol) in THF (2 mL) was stirred under nitrogen at rt. ADDP (0.5 mmol) dissolved in THF (2 mL) was added and the reaction mixture was stirred at rt for 16 h, filtered, evaporated to dryness and redissolved in MeCN and purified by preparative HPLC (Gilson).

20

General procedure 26

A solution of an alcohol (0.4 mmol), a phenol/thiophenol/N-hydroxyazole/azole or imide (0.4 mmol), diisopropylethylamin (0.44 mmol) and solid supported triphenylphosphine (3 mmol/g, 1.2 mmol) in CH₂Cl₂ (2 mL) was stirred under nitrogen at rt. Di-tert-butylazodicarboxylate (DBAD, 1.2 mmol) dissolved in CH₂Cl₂ (1 mL) was added and the reaction mixture was stirred at rt for 16 h. TFA (0.5 mL) was added and the mixture was stirred for further 1 h at rt. Addition of EtOAc, filtration, followed by evaporation to dryness gave a crude which was either purified by flash chromatography (Quad flash 12, EtOAc-heptane) or redissolved in MeCN and purified by preparative HPLC (Gilson).

25

30

Preparation of 1-methyl-3H-imidazol-1-ium iodides**1-Methyl-3-(7-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carbonyl)-3H-imidazol-1-ium iodide****5 Step A: Imidazol-1-yl-(7-trifluoromethyl-3,4-dihydro-2H-quinolin-1-yl)-methanone**

The title product was prepared from 7-(trifluoromethyl)-1,2,3,4-tetrahydroquinoline, as described in the general procedure 4. Light yellow oil. HPLC-MS: m/z = 296.1 (M+1); Rt: 2.85 min.

10 δ_H (300MHz; CDCl₃): 2.11 (qi, 2H), 2.93 (t, 2H), 3.90 (t, 2H), 7.01 (s, 2H), 7.05 (s, 1H), 7.34 (s, 1H), 7.34 (d, 1H), 7.77 (s, 1H).

Step B: 1-Methyl-3-(7-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carbonyl)-3H-imidazol-1-ium iodide

15 The title product was prepared from imidazol-1-yl-(7-trifluoromethyl-3,4-dihydro-2H-quinolin-1-yl)-methanone, as described in the general procedure 5. Light yellow crystals. HPLC-MS: m/z = 310.2 (M+1), Rt: 1.84 min.

δ_H (300MHz; CDCl₃): 2.01 (qi, 2H), 2.94 (t, 2H), 3.83 (t, 2H), 3.92 (s, 3H), 7.52 (s, 2H), 7.76 (s, 1H), 7.80 (s, 1H), 7.85 (s, 1H), 9.62 (s, 1H).

20

3-(cyclohexyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide**Step A: Imidazole-1-carboxylic acid cyclohexyl-methyl-amide**

25 The title product was prepared from cyclohexyl-methyl-amine, as described in the general procedure 4. Off-white crystals. HPLC-MS: m/z = 208.1 (M+1), Rt: 1.85 min

δ_H (300MHz; CDCl₃): 1.05-1.22 (m, 1H), 1.25-1.45 (m, 2H), 1.50-1.61 (dt, 2H), 1.63-1.75 (d, 1H), 1.76-1.95 (m, 4H), 3.80-3.95 (m, 1H), 7.09 (bs, 1H), 7.21 (bs, 1H), 7.87 (bs, 1H).

Step B: 3-(cyclohexyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

The title product was prepared from imidazole-1-carboxylic acid cyclohexyl-methyl-amide, as described in the general procedure 5. Yellow crystals; HPLC-MS: $m/z = 222.2$ ($M+1$), Rt: 1.15 min.

δ_H (300MHz; $CDCl_3$): 1.03-1.25, (m, 1H), 1.30-1.60 (m, 4H), 1.62-1.1.78 (m, 1H), 1.82-2.00 (t, 4H), 3.21 (s, 3H), 3.90-4.10 (m, 1H), 4.29 (s, 3H), 7.52 (bs, 1H), 7.66 (bs, 1H), 10.37 (bs, 1H).

3-(2,6-dimethyl-morpholine-4-carbonyl)-1-methyl-3H-imidazol-1-ium iodide

10 Step A: (2,6-Dimethyl-morpholin-4-yl)-imidazol-1-yl-methanone

The title product was prepared from 2,6-dimethyl-morpholine, as described in the general procedure 4. Colourless oil. PPLC-MS: $m/z = 210.10$ ($M+1$), Rt: 0.63 min.

δ_H (300MHz; $CDCl_3$): 1.20 (s, 3H), 1.22 (s, 3H), 2.82 (dd, 2H), 3.60-3.75 (m, 2H), 3.93 (d, 2H), 7.11 (s, 1H), 7.20 (s, 1H), 7.87 (s, 1H).

Step B: 3-(2,6-dimethyl-morpholine-4-carbonyl)-1-methyl-3h-imidazol-1-ium iodide

The title product was prepared from, (2,6-Dimethyl-morpholin-4-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Light yellow oil; HPLC-MS: $m/z = 224.2$ ($M+1$), Rt: 0.40 min.

δ_H (300MHz; $CDCl_3$): 1.23 (s, 3H), 1.25 (s, 3H), 2.90-3.10 (m, 2H), 3.7-3.9 (m, 2H), 3.95-4.15 (m, 2H), 4.26 (s, 3H), 7.67 (s, 1H), 7.73 (s, 1H) 10.05 (s, 1H).

3-(Benzyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

25 Step A: Imidazole-1-carboxylic acid benzyl-methyl-amide

The title product was prepared from benzyl-methyl-amine, as described in the general procedure 4. Light yellow crystals. HPLC-MS: $m/z = 216.1$ ($M+1$), Rt: 1.53 min.

δ_H (300MHz; $CDCl_3$): 3.04 (s, 3H), 4.65 (s, 3H), 7.08 (bs, 1H), 7.22-7.34 (m, 3H), 7.34-7.50 (m, 3H), 7.93 (bs, bs, 1H).

Step B: 3-(Benzyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

The title product was prepared from imidazole-1-carboxylic acid benzyl-methyl-amide, as described in the general procedure 5. Light yellow crystals. HPLC-MS m/z = 230.1 ($M+1$), Rt:

5 1.23 min.

δ_H (300MHz; $CDCl_3$): 3.25 (s, 3H), 4.2 (s, 3H), 4.76 (s, 2H), 7.27-7.50 (m, 5H), 7.56 (bs, 1H), 7.70 (bs, 1H), 10.29 (bs, 1H).

3-(Phenyl-ethyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

10

Step A: Imidazole-1-carboxylic acid phenyl-ethyl-amide

The title product was prepared from phenyl-ethyl-amine, as described in the general procedure 4. Light brown oil. HPLC-MS m/z = 216.1 ($M+1$), Rt: 1.75 min.

15 δ_H (300MHz; $CDCl_3$): 1.26 (t, 3H), 3.92 (q, 2H), 6.79 (s, 1H), 6.84 (s, 1H), 7.10 (d, 2H), 7.27-7.45 (m, 3H), 7.55 (s, 1H).

Step B: 3-(Phenyl-ethyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

The title product was prepared from imidazole-1-carboxylic acid phenyl-ethyl-amide, as described in the general procedure 5. Light yellow crystals. HPLC-MS m/z = 230.2 ($M+1$), Rt:

20 1.16 min.

δ_H (300MHz; $CDCl_3$): 1.28 (t, 3H), 3.96 (q, 2H), 4.10 (s, 3H), 7.03 (s, 1H), 7.27 (s, 1H), 7.35-7.60 (m, 6 H), 9.70 (s, 1H).

3-(2,3-Dihydro-indole-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide

25

Step A: (2,3-Dihydro-indol-1-yl)-imidazol-1-yl-methanone

The title product was prepared from Indoline, as described in the general procedure 4. Pink crystals. HPLC-MS m/z = 214.1 ($M+1$), Rt: 1.62 min.

δ_H (300MHz; CDCl₃): 3.21 (t, 2H), 4.21 (t, 2H), 7.21 (dt, 1H), 7.15 (s, 1H), 7.17-7.29 (m, 2H), 7.36 (t, 1H), 7.40 (d, 1H), 8.03 (bs, 1H).

Step B: 3-(2,3-Dihydro-indole-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide

- 5 The title product was prepared from (2,3-dihydro-indol-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Light brown crystals. HPLC-MS m/z = 228.1 (M+1), Rt: 0.94 min.

δ_H (300MHz; CDCl₃): 3.42 (t, 2H), 4.34 (s, 3H), 4.71 (t, 2H), 7.17-7.36 (m, 4H), 7.78 (bs, 1H), 7.86 (d, 1H), 10.76 (bs, 1H).

10

3[(4-Chlorophenyl)-methyl-carbamoyl]-1-methyl-3H-imidazol-1-ium iodide

Step A: Imidazole-1-carboxylic acid (4-chloro-phenyl)-methyl-amide

- 15 The title product was prepared from 4-chloro-N-methylaniline, as described in the general procedure 4. Light yellow crystals. HPLC-MS m/z = 236.1 g/mol (M+1), Rt: 1.91 min.

δ_H (300MHz; CDCl₃): 3.47 (s, 3H), 6.85 (s, 1H), 6.87 (s, 1H), 7.06 (d, 2H), 7.36 (d, 2H), 7.60 (s, 1H).

Step B: [(4-Chlorophenyl)-methyl-carbamoyl]-1-methyl-3H-imidazol-1-ium iodide

- 20 The title product was prepared from imidazole-1-carboxylic acid (4-chloro-phenyl)-methyl-amide, as described in the general procedure 5. Orange crystals. HPLC-MS m/z = 250.1 (M+1), Rt: 1.06 min.

δ_H (300MHz; CDCl₃): 3.55 (s, 3H), 4.12 (s, 3H), 7.16 (bs, 1H), 7.35 (t, 1H), 7.41 (d, 1H), 7.45 (t, 1H), 7.51 (t, 1H), 7.53-7.55 (m, 1H), 9.92 (bs, 1H).

25

3-(isopropyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

Step A: Imidazole-1-carboxylic acid isopropyl-methyl-amide

The title product was prepared from isopropyl-methylamine, as described in the general procedure 4. Light yellow oil. HPLC-MS m/z = 168.1 (M+1), Rt: 0.51 min.

δ_H (300MHz; $CDCl_3$): 1.25 (s, 3H), 1.27 (s, 3H), 2.93 (s, 3H), 4.36 (Q, 1H), 7.08 (bs, 1H), 7.22 (bs, 1H), 7.88 (bs, 1H).

5 Step B: 3-(isopropyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide

The title product was prepared from imidazole-1-carboxylic acid isopropyl-methyl-amide, as described in the general procedure 5. Light yellow crystals. HPLC-MS m/z = 182.2 (M+1), Rt: 0.41 min.

10 δ_H (300MHz; $CDCl_3$): 1.31 (s, 3H), 1.35 (s, 3H), 3.17 (s, 3H), 4.29 (s, 3H), 4.30-4.50 (m, 2H), 7.62 bs, 1H), 7.71 (bs, 1H), 10.29 (bs, 1H).

3-(1,3-Dihydroisoindole-2-carbonyl)-1-methyl-3H-imidazol-1-ium iodide

Step A: (1,3-Dihydroisoindole-1-yl)-imidazol-1-yl-methanone

15

The title product was prepared from isoindole, as described in the general procedure 4. Oil.

Step B: 3-(1,3-Dihydroisoindole-2-carbonyl)-1-methyl-3H-imidazol-1-ium iodide

20 The title product was prepared from (1,3-dihydroisoindole-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Crystals.

δ_H (300MHz; $CDCl_3$): 3.96 (s, 3H), 4.98 (s, 2H), 5.04 (s, 2H), 7.35 (bs, 3H), 7.44 (bs, 1H), 7.91 (s, 1H), 8.21 (s, 1H), 9.74 (s, 1H).

1-Methyl-3-(piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

25 Step A: Piperidin-1-yl-imidazol-1-yl-methanone

The title product was prepared from piperidine, as described in the general procedure 4. Oil.

Step B: 1-Methyl-3-(piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

The title product was prepared from piperidin-1-yl-imidazol-1-yl-methanone, as described in the general procedure 5. Oil.

5 δ_H (300MHz; CDCl₃): 1.74 (s, 6H), 3.66 (bs, 4H), 4.28 (s, 3H), 7.78 (s, 1H), 7.83 (s, 1H), 10.07 (s, 1H).

1-Methyl-3-(2-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

Step A: (2-Methyl-piperidin-1-yl)-imidazol-1-yl-methanone

10 The title product was prepared from 2-methyl-piperidine, as described in the general procedure 4. light yellow oil. HPLC-MS m/z = 194.2 (M+1), Rt: 0.92 min.

δ_H (300MHz; CDCl₃): 1.33 (d, 3H) 1.45-1.67 (m, 2H), 1.68-1.85 (m, 4H), 3.17 (dt, 1H), 3.86 (dd, 1H), 4.35-4.50 (m, 1H), 7.09 (s, 1H), 7.18 (s, 1H), 7.84 (s, 1H).

Step B: 1-Methyl-3-(2-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

15

The title product was prepared from (2-methyl-piperidin-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Orange solid. HPLC-MS m/z = 208.1 (M+1), Rt: 0.57min.

20 δ_H (300MHz; CDCl₃): 1.40 (d, H), 1.60-1.98 m, 6H), 3.45 (t, 1H), 3.90 (d, 1H), 4.30 (s, 3H), 4.45-4.60 (m, 1H), 7.59 (s, 1H), 7.62 (s, 1H), 10.06 (s, 1H).

1-Methyl-3-(3-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

Step A: (3-Methyl-piperidin-1-yl)-imidazol-1-yl-methanone

25 The title product was prepared from 3-methyl-piperidine, as described in the general procedure 4. light yellow oil. HPLC-MS m/z = 194.2 (M+1), Rt: 1.15 min.

δ_H (300MHz; CDCl₃): 0.94 (d, 3H), 1.05-1.35 (m, 1H), 1.50-2.00 (m, 4H), 2.67 (t, 1H), 3.01 (dt, 1H), 3.98 (t, 2H), 7.09 (s, 1H), 7.19 (s, 1H), 7.85 (s, 1H).

Step B: 1-Methyl-3-(3-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

The title product was prepared from (3-methyl-piperidin-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Yellow oil. HPLC-MS m/z = 208.1 ($M+1$), R_t : 0.69min. δ_H (300MHz; $CDCl_3$): 0.97 (d, 3H), 1.15-1.40 (m, 1H), 1.55-2.00 (m, 4H), 2.92 (t, 1H), 3.28 (t, 1H), 3.90-4.15 (m, 2H), 4.28 (s, 3H), 7.60-7.75 (m, 2H), 10.14 (s, 1H).

1-Methyl-3-(4-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

Step A: (4-Methyl-piperidin-1-yl)-imidazol-1-yl-methanone

The title product was prepared from 4-methyl-piperidine, as described in the general procedure 4. light yellow oil. HPLC-MS m/z = 194.2 ($M+1$), R_t : 1.32 min. δ_H (300MHz; $CDCl_3$): 1.00 (d, 3H), 1.15-1.35 (m, 2H), 1.55-1.85 (m, 3H), 3.02 (dt, 2H), 4.08 (d, 2H), 7.08 (s, 1H), 7.19 (s, 1H), 7.85 (s, 1H).

Step B: 1-Methyl-3-(4-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

The title product was prepared from (4-methyl-piperidin-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Yellow oil. HPLC-MS m/z = 208.1 ($M+1$), R_t : 0.65min. δ_H (300MHz; $CDCl_3$): 1.00 (d, 3H), 1.20-1.50 (m, 2H), 1.66-1.90 (m, 3H), 3.32 (t, 2H), 4.13 (d, 2H), 4.28 (s, 3H), 7.58 (s, 1H), 7.64 (s, 1H), 10.15 (s, 1H).

1-Methyl-3-(4-benzyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

Step A: (4-Benzyl-piperidin-1-yl)-imidazol-1-yl-methanone

The title product was prepared from 4-methyl-piperidine, as described in the general procedure 4. light yellow oil. HPLC-MS m/z = 270.2 ($M+1$), R_t : 2.58 min. δ_H (300MHz; $CDCl_3$): 1.10-1.50 (m, 2H), 1.65-2.00 (m, 3H), 2.59 (d, 2H), 2.97 (dt, 2H), 4.08 (d, 2H), 7.05-7.40 (m, 7H), 7.84 (s, 1H).

Step B: 1-Methyl-3-(4-benzyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide

The title product was prepared from (4-benzyl-piperidin-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Yellow oil. HPLC-MS m/z = 208.1 ($M+1$), R_t : 0.65min.

δ_{H} (300MHz; CDCl_3): 1.30-1.50 (m, 2H), 1.75-1.95 (m, 3H), 2.59 (d, 2H), 3.15-3.40 (m, 2H), 4.05-4.20 (m, 2H), 4.25 (s, 3H), 7.10-7.35 (m, 5H), 7.45 (bs, 1H), 7.60 (bs, 1H), 10.22 (s, 1H).

5 **1-Methyl-3-(1,2,3,4-tetrahydroisoquinoline-1-carbonyl)-3H-imidazol-1-ium iodide**

Step A: (1,2,3,4-Tetrahydroisoquinoline-1-yl)-imidazol-1-yl-methanone

The title product was prepared from 1,2,3,4-tetrahydroisoquinoline, as described in the general procedure 4. Oil.

10 Step B: 1-Methyl-3-(1,2,3,4-tetrahydroisoquinoline-1-carbonyl)-3H-imidazol-1-ium iodide

The title product was prepared from (1,2,3,4-tetrahydroisoquinoline-1-yl)-imidazol-1-yl-methanone, as described in the general procedure 5. Oil.

δ_{H} (300MHz; CDCl_3): 2.97 (t, 2H), 3.73 (bs, 2H), 3.94 (s, 3H), 4.75 (s, 2H), 7.15-7.35 (m, 4H),
15 7.88 (d, 1H), 8.09 (d, 1H), 9.63 (s, 1H).

Preparation of Phenols

20 **1-(4-Hydroxy-phenyl)-4,4-dimethyl-piperidine-2,6-dione**

A mixture of 4-aminophenol (3.27 g, 30.0 mmol) and 3,3-dimethylglutaric anhydride (4.26 g, 30.0 mmol) was heated in a round bottom flask at 165 °C for 1 h, followed by heating at 180 °C for 7 h. After cooling to room temperature the solid material was dissolved in hot ethanol,
25 activated charcoal was added and the solution was heated at reflux for 1 h. The solid material was removed by hot filtration. The solvent was evaporated and the residue was crystallised from water/ethanol yielding the title compound (3.51 g, 50% yield, pink solid).

^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 1.08 (s, 6H), 2.63 (s, 4H), 6.77 + 6.86 (AB-system, 4H), 9.56 (s, 1H).

***cis*-2-(4-Hydroxy-phenyl)-hexahydro-isoindole-1,3-dione**

A mixture of 4-aminophenol (5.46 g, 50.0 mmol) and *cis*-1,2-cyclohexanedicarboxylic anhydride (7.71 g, 50.0 mmol) was heated in a round bottom flask at 170 °C for 2 h. After cooling to room temperature the solid material was dissolved in hot ethanol (200 ml), activated charcoal was added and the solution was heated at reflux for 1 h. The solid material was removed by hot filtration. The solvent was partially evaporated. The solids were collected by filtration, washed quickly with a small amount of ethanol and dried *in vacuo* at 40 °C yielding the title compound (8.52 g, 69% yield, pink solid).

¹H NMR (300 MHz, DMSO-*d*₆): δ 1.38 (m, 4H), 1.73 (m, 4H), 3.02 (m, 2H), 6.82 (d, 2H), 7.02 (d, 2H), 9.66 (s, 1H, OH); HPLC-MS: *m/z* = 246 (M+1); R_t = 2.53 min.

Cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide

15

To a solution of 4-aminophenol (5.00 g, 45.8 mmol) in dichloromethane (50 ml) were added cyclohexanecarbonyl chloride (6.72 g, 45.8 mmol) and pyridine (3.70 ml, 45.8 mmol), while cooling the reaction mixture in an ice bath. After the addition was completed, the cooling bath was removed and stirring was continued overnight at room temperature. Water (100 ml) was added, the organic phase was removed and the resulting solution was extracted with ethyl acetate (3 x 300 ml). The combined organic phases were washed with water (2 x 200 ml), dried, filtered and evaporated, yielding an off-white solid. The crude product was purified by flash column chromatography (SiO₂, ethyl acetate/heptane (40:60)), yielding a mixture of two compounds, which were dissolved in THF. 6N NaOH (aq, 32 ml) was added and the mixture was stirred at room temperature for 2.5 h. The solution was acidified with concentrated hydrochloric acid and the organic solvent was removed by evaporation. The solid material was collected by filtration, dried and recrystallised from ethyl acetate/heptane, yielding the title compound (4.20 g, 41%, off-white solid).

¹H NMR (300 MHz, DMSO-*d*₆): δ 1.12-1.48 (m, 5H), 1.65 (m, 1H), 1.70-1.82 (m, 4H), 2.27 (m, 1H), 6.66 (d, 2H), 7.36 (d, 2H), 9.10 (s, 1H), 9.50 (s, 1H); HPLC-MS: *m/z* = 220 (M+1); R_t = 2.69 min.

2-Cyclohexyl-*N*-(4-hydroxy-phenyl)-acetamide

To a solution of 4-aminophenol (3.83 g, 35.1 mmol) in dichloromethane (50 ml) were added
5 cyclohexylacetyl chloride (11.26 g, 70.1 mmol) and pyridine (5.67 ml, 70.1 mmol), while cool-
ing the reaction mixture in an ice bath. After the addition was completed, the cooling bath
was removed and stirring was continued overnight at room temperature. The solvent was
removed and the residue was dissolved in THF (300 ml). 6N NaOH (aq, 41 ml) was added
and the mixture was stirred at room temperature for 4 h. The solution was acidified with 1 N
10 hydrochloric acid. The solvent was removed by evaporation. The solid material was collected
by filtration, dried *in vacuo* at 40 °C and dissolved in methanol (100 ml). A solution of KOH
(5.5 g) in methanol (50 ml) was added. After stirring for 1 h at room temperature water (200
ml) was added and the organic solvent was removed by evaporation. The aqueous phase
was acidified with 1 N HCl. The solid material was isolated by filtration and dried *in vacuo* at
15 40 °C yielding the title compound (6.31 g, 77% yield, pink crystals).
¹H NMR (200 MHz, DMSO-*d*₆): δ 0.82-1.32 (m, 5H), 1.54-1.76 (m, 6H), 2.12 (d, 2H), 6.66 (d,
2H), 7.32 (d, 2H), 9.12 (s, 1H), 9.57 (s, 1H); HPLC-MS: *m/z* = 234 (M+1); R_t = 3.09 min.

***cis/trans*-4-*tert*-Butyl-cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide**

20

To a solution of 4-aminophenol (3.08 g, 28.2 mmol) in dichloromethane (50 ml) were added
cis/trans-4-*tert*-butyl-cyclohexanecarbonyl chloride (11.43 g, 56.4 mmol) and pyridine (4.56
ml, 56.4 mmol), while cooling the reaction mixture in an ice bath. After the addition was com-
pleted, the cooling bath was removed and stirring was continued overnight at room tempera-
25 ture. The solvent was removed by evaporation and the residue was dissolved in THF (300
ml). 6N NaOH (aq, 33 ml) was added and the mixture was stirred at room temperature over-
night. The organic phase was removed by evaporation. Water (200 ml) was added and the
solid material was collected by filtration, washed with water, dried *in vacuo* at 40 °C and dis-
solved in methanol (100 ml). A solution of KOH (2.4 g) in methanol (50 ml) was added. After
30 stirring for 2 h at room temperature water (200 ml) was added and the organic phase was
removed by evaporation. The aqueous phase was acidified with 1 N HCl and extracted with

ethyl acetate (3 x 300 ml). The combined organic phases were washed with saturated sodium bicarbonate, dried, filtered and evaporated, yielding a pink oil, which was dried *in vacuo* at 40 °C. The solid material was crystallised from ethyl acetate/heptane yielding the title compound (2.03 g, 26%, pink crystals). From the first aqueous extract a second portion of product was isolated by extraction with ethyl acetate (3 x 250 ml). The combined organic phases were washed with water (400 ml), saturated sodium bicarbonate (2 x 400 ml), dried, filtered and evaporated, yielding a pink thick oil. Crystallisation from ethyl acetate/heptane yielded a further amount of title compound (2.75 g, 35%).

¹H NMR (300 MHz, DMSO-*d*₆): δ 0.80 + 0.84 (2 x s, 9H), 0.98 (m, 2H), 1.23-1.57 (m, 4H), 1.76-1.90 (m, 2H), 2.02-2.14 (m, 1.5H), 2.57 (m, 0.5H), 6.65 (d, 2H), 7.34 (d x d, 2H), 9.09 (s, 1H), 9.36 + 9.50 (2 x s, 1H); HPLC-MS: *m/z* = 276 (M+1); *R*_t = 4.19 and 4.27 min.

***N*-(4-Hydroxy-phenyl)-3,3-dimethyl-butyramide**

15. To a solution of 4-aminophenol (3.27 g, 30.0 mmol) in dichloromethane (50 ml) were added 3,3-dimethyl-butyryl chloride (8.08 g, 60.0 mmol) and pyridine (4.85 ml, 60.0 mmol), while cooling the reaction mixture in an ice bath. After the addition was completed, the cooling bath was removed and stirring was continued overnight at room temperature. The solvent was removed by evaporation and the residue was dissolved in THF (300 ml). 6N NaOH (aq, 35 ml) was added and the mixture was stirred at room temperature overnight. The organic phase was removed by evaporation. Water (200 ml) was added and the solid material was collected by filtration, washed with water, dried *in vacuo* at 40 °C and dissolved in methanol (100 ml). A solution of KOH (3.37 g) in methanol (50 ml) was added. After stirring for 2 days at room temperature water (300 ml) was added and the organic solvent was removed by evaporation. The aqueous phase was acidified with 1 N HCl. The solids were collected by filtration and dried under vacuum at 40 °C yielding the title compound (1.97 g, 31%, pink solid). The mother liquor was extracted with ethyl acetate (3 x 250 ml). The combined organic phases were washed with saturated sodium bicarbonate (2 x 250 ml), dried *in vacuo*, filtered and evaporated yielding a second amount of the title compound (0.67 g, 10%). From the first aqueous extract another portion of product was isolated by extraction with ethyl acetate (4 x 250 ml). The combined organic phases were washed with water (400 ml), saturated sodium bicarbonate (2 x 400 ml), dried, filtered and evaporated yielding a pink thick oil. Cry-

tallisation from ethyl acetate/heptane yielded a third amount of the title compound (2.11 g, 34%).

^1H NMR (300 MHz, DMSO- d_6): δ 1.00 (s, 9H), 2.12 (s, 2H), 6.67 (d, 2H), 7.33 (d, 2H), 9.12 (s, 1H), 9.50 (s, 1H); HPLC-MS: m/z = 208 (M+1); R_t = 2.50 min.

5

1-(4-Hydroxy-phenyl)-4,4-dimethyl-piperidine-2,6-dione

A mixture of 4-aminophenol (3.27 g, 30.0 mmol) and 3,3-dimethylglutaric anhydride (4.26 g, 30.0 mmol) was heated in a round bottom flask at 165 °C for 1 h, followed by heating at 180
10 °C for 7 h. After cooling to room temperature the solid material was dissolved in hot ethanol, activated charcoal was added and the solution was heated at reflux for 1 h. The solid material was removed by hot filtration. The solvent was evaporated and the residue was crystallised (water/ethanol) yielding the title compound (3.51 g, 50%, pink solid).

^1H NMR (300 MHz, DMSO- d_6): δ 1.08 (s, 6H), 2.63 (s, 4H), 6.77 + 6.86 (AB-system, 4H),
15 9.56 (s, 1H).

cis-2-(4-Hydroxy-phenyl)-hexahydro-isoindole-1,3-dione

A mixture of 4-aminophenol (5.46 g, 50.0 mmol) and *cis*-1,2-cyclohexanedicarboxylic anhydride (7.71 g, 50.0 mmol) was heated in a round bottom flask at 170 °C for 2 h. After cooling
20 to room temperature the solid material was dissolved in hot ethanol (200 ml), activated charcoal was added and the solution was heated at reflux for 1 h. The solid material was removed by hot filtration. The solvent was partially evaporated. The solids were collected by filtration, washed quickly with a small amount of ethanol and dried *in vacuo* yielding the title
25 compound (8.52 g, 69%, pink solid).

^1H NMR (300 MHz, DMSO- d_6): δ 1.38 (m, 4H), 1.73 (m, 4H), 3.02 (m, 2H), 6.82 (d, 2H), 7.02 (d, 2H), 9.66 (s, 1H, OH); HPLC-MS: m/z = 246 (M+1); R_t = 2.53 min.

Cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide

To a solution of 4-aminophenol (5.00 g, 45.8 mmol) in dichloromethane (50 ml) were added
5 cyclohexanecarbonyl chloride (6.72 g, 45.8 mmol) and pyridine (3.70 ml, 45.8 mmol), while
cooling the reaction mixture in an ice bath. After the addition was completed, the cooling bath
was removed and stirring was continued overnight at room temperature. Water (100 ml) was
added, the dichloromethane was removed by evaporation and the resulting solution was ex-
tracted with ethyl acetate (3 x 300 ml). The combined organic phases were washed with wa-
10 ter (2 x 200 ml), dried, filtered and evaporated, yielding an off-white solid. The crude product
was purified by flash column chromatography (SiO₂, ethyl acetate/heptane (2:3)), yielding a
mixture of two compounds, which were dissolved in THF. 6N NaOH (aq, 32 ml) was added
and the mixture was stirred at room temperature for 2.5 h. The solution was acidified with
concentrated hydrochloric acid. The THF was removed by evaporation. The solid material
15 was collected by filtration, dried *in vacuo* and recrystallised (ethyl acetate/heptane), yielding
the title compound (4.20 g, 41%, off-white solid).

¹H NMR (300 MHz, DMSO-*d*₆): δ 1.12-1.48 (m, 5H), 1.65 (m, 1H), 1.70-1.82 (m, 4H), 2.27
(m, 1H), 6.66 (d, 2H), 7.36 (d, 2H), 9.10 (s, 1H), 9.50 (s, 1H); HPLC-MS: *m/z* = 220 (M+1); *R*_t
= 2.69 min.

20

2-Cyclohexyl-N-(4-hydroxy-phenyl)-acetamide

To a solution of 4-aminophenol (3.83 g, 35.1 mmol) in dichloromethane (50 ml) were added
cyclohexylacetyl chloride (11.26 g, 70.1 mmol) and pyridine (5.67 ml, 70.1 mmol), while cool-
25 ing the reaction mixture in an ice bath. After the addition was completed, the cooling bath
was removed and stirring was continued overnight at room temperature. The solvent was
removed by evaporation and the residue was dissolved in THF (300 ml). 6N NaOH (aq, 41
ml) was added and the mixture was stirred at room temperature for 4 h. The solution was
acidified with 1 N hydrochloric acid and the organic phase was removed by evaporation. The
30 solid material was collected by filtration, dried and dissolved in methanol (100 ml). A solution
of KOH (5.5 g) in methanol (50 ml) was added. After stirring for 1 h at room temperature wa-
ter (200 ml) was added and the organic solvent was removed by evaporation. The aqueous

phase was acidified with 1 N HCl. The title product was isolated by filtration and dried *in vacuo* (6.31 g, 77%, pink crystals).

¹H NMR (200 MHz, DMSO-*d*₆): δ 0.82-1.32 (m, 5H), 1.54-1.76 (m, 6H), 2.12 (d, 2H), 6.66 (d, 2H), 7.32 (d, 2H), 9.12 (s, 1H), 9.57 (s, 1H); HPLC-MS: *m/z* = 234 (M+1); *R*_t = 3.09 min.

5

***cis/trans*-4-*tert*-Butyl-cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide**

To a solution of 4-aminophenol (3.08 g, 28.2 mmol) in dichloromethane (50 ml) were added *cis/trans*-4-*tert*-butyl-cyclohexanecarbonyl chloride (11.43 g, 56.4 mmol) and pyridine (4.56
10 ml, 56.4 mmol), while cooling the reaction mixture in an ice bath. After the addition was completed, the cooling bath was removed and stirring was continued overnight at room temperature. The solvent was removed by evaporation and the residue was dissolved in THF (300 ml). 6N NaOH (aq, 33 ml) was added and the mixture was stirred at room temperature overnight. The organic phase was removed by evaporation. Water (200 ml) was added and the
15 solid material collected by filtration, washed with water, dried and dissolved in methanol (100 ml). A solution of KOH (2.4 g) in methanol (50 ml) was added. After stirring for 2 h at room temperature water (200 ml) was added and the organic solvent was removed by evaporation. The aqueous phase was acidified with 1 N HCl and extracted with ethyl acetate (3 x 300 ml). The combined organic phases were dried and evaporated, yielding a pink oil, which was
20 dried *in vacuo*. The solid material was crystallised from ethyl acetate/heptane yielding the title compound (2.03 g, 26%) as pink crystals. From the first aqueous extract a second portion of product was isolated by extraction with ethyl acetate (3 x 250 ml). The combined organic layers were washed with water (400 ml), saturated sodium bicarbonate (2 x 400 ml), dried, filtered and evaporated, yielding a pink thick oil. Crystallisation from ethyl acetate/heptane yielded a second amount of the title compound (2.75 g, 35%).
25

¹H NMR (300 MHz, DMSO-*d*₆): δ 0.80 + 0.84 (2 x s, 9H), 0.98 (m, 2H), 1.23-1.57 (m, 4H), 1.76-1.90 (m, 2H), 2.02-2.14 (m, 1.5H), 2.57 (m, 0.5H), 6.65 (d, 2H), 7.34 (d x d, 2H), 9.09 (s, 1H), 9.36 + 9.50 (2 x s, 1H); HPLC-MS: *m/z* = 276 (M+1); *R*_t = 4.19 and 4.27 min.

30 ***N*-(4-Hydroxy-phenyl)-3,3-dimethyl-butylamide**

To a solution of 4-aminophenol (3.27 g, 30.0 mmol) in dichloromethane (50 ml) were added 3,3-dimethyl-butyryl chloride (8.08 g, 60.0 mmol) and pyridine (4.85 ml, 60.0 mmol), while cooling the reaction mixture in an ice bath. After the addition was completed, the cooling bath was removed and stirring was continued overnight at room temperature. The solvent was removed by evaporation and the residue was dissolved in THF (300 ml). 6N NaOH (aq, 35 ml) was added, the mixture was stirred at room temperature overnight and the solvent was removed by evaporation. Water (200 ml) was added and the solid material is collected by filtration, washed with water, dried *in vacuo* at 40 °C and dissolved in methanol (100 ml). A solution of KOH (3.37 g) in methanol (50 ml) was added. After stirring for 2 days at room temperature water (300 ml) was added, the organic solvent was removed and the aqueous phase was acidified with 1 N HCl. The solids were collected and dried yielding the title compound (1.97 g, 31% yield, pink solid). The mother liquor was extracted with ethyl acetate (3 x 250 ml). The combined organic phases were washed with saturated sodium bicarbonate (2 x 250 ml), dried over sodium sulphate, filtered and evaporated yielding a second amount of the title compound (0.67 g, 10%). From the first aqueous extract another portion of product was isolated by extraction with ethyl acetate (4 x 250 ml). The combined organic phases were washed with water (400 ml), dried, filtered and evaporated yielding a pink thick oil. Crystallisation from ethyl acetate/heptane yielded a third amount of the title compound (2.11 g, 34%). ¹H NMR (300 MHz, DMSO-*d*₆): δ 1.00 (s, 9H), 2.12 (s, 2H), 6.67 (d, 2H), 7.33 (d, 2H), 9.12 (s, 1H), 9.50 (s, 1H); HPLC-MS: *m/z* = 208 (M+1); R_t = 2.50 min.

4-(3-Trifluoromethyl-phenoxy)-phenol

Hydroquinone monobenzylether (1 g, 5.0 mmol), 3-(trifluoromethyl)-phenyl boronic acid (1.9 g, 10.0 mmol), copper (II) acetate (0.91 g, 5.0 mmol) and triethylamine (2.53 g, 25.0 mmol) were dissolved/suspended in dichloromethane (50 ml). The reaction mixture was stirred for 70 h. at room temperature and evaporated to dryness. The crude intermediate was subjected to flash chromatography (ethyl acetate/heptane (1:4)) (42%) and hydrogenated (10% Pd/C) using ethanol as a solvent. The organic phase was evaporated and aqueous sodium hydroxide (1N, 30 ml) was added together with dichloromethane. The two phases were separated and the aqueous phase extracted with dichloromethane (30 ml x 2). The aqueous phase was acidified with aqueous hydrochloric acid (2N) and extracted with dichloromethane (30 ml x 5). The organic phase was dried and evaporated to give the crude product (47%). HPLC-MS *m/z* = 254.9 (M+1), R_t: 4.39 min.

δ_{H} (300MHz; CDCl_3): 6.85 (dt, 2H), 6.94 (dt, 2H), 7.10 (dd, 1H), 7.16 (bs, 1H), 7.24-7.30 (m, 1H), 7.39 (t, 1H).

4-Hydroxy-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester

5

4-Hydroxybenzoic acid (30 g, 0.217 mmol) and 4-hydroxysuccinamide (25.32 g, 0.220mmol) were dissolved in 1,4-dioxane (550 ml) at room temperature. After 20 min. the clear solution was cooled to 15°C and dicyclohexylcarbodiimide (44.82 ml, 0.217 mmol) was added. The reaction mixture was stirred for 18 hours and filtered. The organic phase was evaporated to dryness (86 g). Ethanol (250 ml) was added to the crude product and the mixture heated to reflux. The crude product was crystallized from ethanol/water (5:1) (22 g, 43%), and the mother liquor recrystallized from ethanol/water (25 g, 49%). HPLC-MS: $m/z = (M+1)$; R_t : min.

10

N-(6-Methoxy-pyridin-3-yl)-benzamide

15

A solution of 5-amino-2-methoxypyridine (2.48 g, 20.0 mmol) and *N*-ethyldiisopropylamine (2.84 g, 22.0 mmol) in dichloromethane (20 ml) was cooled in an ice-bath. Benzoyl chloride (3.09 g, 22 mmol) was slowly added by means of a syringe. The cooling bath was removed and stirring was continued at room temperature for 18 hours. Dichloromethane was added and the solution was extracted with water. The organic layer was dried over sodium sulphate, filtered and evaporated *in vacuo* leaving a dark solid. Crystallisation from ethyl acetate:heptane yielded the title compound (3.44 g, 75% yield).

20

^1H NMR (300MHz, CDCl_3): δ 3.93 (s, 3H), 6.77 (d, 1H), 7.44-7.59 (m, 3H), 7.81 (br.s, 1H), 7.87 (d, 2H), 8.01 (dd, 1H), 8.16 (d, 1H); HPLC-MS (Method A): $m/z = 229$ (M+H); $R_t = 2.52$ min.

25

Cyclohexanecarboxylic acid (6-methoxy-pyridin-3-yl)-amide hydrochloride

5-Amino-2-methoxypyridine (3.72 g, 30.0 mmol), dissolved in a small amount of tetrahydrofuran, was added slowly to a solution of cyclohexanecarbonyl chloride (4.40 g, 30.0 mmol) in tetrahydrofuran (25 ml). After standing for 0.5 hours diethyl ether (250 ml) was added and the solids were collected by suction yielding the title compound (8.12 g, 100% yield) as a purple solid.

30

¹H NMR (300MHz, CDCl₃): δ 1.11-1.48 (m, 5H), 1.63 (m, 1H), 1.68-1.83 (m, 4H), 2.32 (m, 1H), 3.81 (s, 3H), 6.80 (d, 1H), 7.92 (dd, 1H), 8.00 (br.s, 1H), 8.38 (d, 1H), 9.92 (s, 1H); HPLC-MS (Method A): m/z = 235 (M+H); R_t = 2.89 min.

5 **6'-Methoxy-4,4-dimethyl-4,5-dihydro-3H-[1,3']bipyridinyl-2,6-dione**

A mixture of 5-amino-2-methoxypyridine (3.72 g, 30.0 mmol) and 3,3-dimethylglutaric anhydride (4.26 g, 30.0 mmol) was heated at 175 °C for 7 hours. After cooling down to room temperature the solid material was dissolved in a small amount of dichloromethane and purified
10 by flash column chromatography (SiO₂, ethyl acetate:heptane (40:60)) yielding the title compound (2.56 g, 34% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 1.20 (s, 6H), 2.69 (s, 4H), 3.95 (s, 3H), 3.81 (d, 1H), 7.28 (dd, 1H), 7.89 (d, 1H); HPLC-MS (Method A): m/z = 249 (M+H); R_t = 2.43 min.

15 **N-(6-Methoxy-pyridin-3-yl)-2,2-dimethyl-propionamide hydrochloride**

5-Amino-2-methoxypyridine (3.72 g, 30.0 mmol), dissolved in a small amount of tetrahydrofuran, was added slowly to a solution of 2,2-dimethylpropionyl chloride (3.62 g, 30.0 mmol) in tetrahydrofuran (25 ml). After standing for 0.5 hours diethyl ether (250 ml) was added and a
20 thick oil precipitated. The solvent was decanted and the residue was dried under reduced pressure yielding the title compound (5.50 g, 75% yield) as a purple foam.

¹H NMR (300MHz, CDCl₃): δ 1.22 (s, 9H), 3.83 (s, 3H), 6.86 (d, 1H), 8.00 (dd, 1H), 8.42 (d, 1H), 9.41(s, 1H), 9.54 (br.s, 1H); HPLC-MS (Method A): m/z = 209 (M+H); R_t = 2.28 min.

25 **2-Cyclohexyl-N-(6-methoxy-pyridin-3-yl)-acetamide hydrochloride**

5-Amino-2-methoxypyridine (3.72 g, 30.0 mmol), dissolved in a small amount of tetrahydrofuran, was added slowly to a solution of cyclohexylacetyl chloride (4.82 g, 30.0 mmol) in tetrahydrofuran (25 ml). After standing for 0.5 hours diethyl ether (250 ml) was added and the sol-
30 ids were collected by suction yielding the title compound (8.54 g, 100% yield) as a purple solid.

¹H NMR (300MHz, DMSO-d₆): δ 0.88-1.04 (m, H), 1.09-1.32 (m, 3H), 1.54-1.82 (m, 6H), 2.18 (d, 2H), 3.84 (s, 3H), 6.85 (d, 1H), 7.98 (dd, 1H), 8.41 (d, 1H), 9.81 (br.s, 1H), 10.10 (s, 1H); HPLC-MS (Method A): m/z = 249 (M+H); R_t = 3.32 min.

***N*-(6-Hydroxy-pyridin-3-yl)-benzamide**

N-(6-Methoxy-pyridin-3-yl)-benzamide (2.38 g, 10.4 mmol) was dissolved in a mixture of tetrahydrofuran and diethyl ether. HCl-gas was bubbled into the solution for 5 minutes. More diethyl ether was added and the white precipitate was collected by suction, washed twice with diethyl ether and heated in a kugelrohr apparatus at 180 °C for 0.5 hours. The solid material was crystallised from methanol:water, washed twice with water and dried overnight in a vacuum oven, yielding the title compound (1.19 g, 53% yield) as a grey solid.

¹H NMR (300MHz, CDCl₃): δ 6.39 (d, 1H), 7.47-7.61 (m, 3H), 7.18 (dd, 1H), 7.91 (d, 2H), 7.96 (d, 1H); HPLC-MS (Method A): *m/z* = 215 (M+H); R_t = 1.52 min.

Cyclohexanecarboxylic acid (6-hydroxy-pyridin-3-yl)-amide

Cyclohexanecarboxylic acid (6-methoxy-pyridin-3-yl)-amide hydrochloride (8.12 g, 30.0 mmol) was heated in a kugelrohr apparatus at 190 °C for 25 minutes. After cooling to room temperature the solid material was crystallised from methanol:water, washed twice with water and dried overnight in a vacuum oven, yielding the title compound (3.09 g, 47% yield) as a purple solid.

¹H NMR (300MHz, CDCl₃): δ 1.09-1.46 (m, 5H), 1.63 (m, 1H), 1.68-1.80 (m, 4H), 2.23 (m, 1H), 6.33 (d, 1H), 7.44 (dd, 1H), 7.87 (d, 1H), 9.54 (s, 1H), 11.29 (br.s, 1H); HPLC-MS (Method A): *m/z* = 221 (M+H); R_t = 1.84 min.

6'-Hydroxy-4,4-dimethyl-4,5-dihydro-3*H*-[1,3']bipyridinyl-2,6-dione

6'-Methoxy-4,4-dimethyl-4,5-dihydro-3*H*-[1,3']bipyridinyl-2,6-dione (2.56 g, 10.3 mmol) was dissolved in a mixture of tetrahydrofuran and diethyl ether. HCl-gas was bubbled into the solution for 5 minutes. More diethyl ether was added and the white precipitate was collected by suction, washed twice with diethyl ether and heated in a kugelrohr apparatus at 190 °C for 15 minutes yielding the title compound (2.16 g, 89% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 1.08 (s, 6H), 2.52 (s, 4H), 6.34 (d, 1H), 7.19 (dd, 1H), 7.31 (d, 1H), 11.42 (br.s, 1H); HPLC-MS (Method A): *m/z* = 235 (M+H); R_t = 1.32 min.

***N*-(6-Hydroxy-pyridin-3-yl)-2,2-dimethyl-propionamide**

Under a stream of nitrogen gas *N*-(6-methoxy-pyridin-3-yl)-2,2-dimethyl-propionamide hydrochloride (5.50 g, 22.5 mmol) was heated in a round bottom flask at 180 °C for 15 minutes. After cooling to room temperature the solid material was crystallised from a methanol-water mixture, yielding the title compound (1.13 g, 26% yield) as a dark grey solid.

- 5 ¹H NMR (300MHz, CDCl₃): δ 1.18 (s, 9H), 6.32 (d, 1H), 7.53 (dd, 1H), 7.76 (d, 1H), 8.97 (br.s, 1H), 11.30 (br.s, 1H); HPLC-MS (Method A): *m/z* = 195 (M+H); *R*_t = 1.15 min.

2-Cyclohexyl-*N*-(6-hydroxy-pyridin-3-yl)-acetamide

- 10 2-Cyclohexyl-*N*-(6-methoxy-pyridin-3-yl)-acetamide hydrochloride (8.54 g, 30.0 mmol) was heated in a kugelrohr apparatus at 160 °C for 0.5 hours. After cooling to room temperature the solid material was crystallised from a methanol-water mixture, washed twice with water and dried overnight in a vacuum oven, yielding the title compound (4.53 g, 64% yield) as a grey solid.
- 15 ¹H NMR (300MHz, CDCl₃): δ 0.90-1.38 (m, 5H), 1.60-1.92 (m, 6H), 2.17 (d, 2H), 6.50 (d, 1H), 7.50 (dd, 1H), 7.97 (d, 1H), 8.80 (br.s, 1H), 11.81 (br.s, 1H); HPLC-MS (Method A): *m/z* = 235 (M+H); *R*_t = 2.29 min.

3-Dimethylamino-2-(4-methoxy-phenoxy)-propenal

- 20 Phosphorus oxychloride (18.4 g, 120 mmol) was added to dimethylformamide (8.8 g, 120 mmol), maintaining the temperature below 25 °C by an external ice-bath. Upon completion of the addition the reaction mixture was heated to 50 °C for 45 minutes and then cooled to room temperature. Chloroform (35 ml) was added and the resulting solution was brought to reflux.
- 25 4-Methoxyphenoxyacetaldehyde diethylacetal (9.61 g, 40.0 mmol) was added in portions. After heating for 3 hours at reflux the solution was cooled to room temperature and poured carefully onto a solution of potassium carbonate (115 g) in water (100 ml). The mixture was cooled in an ice-bath to around room temperature and extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo* yielding a brown oil. The residue was heated with ethyl acetate:heptane and decanted, leaving a brown oil. The solvent was removed *in vacuo* to give a brown oil, which was purified by flash column chromatography (SiO₂, ethyl acetate) yielding the title compound (3.87 g, 44% yield) as a white solid.
- 30 ¹H NMR (300MHz, CDCl₃): δ 3.00 (s, 6H), 3.75 (s, 3H), 6.54 (s, 1H), 7.80 + 7.87 (AB-system, 4H), 8.80 (s, 1H); HPLC-MS (Method A): *m/z* = 222 (M+H); *R*_t = 1.73 min.
- 35

2-(3,4-Dichloro-phenoxy)-3-dimethylamino-propenal

Phosphorus oxychloride (18.4 g, 120 mmol) was added to dimethylformamide (8.8 g, 120
5 mmol), maintaining the temperature below 25 °C by an external ice-bath. Upon completion of
the addition the reaction mixture was heated to 50 °C for 45 minutes and then cooled to room
temperature. Chloroform (35 ml) was added and the resulting solution was brought to reflux.
3,4-Dichlorophenoxyacetaldehyde diethylacetal (9.61 g, 40.0 mmol) was added in portions.
After heating for 3 hours at reflux the solution was cooled to room temperature and poured
10 carefully onto a solution of potassium carbonate (115 g) in water (100 ml). The mixture was
cooled in an ice-bath to around room temperature and extracted twice with dichloromethane.
The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*
yielding a brown oil, which was purified by flash column chromatography (SiO₂, ethyl
acetate) yielding the title compound (5.38 g, 52% yield) as a brown solid.
15 ¹H NMR (300MHz, CDCl₃): δ 3.10 (s, 6H), 6.58 (s, 1H), 6.82 (dd, 1H), 7.03 (d, 1H), 7.30 (d,
1H), 8.80 (s, 1H); HPLC-MS (Method A): *m/z* = 260 (M+H); R_t = 3.14 min.

5-(4-Methoxy-phenoxy)-pyrimidin-2-ol

20 A solution of sodium ethoxide, prepared from sodium (0.80 g, 35.0 mmol), 3-dimethylamino-
2-(4-methoxy-phenoxy)-propenal (3.87 g, 17.5 mmol) and urea (2.10 g, 35.0 mmol) in etha-
nol (25 ml) was heated at reflux for 4 hours. Water (1 ml) was added and heating was con-
tinued for an additional 2 hours. The solution was cooled to room temperature and neutral-
ised with glacial acetic acid. Most of the solvent was removed by evaporation *in vacuo*. Wa-
25 ter was added and the precipitate was isolated by suction, followed by drying in a vacuum
oven, yielding the title compound (0.80 g, 21% yield) as a yellow solid.
¹H NMR (300MHz, CDCl₃): δ 3.80 (s, 3H), 6.85-7.95 (AB-system, 4H), 8.12 (s, 2H); HPLC-
MS (Method A): *m/z* = 219 (M+H); R_t = 1.77 min.

5-(3,4-Dichloro-phenoxy)-pyrimidin-2-ol

A solution of sodium ethoxide, prepared from sodium (0.95 g, 41.4 mmol), 2-(3,4-dichloro-
phenoxy)-3-dimethylamino-propenal (5.38 g, 20.7 mmol) and urea (2.48 g, 41.4 mmol) in
ethanol (25 ml) was heated at 60 °C for 4 hours. Water (1 ml) was added and heating was
35 continued for an additional 2 hours. The solution was cooled to room temperature and neu-

tralised with glacial acetic acid. Most of the solvent was removed by evaporation *in vacuo*. Water was added and the precipitate was isolated by suction, followed by drying in a vacuum oven, yielding the title compound (0.92 g, 17% yield) as a brown solid.

¹H NMR (300MHz, DMSO-*d*₆): δ 7.08 (dd, 1H), 7.38 (d, 1H), 7.59 (d, 1H), 8.35 (s, 2H), 12.06 (br.s, 1H); HPLC-MS (Method A): *m/z* = 257 (M+H); *R*_t = 2.75 min.

5-(2-Nitro-phenyl)-pyrimidin-2-ol

10 A solution of 3-(dimethylamino)-2-(2-nitrophenyl)acrylaldehyde (2.00 g, 9.08 mmol), urea (0.60 g, 9.99 mmol) and concentrated hydrochloric acid (0.50 ml) in ethanol (25 ml) was heated at 60 °C for 18 hours under a nitrogen atmosphere. An additional aliquot of concentrated hydrochloric acid (0.50 ml) was added followed by heating at 70 °C for 24 hours. The solvent was removed by evaporation under reduced pressure. The residue was crystallised
15 from methanol yielding the title compound (0.35 g, 18% yield) as a yellow solid.

¹H NMR (300MHz, DMSO-*d*₆): δ 7.54 (dd, 1H), 7.62 (dt, 1H), 7.76 (dt, 1H), 8.06 (dd, 1H), 8.29 (s, 2H); HPLC-MS (Method A): *m/z* = 218 (M+H); *R*_t = 1.26 min.

20 N-(6-Hydroxy-pyridin-3-yl)-3,3-dimethyl-butyramide

3,3-Dimethylbutyroyl chloride (4.04 g, 30.0 mmol) was added dropwise to a stirred solution of 5-amino-2-methoxypyridine (3.72 g, 30.0 mmol) in tetrahydrofuran (25 mL). After stirring for 1 hour at room temperature, diethyl ether was added and the solid material was isolated by
25 suction. The *N*-(6-methoxy-pyridin-3-yl)-3,3-dimethyl-butyramide hydrochloride (4.13 g, 15.96 mmol) was heated at 180 °C for 15 minutes. After cooling to room temperature the product was dissolved in methanol. Partial evaporation of the solvent yielded the title compound (1.15 g, 35% yield) as a solid.

¹H NMR (300MHz, DMSO-*d*₆): δ = 1.00 (s, 9H), 2.12 (s, 2H), 6.38 (d, 1H), 7.44 (dd, 1H), 7.89 (d, 1H), 9.59 (s, 1H), 11.42 (br.s, 1H); HPLC-MS (Method A): *m/z* = 209 (M+H)⁺; *R*_t = 1.71
30 min.

Pyridine-2-carboxylic acid (6-methoxy-pyridin-3-yl)-amide dihydrochloride

5-Amino-2-methoxypyridine (4.40 g, 35.4 mmol), dissolved in a small amount of tetrahydrofuran, was added slowly to a stirred solution of pyridine-2-carbonyl chloride hydrochloride (7.12 g, 40.0 mmol) in tetrahydrofuran (75 mL). After stirring overnight at room temperature diethyl ether was added. The solids were isolated by suction, washed with diethyl ether and dried in a vacuum oven at 45 °C, yielding the title compound (7.89 g, 75%).

¹H NMR (300MHz, CDCl₃): δ = 3.87 (s, 3H), 6.90 (d, 1H), 7.03 (br.s, 2 HCl + H₂O), 7.70 (m, 1H), 8.09 (dt, 1H), 8.17 (d, 1H), 8.22 (dd, 1H), 8.68 (d, 1H), 8.73 (m, 1H), 10.82 (s, 1H);
HPLC-MS (Method A): m/z = 230 (M+H)⁺; Rt = 2.45 min and 264 + 266; Rt = 3.15 min.

Pyridine-2-carboxylic acid (6-hydroxy-pyridin-3-yl)-amide hydrochloride

Pyridine-2-carboxylic acid (6-methoxy-pyridin-3-yl)-amide dihydrochloride (0.66 g, 1.99 mmol) was heated at 180 °C for 10 minutes. After cooling to room temperature, the title compound was obtained and used without further purification.
HPLC-MS (Method A): m/z = 216 (M+H)⁺; Rt = 2.14 min.

5-Methoxy-pyrimidin-2-ylamine

Under a nitrogen atmosphere phosphorus pentachloride (21 g, 0.10 mol) was added portionwise to 1,1,3-trimethoxyethane (12.0 g, 0.10 mol) with stirring and external cooling (ice-bath). After addition was completed, stirring was continued for an additional 30 minutes at room temperature. Dimethylformamide (22.5 mL) was added by means of a dropping funnel, while the reaction mixture was cooled externally with an ice-bath. After completion of the addition, the reaction mixture was heated at 60 °C for 70 minutes. The reaction mixture was then cooled in an ice-bath and methanol (50 mL) was added dropwise. The resulting solution was added dropwise to a stirred solution of sodium hydroxide (24 g) in methanol (80 mL) while cooling in an ice-bath. Guanidine nitrate (20.0 g, 0.16 mol) and sodium hydroxide (7.0 g, 0.175 mol) were added and the solution was stirred for 18 hours at room temperature. Water (150 mL) was added and the solution was extracted three times with dichloromethane. The combined organic layers were evaporated *in vacuo* leaving a brown oil. According to ¹H NMR analysis a mixture of the desired product and the intermediate β -dimethylamine- α -methoxyacroleine was obtained. The mixture was dissolved in methanol (100 mL). Guanidine

nitrate (15.0 g, 0.12 mol) and sodium hydroxide (5.25 g, 0.13 mol) were added and the reaction mixture was heated at 60 °C for 3 hours, followed by stirring at room temperature for 3 days. Water was added and the solution was extracted three times with dichloromethane.

The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo* yielding the title compound (5.43 g, 43% yield) as a yellow solid.

¹H NMR (300MHz, CDCl₃): δ = 3.80 (s, 3H), 5.08 (br.s, 2H), 8.04 (s, 2H); HPLC-MS (Method A): m/z = 126 (M+H)⁺; Rt = 0.39 min.

1-(5-Methoxy-pyrimidin-2-yl)-4,4-dimethyl-piperidine-2,6-dione

- 10 A mixture of 5-methoxy-pyrimidin-2-ylamine (1.00 g, 7.99 mmol) and 3,3-dimethylglutaric anhydride (1.14 g, 7.99 mmol) was heated at 180 °C for 9 hours. After cooling to room temperature the reaction mixture was dissolved in a small amount of dichloromethane and purified by flash column chromatography (SiO₂, ethyl acetate:heptane 70:30), yielding the title compound (0.79 g, 40% yield) as a white solid.

- 15 ¹H NMR (300MHz, CDCl₃): δ = 1.23 (s, 6H), 2.67 (s, 4H), 3.97 (s, 3H), 8.48 (s, 2H); HPLC-MS (Method A): m/z = 250 (M+H)⁺; Rt = 1.86 min.

1-(5-Hydroxy-pyrimidin-2-yl)-4,4-dimethyl-piperidine-2,6-dione

- 20 A mixture of 1-(5-methoxy-pyrimidin-2-yl)-4,4-dimethyl-piperidine-2,6-dione (0.99 g, 3.97 mmol) and pyridine hydrochloride (1.50 g, 7.99 mmol) was heated at 190 °C for 2.5 hours. After cooling to room temperature the reaction mixture was dissolved in a small amount of dichloromethane and filtered over a short pad of silicagel and washed with ethyl acetate. Evaporation of the solvent *in vacuo* yielded the title compound (0.60 g, 64 % yield) as a white solid.

- 25 ¹H NMR (300MHz, CDCl₃): δ = 1.23 (s, 6H), 2.65 (s, 4H), 8.42 (s, 2H), 9.94 (br.s, 1H); HPLC-MS (Method A): m/z = 236 (M+H)⁺; Rt = 1.53 min.

6-Chloro-N-(6-hydroxy-pyridin-3-yl)-nicotinamide

- 30 A solution of 6-chloro-nicotinoyl chloride (0.40 g, 2.27 mmol) and 5-amino-2-hydroxypyridine hydrochloride (0.33 g, 2.25 mmol) in dry tetrahydrofuran (15mL) was stirred at room tempera-

ture for 1 hour. Saturated sodium bicarbonate (aq) was added and the solution was extracted three times with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was dissolved in a mixture of methanol (10 mL) and aqueous sodium hydroxide (1N, 2 mL). After stirring for 2 hours at room temperature water was added and the solution was extracted with dichloromethane. The organic layer was dried over sodium sulphate, filtered and evaporated *in vacuo*, yielding the title compound which was used without further purification.

HPLC-MS (Method A): m/z = 250 (M+H)⁺; Rt = 1.52 min.

N-(2,2-Dimethyl-propyl)-6-hydroxy-nicotinamide

A solution of 6-hydroxynicotinic acid (1.39 g, 10.0 mmol), 1-hydroxy-7-azabenzotriazole (1.50 g, 11.0 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.11 g, 11.0 mmol) in dimethylformamide (25 mL) was stirred at room temperature for 20 minutes. A solution of 2,2-dimethylpropylamine (0.96 g, 11.0 mmol) and *N,N*-diisopropylethylamine (1.42 g, 11.0 mmol) in a small amount of dimethylformamide was added. Stirring was continued for 0.5 hour at room temperature. Ethyl acetate was added and the reaction mixture was extracted twice with water. The solvent was evaporated *in vacuo* yielding the title compound, which was used without further purification.

HPLC-MS (Method A): m/z = 209 (M+H)⁺; Rt = 1.86 min.

3-Chloro-6-(3,4-dichloro-phenoxy)-pyridazine

A solution of 3,6-dichloropyridazine (4.47 g, 30.0 mmol), 3,4-dichlorophenol (4.89 g, 30.0 mmol) and potassium hydroxide (1.68 g, 30.0 mmol) in dimethyl sulfoxide (20 mL) was heated at 60 °C overnight. The solvent was removed by evaporation *in vacuo*. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50). Small amounts of starting material were removed by kugelrohr distillation. Crystallisation from ethyl acetate:heptane yielded the title compound (1.74 g, 21% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 7.10 (dd, 1H), 7.20 (d, 1H), 7.37 (d, 1H), 7.49 (d, 1H), 7.54 (d, 1H); HPLC-MS (Method A): m/z = 275 and 277 (M+H)⁺; Rt = 4.00 min.

6-(3,4-dichloro-phenoxy)-pyridazin-3-ol

A solution of 3-chloro-6-(3,4-dichloro-phenoxy)-pyridazine (1.74 g, 6.32 mmol) in formic acid (25 mL) was heated at 100 °C for 3 hours. The solvent was removed by evaporation *in vacuo* yielding the title compound, which was used without further purification.
HPLC-MS (Method A): m/z = 257 (M+H)⁺; Rt = 3.13 min.

(4-Hydroxy-piperidin-1-yl)-imidazol-1-yl-methanone

A solution of 4-hydroxypiperidine (20.0 g, 198 mmol) and *N,N'*-carbonyldiimidazole (32.06 g, 198 mmol) in tetrahydrofuran (250 mL) was heated overnight at reflux, followed by stirring at room temperature for two days. The solvent was evaporated yielding the title compound, which was used without further purification.
HPLC-MS (Method A): m/z = 196 (M+H)⁺; Rt = 0.39 min.

[4-(tert-Butyl-dimethyl-silanyloxy)-piperidin-1-yl]-imidazol-1-yl-methanone

tert-Butyldimethylsilyl chloride (30.14 g, 0.20 mol) was added to a stirred solution of (4-hydroxy-piperidin-1-yl)-imidazol-1-yl-methanone (39.05 g, 0.20 mol) in dimethylformamide (200 mL). After stirring for 3 days at room temperature, the solvent was removed by evaporation *in vacuo*. The residue was redissolved in dichloromethane, extracted twice with water, dried over sodium sulphate, filtered and evaporated *in vacuo*, yielding the title compound, which was used without further purification.
¹H NMR (300MHz, CDCl₃): δ = 0.07 (s, 6H), 0.90 (s, 9H), 1.64 (m, 2H), 1.81 (m, 2H), 3.53-3.74 (m, 4H), 4.06 (m, 1H), 7.07 (m, 1H), 7.18 (m, 1H), 7.86 (m, 1H); HPLC-MS (Method A): m/z = 310 (M+H)⁺; Rt = 3.40 min.

3-[4-(tert-Butyl-dimethyl-silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium iodide

Methyl iodide (113.5 g, 0.80 mol) was added to a stirred solution of [4-(tert-butyl-dimethyl-silanyloxy)-piperidin-1-yl]-imidazol-1-yl-methanone (61.9 g, 0.20 mol) in acetonitrile (400 mL). After stirring overnight at room temperature the solvent was evaporated *in vacuo*. The resi-

due was washed with ethyl acetate:heptane and dried in a vacuum oven at 45 °C, yielding the title compound (60.92 g, 68% yield over three steps) as a white solid

¹H NMR (300MHz, CDCl₃): δ = 0.08 (s, 6H), 0.88 (s, 9H), 1.57 (m, 2H), 1.83 (m, 2H), 3.45 (m, 2H), 3.65 (m, 2H), 3.93 (s, 3H), 4.06 (m, 1H), 7.87 (m, 1H), 8.03 (m, 1H), 9.56 (s, 1H);

5 HPLC-MS (Method A): m/z = 324 (M-I)⁺; Rt = 2.95 min.

4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide

4-Chlorobenzoyl chloride (1.75 g, 10.0 mmol) was added carefully to a stirred solution of 5-amino-2-methoxypyridine (1.24 g, 10.0 mmol) in dichloromethane (10 mL). After stirring
10 overnight at room temperature, the solvent is evaporated *in vacuo* and the residue is heated in a pre-heated kugelrohr oven at 200 °C for 5-10 minutes under house-vacuum (around 20 mbar) yielding the title compound, which was used without further purification.

HPLC-MS (Method A): m/z = 249 (M+H)⁺; Rt = 2.18 min.

4-Fluoro-N-(6-hydroxy-pyridin-3-yl)-benzamide

15

Starting from 4-fluorobenzoyl chloride (1.59 g, 10.0 mmol) and using the procedure as described for the preparation of 4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide yielded the title compound, which was used without further purification.

HPLC-MS (Method A): m/z = 233 (M+H)⁺; Rt = 1.76 min.

20 N-(6-Hydroxy-pyridin-3-yl)-3-methoxy-benzamide

Starting from 3-methoxybenzoyl chloride (1.71 g, 10.0 mmol) and using the procedure as described for the preparation of 4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide yielded the title compound, which was used without further purification.

25 HPLC-MS (Method A): m/z = 245 (M+H)⁺; Rt = 1.81 min.

N-(6-Hydroxy-pyridin-3-yl)-4-methoxy-benzamide

Starting from 4-methoxybenzoyl chloride (1.71 g, 10.0 mmol) and using the procedure as described for the preparation of 4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide yielded the title compound, which was used without further purification.

5

HPLC-MS (Method A): m/z = 245 (M+H)⁺; Rt = 1.72 min.

N-(6-Hydroxy-pyridin-3-yl)-4-methoxy-benzamide

Starting from 2,4-dichlorobenzoyl chloride (2.10 g, 10.0 mmol) and using the procedure as described for the preparation of 4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide yielded the title compound, which was used without further purification.

10

HPLC-MS (Method A): m/z = 283 (M+H)⁺; Rt = 2.28 min.

N-(6-Hydroxy-pyridin-3-yl)-4-trifluoromethyl-benzamide

Starting from 4-trifluoromethylbenzoyl chloride (2.09 g, 10.0 mmol) and using the procedure as described for the preparation of 4-Chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide yielded the title compound, which was used without further purification.

15

HPLC-MS (Method A): m/z = 283 (M+H)⁺; Rt = 2.28 min.

6'-Hydroxy-4,5-dihydro-3H-[1,3']bipyridinyl-2,6-dione

Glutaric anhydride (1.14 g, 10.0 mmol) was added to a solution of 5-amino-2-methoxypyridine (1.24 g, 10.0 mmol) in dichloromethane (25 mL). After standing for 1 hour at room temperature, thionyl chloride (5.95 g, 50.0 mmol) was added, followed by heating to reflux for 0.5 hour. The solvent and excess thionyl chloride were evaporated *in vacuo*, yielding 4-(6-methoxy-pyridin-3-ylcarbamoyl)-butyryl chloride, which was used without further purification.

20

25

¹H NMR (300MHz, CDCl₃): δ = 1.81 (quintet, 2H), 2.28 (t, 2H), 2.38 (t, 2H), 3.87 (s, 3H), 6.91 (d, 1H), 8.00 (dd, 1H), 8.43 (d, 1H), 10.24 (s, 1H, NH).; HPLC-MS (Method A): m/z = 253 (M+H)⁺; Rt = 1.94 min. (analysed as methyl ester).

The crude 4-(6-methoxy-pyridin-3-ylcarbonyl)-butyryl chloride was redissolved in dichloromethane (25 mL). Thionyl chloride (5.95 g, 50 mmol) was added and the solution was heated to reflux overnight. The solvent and excess thionyl chloride were evaporated *in vacuo*, yielding 6'-methoxy-4,5-dihydro-3*H*-[1,3']bipyridinyl-2,6-dione hydrochloride, which was used

5 without further purification.

¹H NMR (300MHz, CDCl₃): δ = 2.00 (quintet, 2H), 2.72 (t, 4H), 3.88 (s, 3H), 6.90 (d, 1H), 7.51 (dd, 1H), 6.92 (d, 1H), 9.71 (br.s, HCl + H₂O); HPLC-MS (Method A): *m/z* = 221 (M+H)⁺; Rt = 1.38 min.

The crude 6'-methoxy-4,5-dihydro-3*H*-[1,3']bipyridinyl-2,6-dione hydrochloride (2.57 g, 10.0 mmol) was heated in a pre-heated kugerohr oven at 180 °C for 5 minutes. After cooling to room temperature the product was purified by flash column chromatography (SiO₂, ethyl acetate:acetone 25:75), yielding the title compound (0.48 g, 23% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 2.10 (quintet, 2H), 2.81 (t, 4H), 3.53 (br.s, 3H), 7.12 (br.s, 1H), 7.27 (m, 1H), 7.38 (m, 4H), 7.50 (d, 1H), 8.09 (s, 1H); HPLC-MS (Method A): *m/z* = 340 (M+H)⁺; Rt = 2.89 min.

1-(6-Methoxy-pyridin-3-yl)-pyrrolidine-2,5-dione

A mixture of 5-amino-2-methoxypyridine (1.24 g, 10.0 mmol) and succinic anhydride (1.00 g, 10.0 mmol) was heated with a heat gun for 10 minutes. After cooling to room temperature the product was purified by flash column chromatography (SiO₂, ethyl acetate:acetone 25:75). Evaporation of the solvent yielded the title compound as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 2.92 (s, 4H), 3.96 (s, 3H), 6.82 (d, 1H), 7.50 (dd, 1H), 8.11 (d, 1H); HPLC-MS (Method A): *m/z* = 207 (M+H)⁺; Rt = 1.26 min.

1-(6-Hydroxy-pyridin-3-yl)-pyrrolidine-2,5-dione

25

1-(6-Methoxy-pyridin-3-yl)-pyrrolidine-2,5-dione was dissolved in tetrahydrofuran and HCl-gas was bubbled into the solution for 5 minutes. The solvent was evaporated *in vacuo* and the residue was heated for 10 minutes at 180 °C in a pre-heated kugelrohr oven. After cooling to room temperature the residue was purified by flash column chromatography (SiO₂), yielding the title compound (285 mg, 15% yield over two steps).

30

¹H NMR (400MHz, DMSO-*d*₆): δ = 2.72 (s, 4H), 6.40 (d, 1H), 7.31 (dd, 1H), 7.39 (d, 1H), 11.76 (br.s, 1H, OH); HPLC-MS (Method A): *m/z* = 193 (M+H)⁺; Rt = 0.37 min.

2-Methoxy-5-(5-trifluoromethyl-pyridin-2-yloxy)-pyridine

A solution of 5-hydroxy-2-methoxypyridine (1.25 g, 10.0 mmol), 2-chloro-5-trifluoromethylpyridine (1.82 g, 10.0 mmol) and potassium hydroxide (85% pure, 1.08 g, 10.0 mmol) in dimethyl sulfoxide (25 mL) was heated at 90 °C for 2.5 hours. The solution was cooled to room temperature and poured slowly into water (200 mL). After cooling with an external ice-bath, the precipitate was collected by suction, washed thoroughly with water and dried in a vacuum oven at 45 °C, yielding the title compound (2.56 g, 95% yield).

¹H NMR (300MHz, CDCl₃): δ = 3.95 (s, 3H), 6.80 (d, 1H), 7.04 (d, 1H), 7.41 (dd, 1H), 7.91 (dd, 1H), 8.03 (d, 1H), 8.40 (d, 1H); HPLC-MS (Method A): m/z = 271 (M+H)⁺; Rt = 3.88 min.

5-(5-Trifluoromethyl-pyridin-2-yloxy)-pyridin-2-ol

A mixture of 2-methoxy-5-(5-trifluoromethyl-pyridin-2-yloxy)-pyridine (0.28 g, 1.04 mmol) and pyridine hydrochloride (1.00 g, 8.65 mmol) was heated in a kugelrohr oven at 200 °C for 10 minutes. After cooling to room temperature, the reaction mixture is dissolved in dichloromethane and extracted with water, dried over sodium sulphate, filtered and evaporated *in vacuo*, yielding the title compound (180 mg, 68% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 6.63 (m, 1H), 7.06 (d, 1H), 7.42 (m, 2H), 7.92 (dd, 1H), 8.42 (d, 1H); HPLC-MS (Method A): m/z = 257 (M+H)⁺; Rt = 2.32 min.

5-(3,5-Dichloro-pyridin-2-yloxy)-2-methoxy-pyridine

A solution of 5-hydroxy-2-methoxypyridine (1.25 g, 10.0 mmol), 2,3,5-trichloropyridine (1.82 g, 10.0 mmol) and potassium hydroxide (85% pure, 1.08 g, 10.0 mmol) in dimethyl sulfoxide (25 mL) was heated at 90 °C for 1.5 hours. The solution was poured slowly into water (200 mL). The precipitate was collected by suction, washed thoroughly with water and dried in a vacuum oven at 45 °C, yielding the title compound (2.39 g, 88% yield).

¹H NMR (300MHz, CDCl₃): δ = 3.95 (s, 3H), 6.80 (d, 1H), 7.41 (dd, 1H), 7.77 (d, 1H), 7.93 (d, 1H), 8.03 (d, 1H); HPLC-MS (Method A): m/z = 271 (M+H)⁺; Rt = 4.18 min.

5-(3,5-Dichloro-pyridin-2-yloxy)-pyridin-2-ol

A mixture of 5-(3,5-dichloro-pyridin-2-yloxy)-2-methoxy-pyridine (2.39 g, 8.82 mmol) and pyridine hydrochloride (7.00 g, 60.6 mmol) was heated in a kugelrohr apparatus at 200 °C for 25 minutes. After cooling to room temperature dichloromethane and water were added. The solid material, which was insoluble in dichloromethane and water, was isolated by suction and dried in a vacuum oven at 45 °C, yielding the title compound, which was used without further purification.

HPLC-MS (Method A): m/z = 257 (M+H)⁺; Rt = 2.53 min.

4,4-Dimethyl-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-ol

Under a nitrogen atmosphere lithium aluminium hydride (1.90 g, 50.0 mmol) was added portionwise to a stirred suspension of 6'-methoxy-4,4-dimethyl-4,5-dihydro-3H-[1,3']bipyridinyl-2,6-dione (2.85 g, 10.0 mmol) in dry diethyl ether (100 mL). After stirring for 3 hours at room temperature, water (1.90 mL), 15% aqueous sodium hydroxide (1.90 mL) and water (5.70 mL) were added respectively. After stirring for 1 hour at room temperature the salts were removed by filtration and washed three times with diethyl ether. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, dichloromethane followed by ethyl acetate/heptane 25/75) yielding the title compound (1.28 g, 58% yield) as a yellow oil.

¹H NMR (300MHz, CDCl₃): δ = 0.98 (s, 6H), 1.53 (m, 4H), 3.03 (m, 4H), 3.89 (s, 3H), 6.67 (d, 1H), 7.30 (dd, 1H), 7.80 (d, 1H); HPLC-MS (Method A): m/z = 221 (M+H)⁺; Rt = 2.14 min.

4,4'-Dimethyl-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-ol

A mixture of 6'-methoxy-4,4-dimethyl-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl (1.28 g, 5.81 mmol) and pyridine hydrochloride (5.00 g, 43.3 mmol) was heated in a kugelrohr oven at 200 °C for 15 minutes. After cooling to room temperature water was added and the solution was made slightly basic with aqueous 1N sodium hydroxide. The solution was extracted three times with dichloromethane and the combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*, yielding the title compound (0.86 g, 72% yield).

HPLC-MS (Method A): m/z = 207 (M+H)⁺; Rt = 1.31 min.

Methyl-phenyl-carbamic acid 4-amino-phenyl ester

To a solution of N-Boc protected 4-aminophenol (10 mmol) in CH_2Cl_2 (50 mL) was added N-methyl-N-phenylcarbonyl chloride (15 mmol) and DABCO (15 mmol) at room temperature. The reaction mixture was stirred for 16 hours at rt, added CH_2Cl_2 (20 mL) and washed with

5 aqueous citric acid (5%) and brine. The organic phase was separated, dried (MgSO_4) and evaporated to give the crude product which was purified by FC (Quad flash 40 MeOH- CH_2Cl_2 5:95). The purified intermediate was dissolved in CH_2Cl_2 (90 mL). Addition of TFA (6 mL) and stirring for 4 h. The reaction mixture was evaporated to dryness and dried in vacuo at 50 °C overnight producing the title compound in 72% yield as colorless crystals.

10 HPLC-MS : m/z = 243.1 (M+1); R_t = 2.02 min.

Example 1 (General procedure 9)

15 4-[(1,3-Benzodioxol-5-yl)methyl]-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-piperonylpiperazine, yield 67 %. Recrystallisation from 96 % ethanol gave white crystals, m.p. 239 - 240 °C; HPLC-MS m/z = 502 (M+H), 524 (M+Na),

20 R_t = 3.3 min.; ^1H NMR ($\text{DMSO}-d_6$): δ 11.64 (br, 1H, NH), 8.62 - 8.52 (br, 1H, py-H6), 8.31 - 8.19 (m, 1H, py-H4), 7.40 - 7.15 (m, 6H, py-H3 + C_6H_4 + 1 arom.), 7.15 - 6.93 (m, 2H, arom), 4.53 - 3.96 (br, 4H, CH_2 at 4.26 + 2 CH), 3.80 - 2.89 (br, 6H, water at 3.38 + 4C-H); IR (KBr): ν 1724 (C=O).

Example 2 (General procedure 1)

25 Methyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC

30 (36%, white crystals). HPLC-MS m/z = 389.1 (M+1), R_t : 5.13 min.

δ_H (300MHz; CDCl_3): 3.43 (s, 3H), 6.98 (d, 1H, J 8.7), 7.09-7.22 (m, 4H), 7.30-7.34 (m, 1H), 7.35 (d, 2H, J 7.1), 7.40 (t, 2H, J 6.8), 7.88 (dd, 1H, J 8.7 and 2.3), 8.42 (s, 1H).

Example 3 (General procedure 1)

35 Methyl-phenyl-carbamic acid 3-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 3-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad flash 25, dichloromethane) (89%, oil). HPLC-MS m/z = 389.1 (M+1), Rt: 5.08 min. δ_H (300MHz; DMSO- d_6): 3.34 (s, 3H), 7.00-7.15 (m, 3H), 7.27 (t, 2H), 7.35-7.55 (m, 5H), 8.24 (dd, 1H), 8.58 (s, 1H).

Example 4 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(3,5-dichloro-pyridin-2-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (53%, crystals). HPLC-MS m/z = 389.1 (M+1), Rt: 5.1 min. δ_H (300MHz; $CDCl_3$): 3.43 (s, 3H), 7.07-7.20 (m, 4H), 7.27-7.48 (m, 5H), 7.75 (d, 1H, J 2.2), 7.93 (d, 1H, J 2.2).

Example 5 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-ylamino)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-ylamino)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC to give the title product (32%, white crystals). HPLC-MS m/z = 388.2 (M+1), Rt: 4.72 min. δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 6.76 (d, 1H), 6.81 (bs, 1H), 7.12 (d, 2H), 7.27-7.45 (m, 6H), 7.63 (dd, 1H), 8.42 (bs, 1H).

Example 6 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(3,5-dichloro-pyridin-4-yloxy)-phenyl ester

The title compound was prepared from 4-(3,5-dichloro-pyridine-4-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (41%, white crystals). HPLC-MS m/z = 389.1 (M+1), Rt: 4.97 min. δ_H (300MHz; $CDCl_3$): 3.42 (s, 3H), 6.81 (d, 2H, J 9.0), 7.07 (d, 2H, J 7.9), 7.25-7.43 (M, 5H), 8.55 (s, 2H).

Example 7 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester

The title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (78%, white crystals). HPLC-MS m/z = 388.0 (M+1), Rt: 5.59 min.

δ_H (300MHz; $CDCl_3$): 3.43 (s, 3H), 7.02 (d, 4H, J 8.7), 7.14 (d, 2H, J 8.7), 7.31 (1H, d, J 6.8), 7.35 (d, 2H, J 7.2), 7.41 (t, 2H, J 7.1), 7.55 (d, 2H, J 8.6).

Example 8 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(3-trifluoromethyl-phenoxy)-phenyl ester

The title compound was prepared from 4-(3-trifluoromethyl-phenoxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (73%, white crystals). HPLC-MS m/z = 388.2 (M+1), Rt: 5.37 min.

Example 9 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(2-cyano-5-trifluoromethyl-pyridine-3-yloxy)-phenyl ester

The title compound was prepared from 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (74%, colourless oil). HPLC-MS m/z = 414.1 (M+1), Rt: 4.8 min.

δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.11 (d, 2H, J 9), 7.32-7.1 (m, 3H), 7.32-7.50 (m, 5H), 8.63 (s, 1H).

Example 10 (General procedure 1)

Methyl-phenyl-carbamic acid 2-benzenesulfonyl-4-(3-chloro-5-trifluoromethyl-pyridine-2-yloxy)-phenyl ester

The title compound was prepared from 2-benzenesulfonyl-4-(3-chloro-5-trifluoromethyl-pyridine-2-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was purified by preparative HPLC (68%, white crystals). HPLC-MS m/z = 563.1 (M+1), Rt: 5.3 min.

δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.20-7.50 (m, 12H), 7.91 (bs, 1H), 8.00 (d, 1H, J 2.3), 8.23 (s, 1H).

Example 11 (General procedure 1)**Methyl-phenyl-carbamic acid 4-tert-butoxy-phenyl ester**

The title compound was prepared from 4-tert-butoxy-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad12/25, flash 12, dichloromethane) followed by a recrystallization from ethanol (41%, crystals). HPLC-MS m/z = 300.3 ($M+1$), R_t : 4.7 min.

δ_H (300MHz; $CDCl_3$): 1.31 (s, 9H), 3.41 (s, 3H), 6.90-7.07 (m, 4H), 7.20-7.28 (m, 1H), 7.32-7.43 (m, 4H).

Example 12 (General procedure 1)**Methyl-phenyl-carbamic acid 3-(4-fluorobenzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester**

The title compound was prepared from 3-(4-fluorobenzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and N-methyl-N-phenylcarbonyl chloride.

The crude product was of high purity and tested without purification (\approx 100%, crystals); mp: 185-186°C. HPLC-MS m/z = 418.2 ($M+1$), R_t : 5.2 min.

δ_H (300MHz; $CDCl_3$): 2.43 (s, 3H), 3.43 (s, 3H), 6.95 (dt, 2H, J 8.7 and 2.3), 7.10 (m, 2H), 7.21 (dt, 2H, J 7.2 and 2.3), 7.27-7.45 (m, 5H), 7.58 (d, 1H, J 8.3).

Example 13 (General procedure 1)**Methyl-phenyl-carbamic acid 4-phenoxy-phenyl ester**

The title compound was prepared from 4-phenoxy-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was recrystallized (ethanol/water) (86%, white crystals). HPLC-MS m/z = 320.1 ($M+1$), R_t : 5.13 min.

δ_H (300MHz; $CDCl_3$): 3.43 (s, 3H), 6.94-7.02 (m, 4H), 7.08 (t, 2H, J 6.8), 7.04-7.12 (m, 1H), 7.32 (t, 2H, J 7.5), 7.28-7.35 (m, 1H), 7.35-7.45 (m, 4H).

Example 14 (General procedure 1)**Methyl-phenyl-carbamic acid 4-(4-chlorobenzoyl)-phenyl ester**

The title compound was prepared from 4-(4-chlorobenzoyl)-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was recrystallized (ethanol/water) (90%, white crystals). HPLC-MS m/z = 366.1 ($M+1$), R_t : 5.19 min.

δ_H (300MHz; $CDCl_3$): 3.45 (s, 3H), 7.19-7.37 (m, 4H), 7.40 (t, 2H, J 7.2), 7.37-7.40 (m, 1H), 7.46 (dt, 2H, J 8.7 and 2.3), 7.73 (dt, 2H, J 8.7 and 2.2), 7.79 (d, 2H, J 8.7).

Example 15 (General procedure 1)

5 Methyl-phenyl-carbamic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-yloxy)-phenyl ester

The title compound was prepared from 4-(3-chloro-5-trifluoromethyl)-pyridine-2-yloxy)-phenol and N-methyl-N-phenylcarbonyl chloride. Prepared as described in general procedure 1.

The crude product was recrystallized (ethanol/water (78%, white crystals). HPLC-MS m/z =
10 423.1 (M+1), Rt: 5.31 min.

δ_H (300MHz; $CDCl_3$): 3.43 (s, 3H), 7.10-7.23 (m, 4H), 7.27 (t, 1H, J 6.8), 7.35 (d, 2H, J 7.5), 7.41 (t, 2H, J 7.4), 7.96 (d, 1H, J 1.9), 8.23 (bs, 1H).

Example 16 (General procedure 1)

15 Methyl-phenyl-carbamic acid 4-[4-(4-chloro-phenyl)-thiazol-2-yl]-phenyl ester

The title compound was prepared from 4-[4-(4-chloro-phenyl)-thiazol-2-yl]-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was recrystallized (ethanol/water) (59%, white crystals). HPLC-MS m/z = 421.1 (M+1), Rt: 5.85 min

20 δ_H (300MHz; $CDCl_3$): 3.45 (s, 3H), 7.18-7.25 (m, 2H), 7.27-7.33 (m, 1H), 7.34-7.41 (m, 4H), 7.43 (d, 2H, J 6.8), 7.41-7.46 (m, 1H), 7.92 (dt, 2H, J 8.6 and 2.2), 8.00 (d, 2H, J 8.7).

Example 17 (General procedure 1)

Methyl-phenyl-carbamic acid 4-pyrrol-1-yl-phenyl ester

25

The title compound was prepared from 4-pyrrol-1-yl-phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was recrystallized (ethanol/water) (27%, off-white crystals). HPLC-MS m/z = 293.2 (M+1), Rt: 4.51 min.

30 δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 6.33 (t, 2H, J 2.2), 7.03 (t, 2H, J 2.2), 7.17 (bd, 2H, J 8.3), 7.29 (d, 1H, J 6.8), 7.31-7.38 (m, 4H), 7.41 (t, 2H, J 6.8).

Example 17a (General procedure 1)

Methyl-phenyl-carbamic acid 4-imidazol-1-yl-phenyl ester

35 The title compound was prepared from 4-imidazol-1-yl-phenol and N-methyl-N-

phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (4%, clear oil). HPLC-MS m/z = 294.1 (M+1), Rt: 2.25 min.

δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.27-7.39 (m, 4H), 7.39-7.50 (m, 5H), 7.53 (bs, 1H), 8.83 (bs, 1H).

5

Example 18 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(3-chloro-5-trifluoromethyl)-pyridine-2-ylmethyl)-phenyl ester

The title compound was prepared from 4-(3-chloro-5-trifluoromethyl)-pyridine-2-ylmethyl)-phenol and N-methyl-N-phenylcarbamoyl chloride.

10

The crude product was recrystallized (ethanol/water) (74%, white crystals). HPLC-MS m/z = 421.1 (M+1), Rt: 5.23 min.

δ_H (300MHz; $CDCl_3$): 3.40 (s, 3H), 4.33 (s, 2H), 7.03 (d, 2H, J 8.3), 7.20-7.30 (m, 3H), 7.3 (d, 2H, J 7.2), 7.38 (t, 2H, J 7.2), 7.87 (d, 1H, J 1.5), 8.69 (bs, 1H).

15

Example 19 (General procedure 1)

Methyl-phenyl-carbamic acid 4-trifluoromethylsulfanyl-phenyl ester

The title compound was prepared from 4-trifluoromethylsulfanyl-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (70%, clear oil). HPLC-MS m/z = 328.0 (M+1), Rt: 5.16 min.

20

δ_H (300MHz; $CDCl_3$): 3.42 (s, 3H), 7.19 (d, 2H, J 6.4), 7.26-7.7.37 (m, 3H), 7.41 (t, 2H, J 7.9), 7.63 (d, 2H, J 8.3).

25 **Example 20** (General procedure 1)

Methyl-phenyl-carbamic acid 4-pentafluoromethyloxy-phenyl ester

The title compound was prepared from 4-pentafluoromethyloxy-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (66%, clear oil). HPLC-MS m/z = 362.0 (M+1), Rt: 5.31 min.

30

δ_H (300MHz; $CDCl_3$): 3.42 (s, 3H), 7.10-7.22 (m, 4H), 7.29 (d, 1H, J 7.2), 7.34 (d, 2H, J 7.1), 7.41 (t, 2H, J 7.1).

Example 21 (General procedure 1)

35 Methyl-phenyl-carbamic acid 4-benzyloxy-phenyl ester

The title compound was prepared from 4-benzyloxy-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was recrystallized (ethanol/water) (83%, white crystals). HPLC-MS m/z = 334.2 (M+1), Rt: 4.88 min.

5 δ_H (300MHz; $CDCl_3$): 3.41 (s, 3H), 5.03 (s, 2H), 6.92 (dt, 2H, J 9.0 and 2.2), 7.02 (d, 2H, J 8.7), 7.26-7.34 (m, 2H), 7.34-7.38 (m, 4H), 7.38-7.44, m, 4H).

Example 22 (General procedure 1)

Methyl-phenyl-carbamic acid 4-benzyl-phenyl ester

10

The title compound was prepared from 4-benzyl-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was recrystallized (ethanol/water) (56%, white crystals). HPLC-MS m/z = 318.1 (M+1), Rt: 5.05 min.

15 δ_H (300MHz; $CDCl_3$): 3.41 (s, 3H), 3.95 (s, 2H), 7.02 (d, 2H), 7.12-7.19 (m, 5H), 7.20-7.25 (m, 2H), 7.26-7.28 (m, 1H), 7.28-7.34 (m, 1H), 7.34-7.42 (m, 3H).

Example 23 (General procedure 1)

Methyl-phenyl-carbamic acid 4'-cyano-biphenyl-4-yl-ester

20 The title compound was prepared from 4-hydroxy-4'-biphenylcarbonitrile and N-methyl-N-phenylcarbamoyl chloride applying procedure 1. The crude product was recrystallized (ethanol/water) (87%, white crystals). HPLC-MS m/z = 329.2 (M+1), Rt: 4.63 min.

δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.18-7.26 (m, 2H), 7.26-7.32 (m, 2H), 7.34-7.46 (m, 4H), 7.56 (d, 2H), 7.64 (d, 2H), 7.712 (d, 2H).

25

Example 24 (General procedure 1)

Methyl-phenyl-carbamic acid 4'-bromo-biphenyl-4-yl-ester

30 The title compound was prepared from 4-bromo-4'-hydroxybiphenyl and N-methyl-N-phenylcarbamoyl chloride. The crude product was recrystallized (ethanol) (72%, white crystals). HPLC-MS m/z = 382.0 (M+1), Rt: 5.41 min.

δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.15-7.23 (d, 2H), 7.26-7.31 (m, 1H), 7.34-7.45 (m, 6H), 7.47-7.57 (m, 4H).

35 **Example 25** (General procedure 1)

Methyl-phenyl-carbamic acid biphenyl-4-yl-ester

The title compound was prepared from 4-hydroxybiphenyl and N-methyl-N-phenylcarbamoyl chloride applying. The crude product was recrystallized (ethanol) (75%, white crystals).

5 HPLC-MS m/z = 304.2 (M+1), Rt: 4.95 min.

δ_H (300MHz; $CDCl_3$): 3.45 (s, 3H), 7.19 (d, 2H), 7.26-7.37 (m, 2H), 7.37-7.46 (m, 6H), 7.55 (d, 4H).

Example 26 (General procedure 1)

10 **Methyl-phenyl-carbamic acid 4-[3-(4-chlorophenyl)-ureido]-phenyl ester**

The title compound was prepared from 4-[3-(4-chlorophenyl)-ureido]-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (33%, off-white crystals). HPLC-MS m/z = 396.1 (M+1), Rt: 4.40 min.

15 δ_H (300MHz; $CDCl_3$): 3.52 (s, 3H), 6.85-7.03 (m, 7H), 7.09 (d, 2H), 7.30-7.50 (m, 6H).

Example 27 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(4-nitro-phenoxy)-phenyl ester

20 The title compound was prepared from 4-(4-nitro-phenoxy)-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (71%, white crystals). HPLC-MS m/z = 365.0 (M+1), Rt: 4.83 min.
 δ_H (300MHz; $CDCl_3$): 3.44 (s, 3H), 7.00 (d, 2H), 7.06 (d, 2H), 7.26-7.32 (m, 1H), 7.33-7.49 (m, 4H), 8.20 (dt, 2H).

25

Example 28 (General procedure 1)

Methyl-phenyl-carbamic acid 4-heptylsulfanyl-phenyl ester

30 The title compound was prepared from 4-heptylsulfanyl-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (74%, colourless oil, HPLC-MS m/z = 358.2 (M+1), Rt: 6.21 min.
 δ_H (200MHz; $CDCl_3$): 0.87 (t, 3H), 1.15-1.50 (m, 8H), 1.50-1.75 (m, 2H), 2.86 (t, 2H), 3.41 (s, 3H), 7.04 (d, 2H), 7.15-7.50 (m, 7H).

35 **Example 29 (General procedure 1)**

Methyl-phenyl-carbamic acid 4-butoxy-phenyl ester

The title compound was prepared from 4-butoxy-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was recrystallized from ethanol (22%, white crystals). HPLC-MS
5 $m/z = 300.1$ (M+1), R_t : 5.20.

δ_H (300MHz; $CDCl_3$): 0.96 (t, 3H), 1.45 (qi, 2H), 1.74 (qi, 2H), 3.41 (s, 3H), 3.92 (t, 2H), 6.84 (d, 2H), 6.99 (d, 2H), 7.25-7.27 (m, 1H), 7.30-7.45 (m, 4H).

Example 30 (General procedure 1)10 **Methyl-phenyl-carbamic acid 4-(4-chloro-benzenesulfonyl)-phenyl ester**

The title compound was prepared from 4-(4-chloro-benzenesulfonyl)-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (46%, colourless oil). HPLC-MS $m/z = 402.1$ (M+1), R_t : 4.65 min.

15 δ_H (200MHz; $CDCl_3$): 3.41 (s, 3H), 7.21-7.35 (m, 5H), 7.39 (d, 2H), 7.46 (dt, 2H, 7.84 (dt, 2H) 7.90 (d, 2H).

Example 31 (General procedure 1)20 **Methyl-phenyl-carbamic acid 4-(4-chloromethyl-thiazol-2-yl)-phenyl ester**

The title compound was prepared from 4-(4-chloromethyl-thiazol-2-yl)-phenol and N-methyl-N-phenylcarbamoyl chloride. The aqueous phase was adjusted to PH 7.0 (phosphate buffer) before extraction with ethyl acetate. The crude product was subjected to preparative HPLC (21%, white crystals). HPLC-MS $m/z = 359.0$ (M+1), R_t : 4.60.

25 δ_H (200MHz; $CDCl_3$): 3.44 (s, 3H), 4.73 (s, 2H), 7.20 (bd, 2H), 7.26-7.48 (m, 6H), 7.92 (bd, 2H).

Example 32 (General procedure 3)30 **Methyl-phenyl-carbamic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester**

The title product was prepared from 1-(4-hydroxyphenyl)-4,4-dimethylpiperidine-2,6-dione (233 mg, 1.00 mmol)) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (343 mg, 1.00 mmol) by applying general procedure 3 (192 mg, 52%, white solid). HPLC-MS
 $m/z = 367$ (M+1); $R_t = 3.78$ min.

35 1H NMR (300MHz, $CDCl_3$): δ 1.19 (s, 6H), 2.65 (s, 4H), 3.42 (s, 3H), 7.03 (d, 2H), 7.15-7.43

(m, 7H).

Example 33 (General procedure 3)

cis-Methyl-phenyl-carbamic acid 4-(1,3-dioxo-octahydro-isoindol-2-yl)-phenyl ester

5

The title compound (293 mg, 77% yield, white crystals) was prepared from *cis*-2-(4-hydroxy-phenyl)hexahydroisoindole-1,3-dione (245 mg, 1.00 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (343 mg, 1.00 mmol). HPLC-MS *m/z* = 379 (M+1); *R*_t = 4.08 min.

10 ¹H NMR (300MHz, CDCl₃): δ 1.50 (m, 4H), 1.90 (m, 4H), 3.00 (m, 1H), 3.42 (s, 3H), 7.13-7.44 (m, 9H).

Example 34 (General procedure 3)

Methyl-phenyl-carbamic acid 4-(cyclohexanecarbonyl-amino)-phenyl ester

15

The title compound (283 mg, 80% yield, white crystals) was prepared from cyclohexanecarboxylic acid (4-hydroxyphenyl) amide (219 mg, 1.00 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (343 mg, 1.00 mmol). HPLC-MS *m/z* = 353 (M+1); *R*_t = 4.23 min.

20 ¹H NMR (300MHz, CDCl₃): δ 1.27 (m, 3H), 1.52 (m, 2H), 1.70 (m, 1H), 1.76-1.97 (m, 4H), 2.20 (m, 1H), 3.42 (s, 3H), 7.01 (d, 2H), 7.18 (d, 2H), 7.32-7.50 (m, 7H).

Example 35 (General procedure 3)

Methyl-phenyl-carbamic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester

25

The title compound (284 mg, 77% yield, white crystals) was prepared from 2-cyclohexyl-*N*-(4-hydroxy-phenyl)-acetamide (233 mg, 1.00 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (343 mg, 1.00 mmol). HPLC-MS *m/z* = 367 (M+1); *R*_t = 4.51 min

30 ¹H NMR (300MHz, CDCl₃): δ 0.87-1.03 (m, 2H), 1.07-1.38 (m, 3H), 1.60-1.92 (m, 6H), 2.14 (d, 2H), 3.41 (s, 3H), 6.97 (d, 2H), 7.26 (m, 1H), 7.30-7.44 (m, 6H), 7.55 (br.s, 1H, NH);.

Example 36 (General procedure 3)

cis/trans-Methyl-phenyl-carbamic acid 4-[(4-*tert*-butyl-cyclohexanecarbonyl)-amino]-phenyl ester

35

The title compound (353 mg, 86% yield, white crystals) was prepared from *cis/trans*-4-tert-butyl-cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide (275 mg, 1.00 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (343 mg, 1.00 mmol). HPLC-MS *m/z* = 409 (*M*+1); *R*_t = 5.28 and 5.42 min.

¹H NMR (300MHz, CDCl₃): δ 0.82 + 0.86 (2 x s, 9H), 1.03 (m, 2H), 1.23-1.70 (m, 4H), 1.82-2.22 + 2.58 (m, 4H), 3.42 (s, 3H), 7.01 (m, 2H), 7.26 (m, 1H), 7.31-7.48 (m, 7H, arom. + NH).

Example 37 (General procedure 3)

trans-Methyl-phenyl-carbamic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester

The title compound was obtained from *cis/trans*-4-tert-butyl-cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide by preparative HPLC (method B). *R*_t = 5.50 min.

¹H NMR (300MHz, CDCl₃): δ 0.85 (s, 9H), 1.03 (m, 3H), 1.52 (m, 2H), 1.88 (m, 2H), 2.01 (m, 2H), 2.12 (tt, *J* = 12.1, 3.3 Hz, 1H), 3.41 (s, 3H), 7.01 (br.d, 2H), 7.26 (m, 1H), 7.31-7.48 (m, 7H, arom. + NH).

Example 38 (General procedure 3)

cis-Methyl-phenyl-carbamic acid 4-[(4-tert-butyl-cyclohexanecarbonyl)-amino]-phenyl ester

The title compound was obtained from *cis/trans*-4-tert-butyl-cyclohexanecarboxylic acid (4-hydroxy-phenyl)-amide by preparative HPLC (method B). *R*_t = 6.34 min.

Example 39 (General procedure 3)

Methyl-phenyl-carbamic acid 4-(3,3-dimethyl-butyrylamino)-phenyl ester

The title compound (262 mg, 77% yield) was prepared from *N*-(4-hydroxyphenyl)-3,3-dimethyl-butyramide (275 mg, 1.00 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (343 mg, 1.00 mmol). HPLC-MS *m/z* = 341 (*M*+1); *R*_t = 4.15 min.

¹H NMR (300MHz, CDCl₃): δ 1.07 (s, 9H), 2.16 (s, 2H), 3.43 (s, 3H), 6.98 (d, 2H), 7.26 (m, 1H), 7.32-7.43 (m, 6H), 7.51 (br.s, 1H, NH).

Example 40 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-benzyl-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 400 ($M+1$), R_t : 4.90 min.

5 **Example 41** (General procedure 3)

Methyl-phenyl-carbamic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(3,4-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide (white
10 solid). HPLC-MS m/z = 468 ($M+1$), R_t : 5.47 min.

Example 42 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl
15 ester

The title compound was prepared from 3-(2-chloro-6-fluoro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide by applying procedure 3. HPLC-MS m/z = 452 ($M+1$), R_t : 5.15 min.

20 **Example 43** (General procedure 3)

Methyl-phenyl-carbamic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(2,6-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-
25 MS m/z = 468 ($M+1$), R_t : 5.37 min.

Example 44 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl ester
30

The title compound was prepared from 3-(2,6-dichloro-benzyl)-6-chloro-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 502 ($M+1$), R_t : 5.68 min.

35 **Example 45** (General procedure 3)

Methyl-phenyl-carbamic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(4-fluoro-benzyl)-6-chloro-7-hydroxy-4-n-propyl-2H-chromen-2-one and 3-(methyl-phenyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 480 (M+1), R_t : 5.61 min.

Example 46 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound (90 mg, 43% yield, white solid) was prepared from 7-hydroxy-3-(4-methoxy-phenyl)-4-methyl-chromen-2-one (141 mg, 0.50 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (175 mg, 0.51 mmol).

^1H NMR (300MHz, DMSO- d_6): δ 2.28 (s, 3H), 3.38 (s, 3H), 3.80 (s, 3H), 7.00 (d, 2H), 7.18-7.33 (m, 5H), 7.40-7.53 (m, 4H), 7.83 (d, 1H); HPLC-MS: m/z = 416 (M+1); R_t = 4.67 min.

Example 47 (General procedure 3)

Methyl-phenyl-carbamic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester

The title compound (120 mg, 52% yield, white solid) was prepared from 7-hydroxy-4-methyl-3-phenyl-chromen-2-one (126 mg, 0.50 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (175 mg, 0.51 mmol).

^1H NMR (300MHz, DMSO- d_6): δ 2.26 (s, 3H), 3.37 (s, 3H), 7.16-7.54 (m, 12H), 7.86 (d, 1H);

HPLC-MS: m/z = 386 (M+1); R_t = 4.69 min.

Example 48 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound (30 mg, 13% yield, white solid) was prepared from 3-(2,5-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (175 mg, 0.51 mmol).

^1H NMR (300MHz, CDCl₃): δ 2.21 (s, 3H), 3.45 (br.s, 3H), 3.72 (s, 3H), 3.79 (s, 3H), 6.74 (m, 1H), 6.92 (m, 2H), 7.12 (br.s, 2H), 7.26-7.46 (m, 5H), 7.63 (d, 1H); HPLC-MS: m/z = 446

(M+1); R_t = 4.60 min.

Example 49 (General procedure 3)

Methyl-phenyl-carbamic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2*H*-chromen-7-yl ester.

The title compound (30 mg, 13% yield, oil) was prepared from 3-(3,4-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (175 mg, 0.51 mmol).

^1H NMR (300MHz, CDCl_3): δ 2.31 (s, 3H), 3.45 (br.s, 3H), 3.89 (s, 3H), 4.02 (s, 3H), 6.83 (m, 2H), 6.94 (d, 1H), 7.12 (br.s, 2H), 7.27-7.48 (m, 5H), 7.62 (d, 1H); HPLC-MS: m/z = 446 (M+1); R_t = 4.61 min.

Example 50 (General procedure 2)

4-Chlor-phenyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-[(4-chlorophenyl)-methyl-carbamoyl]-1-methyl-3*H*-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (ethyl acetate/heptan, 1:5) (77%, white crystals). HPLC-

MS m/z = 423.1 (M+1), R_t : 5.3 min.

δ_{H} (300MHz; CDCl_3): 3.42 (s, 3H), 6.99 (d, 1H, J 8.7), 7.10-7.20 (m, 4H), 7.30 (d, 2H, J 8.3), 7.37 (d, 2H, J 8.6), 7.88 (dd, 1H, J 8.7 and 2.2), 8.42 bs, 1H).

Example 51 (General procedure 2)

4-Chlor-phenyl-methyl-carbamic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(3,5-dichloro-pyridin-2-yloxy)-phenol and 3-[(4-chlorophenyl)-methyl-carbamoyl]-1-methyl-3*H*-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (Quad flash 12, dichlormethane (67%). HPLC-MS m/z =

422.9 (M+1), R_t : 5.5 min.

δ_{H} (300MHz; CDCl_3): 3.41 (s, 3H), 7.20-7.08 (m, 4H), 7.29 (d, 2H, J 9), 7.37 (dd, 2H, J 6.4 and 2.2), 7.76 (d, 1H, J 2.3), 7.93 (d, 1H, J 2.3).

Example 52 (General procedure 2)

(4-Chloro-phenyl)-methyl-carbamic acid 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenyl

ester

The title compound was prepared from 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenol and 3-[(4-chlorophenyl)-methyl-carbamoyl]-1-methyl-3H-imidazol-1-ium iodide. The crude
5 product was purified by preparative HPLC (25%). HPLC-MS m/z = 448.2 (M+1), Rt: 5.1 min.
 δ_H (300MHz; $CDCl_3$): 3.43 (s, 3H), 7.12 (d, 2H, J 9.0), 7.19-7.35 (m, 4H), 7.39 (dd, 2H, J 6.6 and 1.8), 7.36-7.41 (m, 1H), 8.63 (d, 1H, J 0.7).

Example 53 (General procedure 1)

10 Ethyl-phenyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(4-trifluoromethyl-pyridin-2-yloxy)-phenol and *N*-ethyl-*N*-phenylcarbamoyl chloride. The crude product was subjected to flash chromatography (ethyl acetate/heptane, 1:5) (78%, white crystals). HPLC-MS m/z = 403.2 (M+1), Rt: 5.17
15 min.

δ_H (300MHz; $CDCl_3$): 1.25 (t, 3H, J 6.8), 3.83 (q, 2H, J 6.8), 6.98 (d, 1H, J 8.6), 7.12 (m, 4H), 7.32 (d, 2H, J 7.1), 7.32 (m, 1H), 7.41 (t, 2H, J 7.5), 7.87 (dt, 1H, J 8.7 and 2.7), 8.42 (bs, 1H).

20 **Example 54** (General procedure 1)

Ethyl-phenyl-carbamic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester

The title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenol and *N*-ethyl-*N*-phenylcarbamoyl chloride. The crude product was subjected to flash chromatography (ethyl
25 acetate/heptane, 1:5) (77%, white crystals. HPLC-MS m/z = 402.1 (M+1), Rt: 5.6 min.

δ_H (300MHz; $CDCl_3$): 1.25 (t, 3H, J 7.2), 3.83 (q, 2H, J 7.2), 7.01 (d, 4H, J 8.6) 7.12 (d, 2H, J 8.3); 7.30 (t, 2H, J 6.8), 7.30 (m, 1H), 7.42 (dt, 2H, J 7.5), 7.54 (d, 2H, J 8.6)

Example 55 (General procedure 2)

30 Benzyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(benzyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (ethyl acetate/heptane, (1:5) (69%, colourless oil) HPLC-MS
35 m/z = 403.2 (M+1), Rt: 5.11min

δ_H (300MHz; $CDCl_3$): 3.03 (d, 3H, J 8.0), 4.64 (d, 2H, J 24.9), 7.00 (d, 1H), 7.10-7.26 (m, 4H), 7.30-7.50 (m, 5H), 7.88 (dd, 1H), 8.43 (s, 1H).

Example 56 (General procedure 2)

5 Benzyl-methyl-carbamic acid 4-(3,5-dichloro pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(3,5-dichloro pyridin-2-yloxy)-phenol and 3-(benzyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide.

10 The crude product was subjected to flash chromatography (Quad flash 12, dichloromethane) (92%, oil). HPLC-MS m/z = 403.2 ($M+1$), Rt: 5.4 min.

δ_H (300MHz; $CDCl_3$): 3.02 (d, 3H, J 7.2), 4.60 (d, 2H, J 24.1), 7.05-7.25 (m, 4H), 7.28-7.45 (m, 5H), 7.76 (d, 1H, J 2.3) 7.95 (d, 1H, J 2.2).

Example 57 (General procedure 2)

15 tert-Butyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(tert-Butyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to preparative HPLC (34%, white crystals). HPLC-MS m/z = 369.1 ($M+1$), Rt: 5.17 min.

20 δ_H (300MHz; $CDCl_3$): 1.47 (s, 9H), 3.08 (s, 3H), 6.99 (d, 1H), 7.09-7.20 (m, 4H), 7.87 (dd, 1H), 8.43 (bs, 1H).

Example 58 (General procedure 2)

25 Isopropyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(isopropyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (ethyl acetate/heptane 1:5) (77%, white crystals). HPLC-MS m/z = 355.1 ($M+1$), Rt: 4.80 min.

30 δ_H (300MHz; $CDCl_3$): 1.21 (m, 6H), 2.91 (d, 3H), 4.49 (qi, 1H), 6.99 (d, 1H), 7.10-7.25 (m, 4H) 7.88 (dd, 1H), 8.44 (s, 1H).

Example 59 (General procedure 2)

35 Cyclohexyl-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(cyclohexyl-methyl-carbamoyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (ethyl acetate/heptane, 1:5) (80%, white crystals). HPLC-
5 MS m/z = 395.2 ($M+1$), Rt: 5.7 min.

δ_H (300MHz; $CDCl_3$): 1.13 (m, 1H), 1.50-1.30 (m, 4H), 1.68 (d, 1H, J 13.2), 1.75-1.95 (m, 4H), 2.93 (d, 2H, J 12.1), 2.90-3.00 (m, 1H), 4.02 (t, 1H, J 12.1), 6.99 (d, 1H, J 8.7), 7.10-7.17 (, 4H), 7.88 (dd, 1H, J 8.7 and 2.3), 8.44 (s, 1H).

10 **Example 60** (General procedure 1)

Dimethyl-carbamic acid 4-(3,5-dichloro pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(3,5-dichloro pyridin-2-yloxy)-phenol and dimethyl-carbamoyl chloride. The crude product was purified by preparative HPLC (38%). HPLC-MS
15 m/z = 327.0 ($M+1$), Rt: 4.7 min.

δ_H (300MHz; $CDCl_3$): 2.92 (s, 3H), 3.05 (s, 3H), 7.10-7.25 (m, 4H), 8.17 (d, 1H), 8.34 (d, 1H).

Example 61 (General procedure 1)

Pyrrolidine1-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester

20

The title compound was prepared from 4-(3,5-dichloro-pyridine-2-yloxy)-phenol and 1-pyrrolidinecarbamoyl chloride. The crude product was purified by preparative HPLC (64%, white crystals). HPLC-MS m/z = 353.0 ($M+1$), Rt: 5.00 min.

25 δ_H (300MHz; $CDCl_3$): 1.95 (qi, 4H, J 6.4), 3.48 (t, 2H, J 6.4), 3.56 (t, 2H, 6.4), 7.10 (t, 1H, J 2.7), 7.13 (t, 1H, J 2.7), 7.17 (t, 1H, J 2.0), 7.20 (t, 1H, J 2.3), 7.76 (d, 1H, J 2.6), 7.95 (d, 1H J, 2.6).

Example 62 (General procedure 2)

2,3-Dihydro-indole-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

30

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(2,3-dihydro-indole-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to column chromatography(ethyl acetate/heptane, 1:5) (73%, crystals. HPLC-MS
35 m/z = 401.1) ($M+1$), Rt: 5.5 min.

δ_H (300MHz; $CDCl_3$): 3.24 (t, 2H, J 8.4), 4.25 (t, 2H, J 8.4), 7.10 (m, 2H), 7.15-7.35 (m, 6H),

7.90 (dd, 2H, J 8.7 and 2.6), 8.44 (bs, 1H).

Example 63 (General procedure 2)

1,3-Dihydro-isoindole-2-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

5

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(1,3-dihydro-isoindole-2-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was recrystallized (ethanol) (≈ 100). HPLC-MS m/z = 401.1 ($M+1$), Rt: 5.1 min.

δ_H (300MHz; $CDCl_3$): 4.85 (s, 2H), 4.95 (s, 2H), 7.01(d, 1H, J 8.69), 7.16 (dt, 2H, J 9.4 and
10 2.7), 7.22-7.29 (m, 2H), 7.30-7.35 (m, 4H), 7.89 (dd, 1H, J 8.6 and 3.0), 8.45 (m, 1H).

Example 64 (General procedure 2)

Piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

15 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methyl-3-(piperidine-1-carbonyl)-3H-imidazol-1-ium iodide. The crude product was recrystallized (ethanol) (45%). HPLC-MS m/z = 367.02 ($M+1$), Rt: 4.9 min.

δ_H (300MHz; $CDCl_3$): 1.65 (bs, 6H), 3.58 (d, 4H, J 21.4), 6.99 (d, 1H, J 8.7), 7.10-7.24 (m, 4H), 7.88 (dd, 1H, J 8.7 and 2.2), 8.44 (bs, 1H).

20

Example 65 (General procedure 2)

2-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-
25 methyl-3-(2-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide. The crude product was purified by flash chromatography using a Quad flash 25 (ethyl acetate/heptane (1:6), (71%, white crystals). HPLC-MS m/z = 381.1 ($M+1$), Rt: 5.2 min.

δ_H (300MHz; $CDCl_3$): 1.26 (d, 3H), 1.40-1.85 (m, 6H), 3.03 (t, 1H), 4.11 (dd, 1H), 4.50-4.65
(m, 1H), 6.95-7.02 (d, 1H), 7.10-7.20 m, 4H), 7.88 (dd, 1H), 8.43 (bs, 1H).

30

Example 66 (General procedure 2)

3-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-
35 methyl-3-(3-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide. The crude product was

purified by flash chromatography using a Quad flash 25 (ethyl acetate/heptane (1:6), (75%, white crystals). HPLC-MS m/z = 381.1 ($M+1$), Rt: 5.4 min.

δ_H (300MHz; $CDCl_3$): 0.94 (d, 3H), 1.05-1.20 (m, 1H), 1.50-1.80 (m, 3H), 1.80-1.95 (m, 1H), 2.45-2.75 (dt, 1H), 2.80-3.00 (m, 1H), 4.00-4.25 (m, 2H), 6.95-7.05 (d, 1H), 7.10-7.25 (m, 4H), 7.88 (dd, 1H), 8.43 (bs, 1H).

Example 67 (General procedure 2)

4-Methyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methyl-3-(4-methyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide. The crude product was purified by flash chromatography using a Quad flash 25 (ethyl acetate/heptane (1:6), (73%, white crystals). HPLC-MS m/z = 381.1 ($M+1$), Rt: 5.4 min.

δ_H (300MHz; $CDCl_3$): 1.00 (d, 3H), 1.23 (dq, 2H), 1.52-1.65 (m, 1H), 1.70 (d, 2H), 2.75-3.05 (m, 2H), 4.15-4.35 (m, 2H), 6.99 (d, 1H), 7.05-7.20 (m, 4H), 7.88 (dd, 1H), 8.43 (s, 1H).

Example 68 (General procedure 2)

4-Benzyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methyl-3-(4-benzyl-piperidine-1-carbonyl)-3H-imidazol-1-ium iodide. The crude product was purified by flash chromatography using a Quad flash 25 (ethyl acetate/heptane (1:6), (72%, white crystals). HPLC-MS m/z = 457.2 ($M+1$), Rt: 6.0 min.

δ_H (300MHz; $CDCl_3$): 1.20-1.40 (m, 2H), 1.65-1.85 (m, 3H), 2.59 (d, 2H), 2.70-3.00 (m, 2H), 4.15-4.35 (m, 2H), 6.99 (d, 1H), 7.05-7.22 (m, 6H), 7.22-7.35 (m, 3H), 7.88 (dd, 1H), 8.43 (bs, 1H).

Example 69 (General procedure 2)

3,4-Dihydro-1H-isoquinoline-2-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(3,4-dihydro-1H-isoquinoline-2-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was recrystallized (ethanol) (53%). HPLC-MS m/z = 415.2 ($M+1$), Rt: 5.3 min.

δ_H (300MHz; $CDCl_3$): 2.96 (d, 2H, J 4.9), 3.86 (dt, 2H, J 23.3 and 6.0), 4.79 (d, 2H, J 35.4),

7.00 (d, 1H, J 8.7), 7.10-7.23 (m, 8H), 7.88 (dt, 1H, J 8.7 and 2.1), 8.44 (s, 1H).

Example 70 (General procedure 2)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester

5

The title compound was prepared from 4-(3,5-dichloro-pyridin-2-yloxy)-phenol and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was recrystallized (ethanol) (68%). HPLC-MS m/z = 415.2 ($M+1$), R_t : 5.6 min.

10 δ_H (300MHz; $CDCl_3$): 2.05 (Qi, 2H, J 6.8), 2.85 (t, 2H, J 6.8), 3.92 (t, 2H, J 6.8), 7.06 (dt, (1H, J 7.4 and 1.1), 7.11-7.19 (m, 4H), 7.21 (t, 1H, J 2.8), 7.24 (t, 1H, J 2.7), 7.77 (d, 1H, J 2.3), 7.75-7.80 (m, 1H), 7.95 (d, 1H, J 2.39).

Example 71 (General procedure 2)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenyl ester

15

The title compound was prepared from 4-(2-cyano-5-trifluoromethyl-pyridin-3-yloxy)-phenol and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was subjected to flash chromatography (Quad flash 12, dichlormethane) (43%, oil).

20 HPLC-MS m/z = 440.2 ($M+1$), R_t : 5.2 min.

δ_H (300MHz; $CDCl_3$): 2.07 (Qi, 2H, J 6.4), 2.87 (t, 2H, J 6.4), 3.94 (t, 2H, J 6.4), 7.25-7.04 m, (5H), 7.26-7.7.36 (m, 2H), 7.44 (d, 1H, J 1.5), 7.76 (bd, 1H, J 7.2), 8.64 (s, 1H).

Example 72 (General procedure 3)

25 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(3,4-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 494 ($M+1$), R_t : 5.86 min.

30

Example 73 (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-benzyl-4-methyl-2-oxo-2H-chromen-7-yl ester

35 The title compound was prepared from 3-benzyl-7-hydroxy-4-methyl-2H-chromen-2-one and

3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 426 (M+1), Rt: 5.34 min.

Example 74 (General procedure 3)

5 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(2-chloro-6-fluoro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium io-
10 dide. HPLC-MS m/z = 478 (M+1), Rt: 5.58 min.

Example 75 (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

15 The title compound was prepared from 3-(2,6-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 494 (M+1), Rt: 5.79 min.

20 **Example 76** (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,6-dichloro-benzyl)-6-chloro-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(2,6-dichloro-benzyl)-6-chloro-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium
25 iodide. HPLC-MS m/z = 530 (M+1), Rt: 6.09 min.

Example 77 (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(4-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(4-fluoro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide.
HPLC-MS m/z = 444 (M+1), Rt: 5.36 min.

Example 78 (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 6-chloro-3-(2-chloro-6-fluoro-benzyl)-4-n-propyl-2-oxo-2H-chromen-7-yl ester

- 5 The title compound was prepared from 3-(4-fluoro-benzyl)-6-chloro-7-hydroxy-4-n-propyl-2H-chromen-2-one and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 506 (M+1), R_t : 6.01 min.

Example 79 (General procedure 3)

- 10 3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound (130 mg, 59% yield, white solid) was prepared from 7-hydroxy-3-(4-methoxy-phenyl)-4-methyl-chromen-2-one (141 mg, 0.50 mmol) and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide (188 mg, 0.51 mmol).

15 ^1H NMR (300MHz, DMSO- d_6): δ 2.00 (quintet, 2H), 2.30 (s, 3H), 2.82 (t, 2H), 3.81 (s, 3H), 3.88 (t, 2H), 7.02 (d, 2H), 7.08 (m, 1H), 7.19 (m, 2H), 7.27 (d, 2H), 7.31 (dd, 1H), 7.41 (d, 1H), 7.72 (d, 1H), 7.88 (d, 1H); HPLC-MS: m/z = 442 (M+1); R_t = 5.13 min.

- 20 **Example 80** (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester

The title compound (150 mg, 73% yield, white solid) was prepared from 7-hydroxy-4-methyl-3-phenyl-chromen-2-one (126 mg, 0.50 mmol) and 3-(3,4-dihydro-2H-quinoline-1-carbonyl)-1-methyl-3H-imidazol-1-ium iodide (188 mg, 0.51 mmol).

25 ^1H NMR (300MHz, DMSO- d_6): δ 1.99 (quintet, 2H), 2.28 (s, 3H), 2.82 (t, 2H), 3.88 (t, 2H), 7.08 (m, 1H), 9.19 (m, 2H), 7.33 (m, 3H), 7.46 (m, 4H), 7.74 (d, 1H), 7.90 (d, 1H); HPLC-MS: m/z = 412 (M+1); R_t = 5.14 min.

30

Example 81 (General procedure 3)

3,4-Dihydro-2H-quinoline-1-carboxylic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound (50 mg, 21% yield, white solid) was prepared from 3-(2,5-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 3-(3,4-dihydro-2H-

35

quinoline-1-carbonyl)-1-methyl-3*H*-imidazol-1-ium iodide (188 mg, 0.51 mmol).

¹H NMR (300MHz, CDCl₃): δ 2.08 (quintet, 2H), 2.34 (s, 3H), 2.87 (t, 2H), 3.73 (s, 3H), 3.80 (s, 3H), 3.94 (t, 2H), 6.74 (m, 1H), 6.93 (m, 2H), 7.07-7.26 (m, 5H), 7.68 (d, 1H), 7.77 (br.d, 1H); HPLC-MS: *m/z* = 472 (M+1); R_t = 5.04 min.

5

Example 82 (General procedure 3)

3,4-Dihydro-2*H*-quinoline-1-carboxylic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2*H*-chromen-7-yl ester

10 The title compound (50 mg, 21% yield, oil) was prepared from 3-(3,4-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 3-(3,4-dihydro-2*H*-quinoline-1-carbonyl)-1-methyl-3*H*-imidazol-1-ium iodide (188 mg, 0.51 mmol).

¹H NMR (300MHz, CDCl₃): δ 2.08 (quintet, 2H), 2.33 (s, 3H), 2.88 (t, 2H), 3.89 (s, 3H), 4.03 (s, 3H), 4.06 (t, 2H), 6.65 (m, 2H), 6.96 (d, 1H), 7.05-7.25 (m, 5H), 7.68 (d, 1H), 7.77 (br.d, 1H); HPLC-MS: *m/z* = 472 (M+1); R_t = 5.14 min.

15

Example 83 (General procedure 2)

7-Trifluoromethyl-3,4-dihydro-2*H*-quinoline-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

20

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methyl-3-(7-trifluoromethyl-3,4-dihydro-2*H*-quinoline-1-carbonyl)-3*H*-imidazol-1-ium iodide. The crude product was subjected to preparative HPLC (66 %). HPLC-MS *m/z*: 483.1 (M+1), R_t: 5.87 min.

25 δ_H(200MHz; CDCl₃): 2.08 (Qi, 2H), 2.90 (t, 2H), 3.98 (t, 2H), 7.03 (d, 1H), 7.1-7.40 (m, 6H), 7.90 (dd, 1H), 8.18 (s, 1H), 8.44 (d, 1H).

Example 84 (General procedure 1)

Morpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and morpholine-4-carbonyl chloride. The crude product was purified by preparative HPLC (37%, white crystals). HPLC-MS *m/z* = 369.1 (M+1), R_t: 4.3 min.

35 δ_H(300MHz; CDCl₃): 3.59 (bs, 2H), 3.67 (bs, 2H), 3.76 (m, 4H), 7.01 (d, 1H, J 8.7), 7.12-7.18 (m, 2H), 7.16 (d, 2H, J 3.7), 7.90 (dd, 1H, J 8.7 and 2.6), 8.43 (s, 1H).

Example 85 (General procedure 1)

Morpholine-4-carboxylic acid 4-(3,5-dichloro-pyridin-4-yloxy)-phenyl ester

- 5 The title compound was prepared from 4-(3,5-dichloro-pyridin-4-yloxy)-phenol and morpholine-4-carbonyl chloride. The crude product was purified by preparative HPLC (66%, white crystals. HPLC-MS m/z = 368.9 (M+1), Rt: 4.0 min.
 δ_H (300MHz; $CDCl_3$): 3.57 (bs, 2H), 3.65 (bs, 2H) 3.74 (m, 4H), 6.83 (d, 1H, J 9), 6.83 (m, 1H) 7.08 (d, 1H, J 9), 7.08 (m, 1H), 8.56 (s, 2H).

10

Example 86 (General procedure 1)

Morpholine-4-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester

- The title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenol and morpholine-4-carbonyl chloride. The crude product was purified by preparative HPLC (74%, white crystals). HPLC-MS m/z = 368.1 (M+1), Rt: 4.85 min.
 δ_H (300MHz; $CDCl_3$): 3.59 (bs, 2H), 3.68 (bs, 2H), 3.75 (m, 4H), 7.04 (d, 2H, J 6.4), 7.04 (m, 2H), 7.14 (d, 1 H, J 9), 7.13 (m, 1H), 7.57 (d, 2H, J 8.3).

- 20
- Example 87**
- (General procedure 1)

Morpholine-4-carboxylic acid 4-tert-butoxy-phenyl ester

- The title compound was prepared from 4-(tert-butoxy)-phenol and 4-morpholine-carbamoyl chloride. The crude product was recrystallized (ethanol) (69%, crystals); mp: 128.8-129.5°C.
25 HPLC-MS m/z = 280.1 (M+1), Rt: 3.6 min.
 δ_H (300MHz; $CDCl_3$): 1.33 (s, 9H,), 3.58 (bs, 2H), 3.66 (bs, 2H), 6.8-7.1 (m, 4H)

Example 88 (General procedure 1)

Morpholine-4-carboxylic acid 4-(3,5-dichloro-pyridin-2-yloxy)-phenyl ester

30

The title compound was prepared from 4-(3,5-dichloro-pyridin-2-yloxy)-phenol and morpholine-4-carbonyl chloride. The crude product was purified by preparative HPLC (45%, crystals. HPLC-MS m/z = 369.1 (M+1), Rt: 4.3 min.

- 35
- Example 89**
- (General procedure 1)

Morpholine-4-carboxylic acid 3-(4-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(4-fluorobenzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and morpholine-4-carbonyl chloride. The crude product was recrystallized (ethanol)

5 (78%, crystals). HPLC-MS m/z = 398.0 (M+1), Rt: 4.3 min.

δ_H (200MHz; DMSO- d_6): 2.47 (s, 3H), 3.45 (bs, 2H), 3.56-3.72 (m, 6H), 3.96 (s, 2H), 7.08 (t, 2H), 7.16-7.35 (m, 4H, 7.83 (d, 1H).

Example 90 (General procedure 1)

10 Morpholine-4-carboxylic acid 4-(5,7-bis-trifluoromethyl-[1,8]naphthypyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5,7-bis-trifluoromethyl-[1,8]naphthypyridin-2-yloxy)-phenol and morpholine-4-carbonyl chloride. The crude product was tested without purification

15 tion (\approx 100%). HPLC-MS m/z = 488.0 (M+1), Rt: 5.0 min.

Example 91 (General procedure 3)

Morpholine-4-carboxylic acid 3-(3,4-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

20 The title compound was prepared from 3-(3,4-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 3-(morpholine-4-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. HPLC-MS m/z = 448 (M+1), Rt: 4.73 min.

Example 92 (General procedure 3)

25 Morpholine-4-carbamic acid 3-(benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS m/z = 380 (M+1), Rt: 4.05 min.

30

Example 93 (General procedure 3)

Morpholine-4-carbamic acid 3-(2-chloro-6-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

35

The title compound was prepared from 3-(2-chloro-6-fluoro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS $m/z = 432$ (M+1), Rt: 4.36 min.

5 **Example 94** (General procedure 3)

Morpholine-4-carbamic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(2,6-dichloro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS
10 $m/z = 448$ (M+1), Rt: 4.59 min.

Example 95 (General procedure 3)

Morpholine-4-carbamic acid 3-(2,6-dichloro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

15 The title compound was prepared from 3-(2,6-dichloro-benzyl)-6-chloro-7-hydroxy-4-methyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS $m/z = 484$ (M+1), Rt: 4.99 min.

Example 96 (General procedure 3)

20 Morpholine-4-carbamic acid 3-(4-fluoro-benzyl)-4-methyl-2-oxo-2H-chromen-7-yl ester

The title compound was prepared from 3-(4-fluoro-benzyl)-7-hydroxy-4-methyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS $m/z = 398$ (M+1), Rt: 4.12 min.

25

Example 97 (General procedure 3)

Morpholine-4-carbamic acid 6-chloro-3-(2,6-dichloro-benzyl)-4-n-propyl-2-oxo-2H-chromen-7-yl ester

30 The title compound was prepared from 3-(4-fluoro-benzyl)-6-chloro-7-hydroxy-4-n-propyl-2H-chromen-2-one and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide. HPLC-MS $m/z = 460$ (M+1), Rt: 4.90 min.

Example 98 (General procedure 3)

35 Morpholine-4-carboxylic acid 4-methyl-2-oxo-3-phenyl-2H-chromen-7-yl ester

The title compound (75 mg, 41% yield, crystals) was prepared from 7-hydroxy-4-methyl-3-phenyl-chromen-2-one (126 mg, 0.50 mmol) and 1-methyl-3-(morpholine-4-carbonyl)-3*H*-imidazol-1-ium iodide (165 mg, 0.51 mmol).

¹H NMR (300MHz, CDCl₃): δ 2.27 (s, 3H), 3.45 (br.s, 2H), 3.62 (br.s, 2H), 3.68 (m, 4H), 7.23 (dd, 1H), 7.32 (m, 3H), 7.43 (m, 3H), 7.87 (d, 1H); HPLC-MS: *m/z* = 366 (M+1); R_t = 3.79 min.

Example 99 (General procedure 3)

Morpholine-4-carboxylic acid 3-(4-methoxy-phenyl)-4-methyl-2-oxo-2*H*-chromen-7-yl ester

The title compound (120 mg, 61% yield, crystals) was prepared from 7-hydroxy-3-(4-methoxy-phenyl)-4-methyl-chromen-2-one (141 mg, 0.50 mmol) and 1-methyl-3-(morpholine-4-carbonyl)-3*H*-imidazol-1-ium iodide (165 mg, 0.51 mmol).

¹H NMR (300MHz, DMSO-*d*₆): δ 2.29 (s, 3H), 3.43 (br.s, 2H), 3.61 (br.s, 2H), 3.66 (m, 4H), 3.81 (s, 3H), 7.01 (d, 2H), 7.20-7.30 (m, 4H), 7.84 (d, 1H); HPLC-MS: *m/z* = 396 (M+1); R_t = 3.80 min.

20

Example 100 (General procedure 3)

Morpholine-4-carboxylic acid 3-(3,4-dimethoxy-phenyl)-4-methyl-2-oxo-2*H*-chromen-7-yl ester

The title compound (120 mg, 56% yield) was prepared from 3-(3,4-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 1-methyl-3-(morpholine-4-carbonyl)-3*H*-imidazol-1-ium iodide (165 mg, 0.51 mmol).

¹H NMR (300MHz, DMSO-*d*₆): δ 2.29 (s, 3H), 3.46 (br.s, 2H), 3.62 (br.s, 2H), 3.67 (m, 4H), 3.74 (s, 3H), 3.81 (s, 3H), 6.83 (dd, 1H), 6.92 (d, 1H), 7.02 (d, 1H), 7.22 (dd, 1H), 7.30 (d, 1H), 7.85 (d, 1H); HPLC-MS: *m/z* = 426 (M+1); R_t = 3.80 min.

30

Example 101 (General procedure 3)

Morpholine-4-carboxylic acid 3-(2,5-dimethoxy-phenyl)-4-methyl-2-oxo-2*H*-chromen-7-yl ester

35

The title compound (120 mg, 56% yield) was prepared from 3-(2,5-dimethoxy-phenyl)-7-hydroxy-4-methyl-chromen-2-one (156 mg, 0.50 mmol) and 1-methyl-3-(morpholine-4-carbonyl)-3H-imidazol-1-ium iodide (165 mg, 0.51 mmol).

¹H NMR (300MHz, CDCl₃): δ 2.23 (s, 3H), 3.60 (br.s, 2H), 3.74 (br.s, 2H + s, 3H), 3.80 (m, 4H + s, 3H), 6.73 (m, 1H), 6.92 (m, 2H), 7.13 (m, 2H), 7.64 (d, 1H); HPLC-MSA: *m/z* = 426 (M+1); R_t = 3.69 min.

Example 102 (General procedure 2)

2.6-dimethyl-morpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-(2,6-dimethyl-morpholine-4-carbonyl)-1-methyl-3H-imidazol-1-ium iodide. The crude product was purified by preparative HPLC (39%, colorless oil). HPLC-MS *m/z* = 397.1 (M+1), R_t: 4.64 min.

δ_H(200MHz; CDCl₃): 1.22 (s, 3H), 1.25 (s, 3H), 2.50-3.00 (m, 2H), 3.55-3.9 (m, 2H), 4.00-3.30 bd, 2H), 6.90-7.50 (m, 5H), 7.91 (dd, 1H), 8.44 (bs, 1H).

Example 103

Dimethylcarbamic acid benzotriazol-1-yl ester

Example 104

2-Oxo-*N-p*-tolylacetamide O-(cyclohexylcarbamoyl)-oxime

Example 105

10,11-Dihydro-dibenzo[a,d]cyclohepten-5-one O-cyclohexylcarbamoyl-oxime

Example 106

1-(4-Chlorophenyl)-non-1-en-3-one O-cyclohexylcarbamoyl-oxime

Example 107

1,7,7-Trimethyl-bicyclo[2.2.1]heptan-2-one O-cyclohexylcarbamoyl-oxime

Example 108

1-(4-Bromophenyl)-6-methyl-hept-1-en-3-one O-isopropylcarbamoyl-oxime

Example 109

1-(4-Chloro-phenyl)-non-1-en-3-one O-isopropylcarbamoyl-oxime

5 Example 110

4-Bromo-2-(4-chlorobenzyl)-2H-pyrazole-3-carbaldehyde O-methylcarbamoyl-oxime

Example 111

1-(4-Bromophenyl)-6-methyl-hept-1-en-3-one O-propylcarbamoyl-oxime

10

Example 112

N-(4-Fluorobenzylcarbamoyloxy)-isobutyrimidoyl chloride

Example 113

15 N-(2-Hydroxy-2-phenylethylcarbamoyloxy)-isobutyrimidoyl chloride

Example 114

Dimethylcarbamic acid 6-methanesulfonyl-indol-1-yl ester

20 Example 115

1-Biphenyl-4-yl-3-methylamino-propenone-cyclohexyl-carbamic acid

Example 116

Dimethylcarbamic acid 4-oxo-1,2,3-benzotriazin-3-yl ester

25

Example 117

[1-(3-Chloro-5-trifluoromethyl-pyridin-2-yl)-1H-pyrrol-2-yl]-carbamic acid 4-chloro-phenyl ester

Example 118 (General procedure 6)

30 N-Methyl-N-phenyl-5-hexylsulfanyl-3-p-tolyl-[1,2,4]triazole-1-carboxamide

The title compound was prepared from p-toluoyl chloride, 1-bromo-hexane and N-methyl-phenyl-carbamoyl chloride. The crude product was recrystallized (ethanol/ water); HPLC-MS $m/z = 409.2$ ($M+1$), R_t : 6.66 min.

35 δ_H (300MHz; [2H_6]DMSO): 0.87 (t, 3H), 1.20-1.35 (m, 4H), 1.35-1.48 (m, 2H), 1.72 (q, 2H), 2.99 (s, 3H), 3.25 (t, 2H), 3.46 (s, 3H), 7.13-7.30 (m, 5H), 7.36 (t, 2H), 7.52 (d, 2H).

Example 119 (General procedure 6)**N-Methyl-N-phenethyl-5-ethyl-3-(4-chlorophenyl)-[1,2,4]triazole-1-carboxamide**

- 5 The title compound was prepared from 4-chlorobenzoyl chloride, 1-bromo-ethane and *N*-methyl-phenethyl-carbamoyl chloride. The crude product was subjected to preparative HPLC; HPLC-MS m/z = 401.1 ($M+1$), R_t : 6.17 min.
- δ_H (300MHz; $[^2H_6]$ DMSO): 1.37 (t, 3H), 2.98 (t, 2H), 3.11 (bs, 3H), 3.23 (Q, 2H), 3.80 (m, 2H), 7.10-7.40 (m, 5H), 7.58 (d, 2H), 8.03 (d, 2H).

10

Example 120 (General procedure 6)**[3-(4-Chlorophenyl)-5-methylsulfanyl-[1,2,4]triazol-1-yl]-morpholin-4-yl-methanone**

- The title compound was prepared from 4-chlorobenzoyl chloride, methyl iodide and morpholine-4-carbonyl chloride. The crude product was used without further purification; HPLC-MS m/z = 339.1 ($M+1$), R_t : 4.68 min.
- δ_H (300MHz; $[^2H_6]$ DMSO): 2.68 (s, 3H), 3.25-3.65 (m, 4H), 3.65-3.85 (m, 4H), 7.58 (d, 2H), 8.03 (d, 2H).

20 **Example 121** (General procedure 6)**N,N-Dimethyl-5-methylsulfanyl-3-naphthalen-2-yl-[1,2,4]triazole-1-carboxamide**

- The title compound was prepared from 2-naphthoyl chloride, methyl iodide and dimethyl carbamoyl chloride. The crude product was recrystallized from ethanol; HPLC-MS m/z = 313.2 ($M+1$), R_t : 4.91 min.
- 25 δ_H (300MHz; $[^2H_6]$ DMSO): 2.74 (s, 3H), 3.18 (s, 6H), 7.53-7.65 (m, 2H), 7.92-8.00 (m, 1H), 8.00-8.17 (m, 3H), 8.62 (s, 1H).

Example 122 (General procedure 6)**N,N-Dimethyl-3-(4-chloro-phenyl)-5-ethylsulfanyl-[1,2,4]triazole-1-carboxamide**

30

- The title compound was prepared from 2-naphthoyl chloride, methyl iodide and dimethyl carbamoyl chloride. The crude product was recrystallized from ethanol; HPLC-MS m/z = 311.0 ($M+1$), R_t : 5.13 min.
- δ_H (300MHz; $[^2H_6]$ DMSO): 1.39 (t, 3H), 3.13 (bs, 6H), 3.26 (q, 2H), 7.58 (d, 2H), 8.03 (d, 2H).

35

Example 123 (General procedure 6)

N,N-Dimethyl-3-biphenyl-4-yl-5-methylsulfanyl-[1,2,4]triazole-1-carboxamide

The title compound was prepared from 3-biphenylcarbonyl chloride, methyl iodide and dimethyl carbamoyl chloride. The crude product was without further purification; HPLC-MS m/z = 339.1 (M+1), R_f: 5.27 in.

δ_H (300MHz; [²H₆]DMSO): 2.70 (s, 3H), 3.16 (bs, 6H), 7.40 (t, 1H), 7.50 (t, 2H), 7.73 (d, 2H), 7.82 (d, 2H), 8.12 (d, 2H).

Example 124

N,N-Dimethyl-3-(4-chloro-phenyl)-5-methylsulfanyl-[1,2,4]triazole-1-carboxamide

Example 126

N-(4-Chlorophenyl)-3-(4-chlorophenyl)-5-(3-hydroxypropyl)-pyrazole-1-carboxamide.

15

Example 127

5-Morpholin-4-yl-3-(4-phenoxyphenyl)-pyrazole-1-carboxylic acid phenethylamide

Example 128

N-Phenyl-4-(4-chlorobenzenesulfonyl)-5-(4-chlorophenyl)-pyrazole-1-carboxamide

20

Example 129

N-(3,4-Dichlorophenyl)-2-phenyl-benzimidazole-1-carboxamide

Example 130

Dimethyl-carbamic acid 5-isopropylsulfanyl-4-(3-trifluoromethyl-phenyl)-4H-[1,2,4]triazol-3-yl ester

25

Example 131

Dimethyl carbamic acid 1-benzyl-2-oxo-3,5-diphenyl-1,2-dihydro-pyridin-4-yl ester

30

Example 132

Dimethyl-carbamic acid 7-methoxy-1-methyl-2-oxo-1,2-dihydro-quinolin-4-yl ester

Example 133

Dimethyl carbamic acid 4-(3-chloro-phenyl)-5-(4-methyl-benzylsulfanyl)-4H-[1,2,4]triazol-3-yl

35

ester

Example 134 (General procedure 7)

5 4-Methyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methylpiperazine. White solid, m.p. 249 - 250 °C (decomp.); HPLC-MS m/z = 458 (M+H), R_t: 3.15 min.; ¹H NMR (DMSO-*d*₆): δ 11.56 (br, 1H, NH), 8.61 - 8.54 (br, 1H, py-H6), 8.30 - 8.20 (m, 1H, py-H4), 7.32 - 7.19 (d+ br s, 5H, py-H3 + C₆H₄), 4.5 - 3.0 (br, 10H, 8 CH + H₂O), 2.79 (s, 3H, CH₃); IR (KBr): ν 1713 (C=O).

Example 135 (General procedure 7)

15 4-Benzyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-benzylpiperazine. White solid, m.p. 229 - 231 °C; HPLC-MS m/z = 458 (M+H), R_t: 3.15 min.; ¹H NMR (DMSO-*d*₆): δ 11.17 (br, 1H, NH), 8.61 - 8.54 (br, 1H, py-H6), 8.30 - 8.20 (m, 1H, py-H4), 7.74 - 7.59 (m, 2H, arom.), 7.52 - 7.41 (m, 3H, arom.), 7.31 - 7.17 (d+ br s, 5H, py-H3 + C₆H₄), 4.5 - 4.0 (br, 4H, CH₂ + 2 CH), 3.8 - 3.0 (br, 8H, 6 CH + H₂O); IR (KBr): ν 1720 (C=O).

Example 136 (General procedure 7)

25 4-(2-Hydroxyethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-(2-hydroxyethyl)-piperazine. White solid, m.p. 255 °C; HPLC-MS m/z = 412 (M+H), R_t: 2.12 min.; ¹H NMR (DMSO-*d*₆): δ 10.66 (br, 1H, NH), 8.61 - 8.54 (br, 1H, py-H6), 8.30 - 8.20 (m, 1H, py-H4), 7.32 - 7.20 (d+ br s, 5H, py-H3 + C₆H₄), 5.39 (br, 1H, OH), 4.18 (br, 2H), 3.87 - 3.70 (br t, 2H), 3.70 - 3.02 (br, 8H + water); IR (KBr): ν 1714 (C=O).

Example 137 (General procedure 7)

35 4-(2-Oxo-2-pyrrolidin-1-yl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-(pyrrolidinocarbonylmethyl)-piperazine. White solid, m.p. 224 °C; HPLC-MS m/z = 479 (M+H), R_t : 2.73 min.; ^1H NMR (DMSO- d_6): δ 10.43 (br, 1H, NH), 8.62 - 8.52 (br, 1H, py-H6), 8.31 - 8.20 (m, 1H, py-H4), 7.34 - 7.18 (d+ br s, 5H, py-H3 + C₆H₄), 4.40 - 3.05 (br, CH₂ at 4.26 + water at 3.37 + N-C-H), 2.03 - 1.72 (m, 4H, CH₂).

Example 138 (General procedure 7)

4-Phenyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-phenyl-piperazine (0.5 mmol). The crude product (0.15 g) was partitioned between ethyl acetate (4 ml) and ethyldiisopropylamine (0.04 ml) dissolved in water (4 ml). The organic layer was washed with water (2 x 5 ml), dried over sodium sulfate. The drying agent was filtered off, and the solvent was removed from the filtrate *in vacuo* to give the title compound (0.095 g). White crystals, m.p. 117 °C; HPLC-MS m/z = 444 (M+H), R_t : 5.12 min.; ^1H NMR (DMSO- d_6): δ 8.61 - 8.55 (br, 1H, py-H6), 8.29 - 8.20 (m, 1H, py-H4), 7.32 - 7.14 (m, 7H, py-H3 + C₆H₄ + 2 arom.), 7.05 - 6.71 (m, 3H, arom.), 3.83 - 3.51 (br, 4H, 2 CH₂), 3.28 - 2.99 (m, 4H, 2 CH₂).

20

Example 139 (General procedure 8)

Methyl-phenyl-carbamic acid pyrazol-1-yl ester

The title compound was prepared from 1-hydroxypyrazole and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad flash 25, EtOAc-heptane) (79%, oil which slowly crystallizes). HPLC-MS m/z = 218.1 (M+1), R_t : 2.82 min. Mp 63-67 °C

25

δ_{H} (300MHz; CDCl₃): 3.45 (bs, 3H), 6.28 (s, 3H), 7.30-7.47 (m, 7H).

Example 140 (General procedure 8)

30 Methyl-phenyl-carbamic acid 4-bromo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-bromopyrazole and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad flash 25, EtOAc-heptane) (89%, oil). HPLC-MS m/z = 298.0 (M+1), R_t : 3.77 min.

δ_{H} (300MHz; CDCl₃): 3.44 (bs, 3H), 7.32-7.47 (m, 7H).

35

Example 141 (General procedure 8)

Methyl-phenyl-carbamic acid 3,4,5-tribromo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3,4,5-bromopyrazole and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad
5 flash 25, EtOAc-heptane) (93%, colorless crystals). HPLC-MS m/z = 455.8 (M+1), Rt: 4.88 min. Mp 115-119 °C
 δ_H (300MHz; $CDCl_3$): 3.44 (bs, 3H), 7.36-7.48 (m, 5H).

Example 142 (General procedure 8)

10 Methyl-phenyl-carbamic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3-(4-methoxyphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad flash 25, EtOAc-heptane) (93%, oil). HPLC-MS m/z = 346.1 (M+Na), Rt: 4.21 min.
 δ_H (300MHz; $CDCl_3$): 3.45 (bs, 3H), 3.84 (s, 3H), 6.50 (d, 1H), 6.91 (d, 2H), 7.30-7.48 (m, 6H),
15 7.70 (d, 2H).

Example 143 (General procedure 8)

Methyl-phenyl-carbamic acid imidazol-1-yl ester

The title compound was prepared from 1-hydroxyimidazole and N-methyl-N-phenylcarbonyl
20 chloride. The crude product was subjected to flash chromatography (Quad flash 25, EtOAc-heptane) (68%, oil). HPLC-MS m/z = 218.1 (M+1), Rt: 1.53 min.
 δ_H (300MHz; $CDCl_3$): 3.45 (bs, 3H), 7.00 (bs, 1H), 7.05 (bs, 1H), 7.32-7.49 (m, 5H), 7.55 (bs, 1H).

25 **Example 144** (General procedure 8)

Methyl-phenyl-carbamic acid [1,2,3]triazol-1-yl ester

The title compound was prepared from 1-hydroxy-1,2,3-triazole and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to flash chromatography (Quad
flash 25, EtOAc-heptane) (80%, oil). HPLC-MS m/z = 219.1 (M+1), Rt: 2.50 min. Mp 105-106
30 °C.
 δ_H (300MHz; $CDCl_3$): 3.46 (bs, 3H), 7.00 (bs, 1H), 7.05 (bs, 1H), 7.31-7.48 (m, 5H), 7.62 (bs, 1H), 7.74 (s, 1H).

Example 145 (General procedure 7)

35 4-(Isopropylcarbonyl-methyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-

yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 1-methyl-piperazine. White solid, m.p. 234 - 235 °C; HPLC-MS m/z = 466 (M+H), R_t: 2.75 min.; ¹H NMR (DMSO-d₆): δ 10.50 (br, 1H, NH), 8.66 - 8.54 (br, 2H, NH + py-H6), 8.30 - 8.20 (dd, 1H, py-H4), 7.31 - 7.20 (d+ br s, 5H, py-H3 + C₆H₄), 4.00 - 3.82 (br m, 5H, methin + 4 CH), 3.70 - 3.09 (br, 9H, 6H + water), 1.11 (d, 6H, CH₃).

Example 146 (General procedure 9)

4-Cyclopentyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-cyclopentyl-piperazine. White solid, m.p. 294 - 295 °C; HPLC-MS m/z = 436 (M+H), R_t: 2.92 min.; ¹H NMR (DMSO-d₆): δ 11.15 (br, 1H, NH), 8.60 - 8.55 (br, 1H, py-H6), 8.29 - 8.20 (m, 1H, py-H4), 7.32 - 7.21 (d+ br s, 5H, py-H3 + C₆H₄), 4.35 - 3.98 (br, 2H), 3.72 - 3.37 (br m, 5H), 3.29 - 2.97 (br, 2H), 2.12 - 1.45 (br m, 8H).

Example 147 (General procedure 9)

4-Butyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-butyl-piperazine. White solid, m.p. 221 - 222 °C; HPLC-MS m/z = 424 (M+H), R_t: 2.94 min.; ¹H NMR (DMSO-d₆): δ 10.86 (br, 1H, NH), 8.60 - 8.55 (br, 1H, py-H6), 8.29 - 8.21 (m, 1H, py-H4), 7.31 - 7.22 (d+ br s, 5H, py-H3 + C₆H₄), 4.32 - 4.03 (br, 2H), 3.65 - 3.44 (br m, 4H), 3.28 - 2.97 (br, 4H), 1.81 - 1.60 (br m, 2H), 1.45 - 1.22 (br, 4H), 0.92 (t, 3H, CH₃).

Example 148 (General procedure 1)

4-(Methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester

The title product was prepared from 4-Hydroxy-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and N-methyl-N-phenylcarbamoyl chloride. The crude product was recrystallized from

methanol (74%, white crystals).

¹H NMR (CDCl₃): 8.13 (d, 2H), 7.38-7.55 (m, 6H), 7.30 (t, 1H), 3.37 (s, 3H), 2.89 (s, 4H).

Example 149 (General procedure 10)

5 4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-hydroxymethylpiperidine. The crude product was subjected to flash chromatography (ethyl acetate/heptane, 1:2 → 2:1) (78%, light yellow oil). The purified product was crystallized from ethyl acetate/heptane (51%, white solid). HPLC-MS: *m/z* = 397.1 (M+1); *R_t*: 4.08 min.

¹H NMR (CDCl₃): δ 8.44 (s, 1H), 7.88 (dd, 1H), 7.17 (d, 2H), 7.13 (d, 2H), 7.00 (d, 2H), 4.45-4.20 (bs, 2H), 3.55 (t, 2H), 3.10-2.75 (m, 2H), 1.95-1.65 (m, 3H), 1.47 (t, 1H), 1.29 (dq, 2H).

15 **Example 150** (General procedure 10)

4-Oxo-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-piperidone monohydrate. The solvent used was a mixture of dichloromethane and dimethylformamide (1:1). The crude product was subjected to preparative HPLC (7%, oil). HPLC-MS: *m/z* = 381.1 (M+1); *R_t*: 4.17 min.

Example 151 (General procedure 10)

25 4-[5-(4-Dimethylamino-phenyl)-1H-pyrazol-3-yl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and N,N-dimethyl-4-[3-(4-piperidiny)-1H-pyrazol-5-yl]aniline. The solvent used was a mixture of dichloromethane and dimethylformamide (1:2). The crude product was subjected to preparative HPLC (4%, oil). HPLC-MS: *m/z* = 552.2 (M+1); *R_t*: 4.17 min.

¹H NMR (CDCl₃): 8.44 (d, 1H), 7.89 (dd, 1H), 7.53 (d, 2H), 7.22-7.10 (m, 4H), 7.00 (d, 1H), 6.73 (d, 2H), 6.27 (s, 1H), 4.45-4.20 (ds, 2H), 3.25-2.95 (m, 3H), 2.88 (s, 6H), 2.15-2.00 (m, 2H), 1.90-1.70 (dq, 2H).

35 **Example 152** (General procedure 10)

4-(5-Furan-2-yl-1H-pyrazol-3-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-[3-(2-furyl)-1H-pyrazol-5-yl]piperidine. The solvent used was a mixture of dichloromethane and dimethylformamide (1:2). The crude product was subjected to preparative HPLC (4%, oil).

HPLC-MS: m/z = 499.1 (M+1); R_t : 4.60 min.

^1H NMR (CDCl_3): 8.44 (d, 1H), 7.89 (dd, 1H), 7.45 ((dd, 1H), 7.24-7.10 (m, 4H), 6.99 (d, 1H), 6.62 (d, 1H), 6.48 (dd, 1H), 6.34 (s, 1H), 4.40-4.25 (bs, 2H), 3.25-2.85 (m, 3H), 2.08 (d, 2H), 1.79 (dq, 2H).

Example 153 (General procedure 10)

4-Benzylamino-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and benzyl-piperidin-4-yl-amine. The crude product was subjected to preparative HPLC (20%, oil).

HPLC-MS: m/z = 472.2 (M+1); R_t : 3.34 min.

Example 154 (General procedure 10)

4-(3,4-Dihydro-1H-isoquinolin-2-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 2-Piperidin-4-ylmethyl-1,2,3,4-tetrahydro-isoquinoline. The crude product was subjected to preparative HPLC (12%, oil). HPLC-MS: m/z = 512.2 (M+1); R_t : 3.21 min.

Example 155 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(1,3,5-trimethyl-1H-pyrazol-4-ylmethyl)-phenyl ester

The title product was prepared from 4-[(1,3,5-trimethyl-1H-pyrazol-4-yl)methyl]phenol and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to preparative HPLC (88%, oil). HPLC-MS: m/z = 350.0 (M+1); R_t : 3.52 min. (66% purity).

Example 156 (General procedure 10)

3-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl es-

ter

The title product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-hydroxymethyl-piperidine. The crude product was subjected to flash chromatography (Quad Flash 12, ethyl acetate/heptane 1:2). (56%, colourless oil). HPLC-MS: $m/z = 397.1$ ($M+1$); R_t : 4.19 min.

^1H NMR ($\text{DMSO}-d_6$, 90 °C): 8.37 (s, 1H), 8.00 (dd, 1H), 6.95-7.10 (m, 5H), 4.09 (bs, 1H), 3.91 (dd, 1H), 3.78 (d, 1H), 3.21 (m, 1H), 3.15 (t, 1H), 2.80-2.90 (m, 1H), 2.65 (t, 1H), 1.45-1.70 (m, 3H), 1.25-1.45 (m, 1H), 1.08 (q, 1H).

Example 157 (General procedure 10)

3-Hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 3-hydroxy-piperidine. The crude product was subjected to flash chromatography (Quad Flash 12, ethyl acetate/heptane (1:2) (56%, colourless oil). HPLC-MS: $m/z = 383.0$ ($M+1$); R_t : 3.98 min.

^1H NMR (CDCl_3): 8.43 (s, 1H), 7.89 (dd, 1H), 7.23-7.10 (m, 4H), 7.00 (d, 1H), 3.80-4.00 (m, 2H), 3.60-3.80 (m, 1H), 3.25-3.50 (m, 2H), 1.80-2.05 (m, 2H), 1.50-1.70 (m, 3H).

Example 158 (General procedure 10)

4-Benzyl-4-hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 4-benzyl-4-hydroxypiperidine. The crude product was subjected to flash chromatography (Quad Flash 12, ethyl acetate/heptane (1:2) (48%, colourless oil). HPLC-MS: $m/z = 473.0$ ($M+1$); R_t : 5.04 min.

^1H NMR (CDCl_3): 8.44 (s, 1H), 7.89 (dd, 1H), 7.28-7.45 (m, 3H), 7.10-7.24 (m, 6H), 7.00 (d, 1H), 4.00-4.14 (bs, 2H), 3.15-3.45 (dt, 2H), 2.81 (s, 2H), 1.66-1.85 (dt, 2H), 1.55-1.65 (m, 2H), 1.27 (s, 1H).

Example 159 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(2-cyano-ethyl)-phenyl ester

The title product was prepared from 3-(4-hydroxyphenyl)proprionitrile and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (11%, colourless oil). HPLC-MS: m/z = 281.2 (M+1); R_t : 3.75 min, purity 80%.

^1H NMR (CDCl_3): 7.26-7.50 (m, 7H), 7.00-7.15 (m, 2H), 3.42 (s, 3H), 2.94 (t, 2H), 2.59 (t, 2H).

Example 160 (General procedure 1)

Methyl-phenyl-carbamic acid 4-([1,2,3,4]thiatriazol-5-ylamino)-phenyl ester

The title product was prepared from 4-(1,2,3,4-thiatriazol-5-ylamino)phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (10%, light brown solid material). HPLC-MS: m/z = 328.0 (M+1); R_t : 3.69 min, purity 81%.

^1H NMR (CDCl_3): 7.61 (d, 2H), 7.38-7.52 (m, 4H), 7.19-7.35 (m, 2H), 6.86-6.95 (m, 1H), 3.35 (s, 3H), 3.33 (s, 1H).

Example 161 (General procedure 1)

Methyl-phenyl-carbamic acid 4-pentyl-phenyl ester

The title product was prepared from 4-pentylphenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (4%, light brown oil). HPLC-MS:

m/z = 298.2 (M+1); R_t : 5.61 min.

^1H NMR (CDCl_3): 7.31-7.50 (m, 4H), 7.18-7.30 (m, 1H), 7.13 (d, 2H), 7.00 (d, 2H), 3.42 (s, 3H), 2.57 (t, 2H), 1.58 (q, 2H), 1.20-1.40 (m, 4H), 0.87 (t, 3H).

Example 162 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(2-methoxy-ethyl)-phenyl ester

The title product was prepared from 4-(2-methoxyethyl)phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (4%, light yellow oil). HPLC-MS: m/z = 286.1 (M+1); R_t : 4.01 min.

^1H NMR (CDCl_3): 7.30-7.48 (m, 4H), 7.12-7.30 (m, 3H), 7.03 (d, 2H), 3.56 (t, 2H), 3.42 (s, 3H), 3.33 (s, 3H), 2.85 (t, 2H).

Example 163 (General procedure 10)

4-Hydroxy-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-hydroxypiperidine. The crude product was subjected to flash chromatography (ethyl acetate/heptane (1:1) (48%, light yellow oil). HPLC-MS: m/z = 383.0 (M+1); R_t : 3.88 min. purity 93%.

5 ^1H NMR (CDCl_3): 8.44 (s, 1H), 7.89 (dd, 1H), 7.08-7.20 (m, 4H), 6.99 (d, 1H), 3.85-4.10 (m, 3H), 3.20-3.45 (m, 2H), 1.85-2.02 (m, 2H), 1.50-1.70 (m, 3H).

Example 164 (General procedure 1)

Methyl-phenyl-carbamic acid 4-acetyl-phenyl ester

10

The title product was prepared from 4'-hydroxyacetophenone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (78%, colourless oil). HPLC-MS: m/z = 270.1 (M+1); R_t : 3.62 min.

^1H NMR (CDCl_3): 7.96 (d, 2H), 7.15-7.7.48 (m, 7 H), 3.43 (s, 3H), 2.58 (s, 3H).

15

Example 165 (General procedure 1)

Methyl-phenyl-carbamic acid pyridin-4-yl ester

20 The title product was prepared from 4-hydroxypyridine and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (11%, yellow solid). HPLC-MS: m/z = 229.2 (M+1); R_t : 1.66 min, purity: 67%.

Example 166 (General procedure 1)

Methyl-phenyl-carbamic acid pyridin-3-yl ester

25

The title product was prepared from 3-hydroxypyridine and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (94%, colourless oil).

HPLC-MS: m/z = 229.2 (M+1); R_t : 2.54 min.

30 ^1H NMR (CDCl_3): 10.70 (bs, 1H), 8.50 (d, 1H), 8.45-8.65 (m, 1H), 7.70-7.90 (m, 1H), 7.51 (dd, 1H), 7.42 (d, 2H), 7.34 (d, 2H), 7.28-7.37 (m, 1H), 3.43 (s, 3H).

Example 167 (General procedure 1)

Methyl-phenyl-carbamic acid 6-methyl-pyridin-3-yl ester

35 The title product was prepared from 3-hydroxy-6-methyl-pyridine and N-methyl-N-

phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (79%, colourless oil). HPLC-MS: m/z = 243.1 (M+1); R_t : 2.24 min.

^1H NMR (CDCl_3): 10.8 (bs, 1H), 8.54 (m, 1H), 7.76 (m, 1H), 7.26-7.50 (m, 6H), 3.42 (s, 3H), 2.69 (s, 3H).

5

Example 168 (General procedure 1)

Methyl-phenyl-carbamic acid isoquinolin-1-yl ester

10 The title product was prepared from and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (15%, colourless oil). HPLC-MS: m/z = 279.1 (M+1); R_t : 3.67 min.

Example 169 (General procedure 1)

Methyl-phenyl-carbamic acid 3-phenoxy-phenyl ester

15

The title product was prepared from 3-phenoxyphenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (79%, colourless oil).

HPLC-MS: m/z = 320.1 (M+1); R_t : 5.16 min.

20 ^1H NMR (CDCl_3): 7.20-7.50 (m, 8 H), 7.11 (t, 1H), 6.95-7.06 (m, 2H), 6.70-6.93 (m, 3H), 3.40 (s, 3H).

Example 170 (General procedure 1)

Methyl-phenyl-carbamic acid 3-acetyl-phenyl ester

25 The title product was prepared from m-hydroxyacetophenone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (62%, colourless oil). HPLC-MS: m/z = 270.1 (M+1); R_t : 3.56 min.

^1H NMR (CDCl_3): 7.78 (d, 1H), 7.68 (s, 1H), 7.22-7.50 (m, 7H), 3.43 (s, 3H), 2.58 (s, 3H).

30 **Example 171** (General procedure 1)

Methyl-phenyl-carbamic acid 4-acetyl-2-carbamoyl-phenyl ester

The title product was prepared from 5-acetylsalicylamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (87%, white solid). HPLC-
35 MS: m/z = 313.1 (M+1); R_t : 2.51 min.

¹H NMR (CDCl₃): 8.47 (bs, 1H), 8.08 (dd, 1H), 7.30-7.52 (m, 6H), 6.05 (bs, 1H), 5.38 (bs, 1H), 3.43 (s, 3H), 2.61 (s, 3H).

Example 172 (General procedure 1)

5 Methyl-phenyl-carbamic acid 4-acetyl-3-methyl-phenyl ester

The title product was prepared from 4'-hydroxy-2'-methylacetophenone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (76%, white crystals). HPLC-MS: *m/z* = 284.2 (M+1); R_t: 3.95 min.

10 ¹H NMR (CDCl₃): 7.72 (d, 1H), 7.22-7.47 (m, 5H), 6.90-7.12 (m, 2H), 3.43 (s, 3H), 2.55 (s, 3H), 2.52 (s, 3H).

Example 173 (General procedure 1)

Methyl-phenyl-carbamic acid 1-oxo-indan-4-yl ester

15

The title product was prepared from 4-hydroxyindanone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (77%, light yellow oil).

HPLC-MS: *m/z* = 282.1 (M+1); R_t: 3.54 min.

¹H NMR (CDCl₃): 7.61 (d, 1H), 7.27-7.50 (m, 7H), 3.45 (s, 3H), 3.00 (ds, 2H), 2.67 (t, 2H).

20

Example 174 (General procedure 1)

Methyl-phenyl-carbamic acid benzothiazol-2-yl ester

The title product was prepared from 2-benzothiazolol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (70%, white crystals). HPLC-MS: *m/z* = 307.1 (M+1); R_t: 4.24 min, purity 85%.

25

¹H NMR (CDCl₃): 7.78 (m, 2H), 7.28-7.50 (m, 7H), 3.40-3.70 (d, 3H).

Example 175 (General procedure 1)

30 Methyl-phenyl-carbamic acid 5-oxo-5,6,7,8-tetrahydro-naphthalen-2-yl ester

The title product was prepared from 6-hydroxy-1-tetralone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (49%, colorless oil). HPLC-MS: *m/z* = 296.2 (M+1); R_t: 3.90 min.

35 ¹H NMR (CDCl₃): 8.04 (d, 2H), 7.27-7.50 (m, 5H), 7.00-7.10 (m, 2H), 3.42 (s, 3H), 2.94 (t,

2H), 2.63 (t, 2H), 2.12 (qu, 2H).

Example 176 (General procedure 1)

Methyl-phenyl-carbamic acid benzo[d]isoxazol-3-yl ester

5

The title product was prepared from benzo[d]isoxazol-3-ol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (53%, colorless oil). HPLC-MS: m/z = 291.1 (M+23); R_t : 4.05 min.

^1H NMR (CDCl_3): 7.60-7.75 (m, 1H), 7.49-7.60 (m, 2H), 7.38-7.48 (m, 4H), 7.28-7.36 (m, 2H),
10 3.46 (s, 3H).

Example 177 (General procedure 1)

Methyl-phenyl-carbamic acid pyridin-2-yl ester

15 The title product was prepared from 2-hydroxypyridine and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (57%, colorless oil). HPLC-MS: m/z = 229.1 (M+23); R_t : 2.80 min.

^1H NMR (CDCl_3): 8.38 (d, 1H), 7.75 (t, 1H), 7.35-7.45 (m, 4H), 7.22-7.34 (m, 1H), 7.14-7.21 (t, 1H), 6.99-7.15 (bs, 1H), 3.44 (s, 3H).

20

Example 178 (General procedure 1)

Methyl-phenyl-carbamic acid 1-(methyl-phenyl-carbamoyl)-1H-benzimidazol-2-yl ester

The title product was prepared from 2-hydroxybenzimidazole and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (14%,
25 white crystals). HPLC-MS: m/z = 401.2 (M+1); R_t : 3.88 min, purity: 83%.

^1H NMR (CDCl_3): 7.32 (dd, 2H), 7.24-7.28 (m, 2H), 7.21-7.24 (m, 2H), 7.29 (t, 1H), 7.12-7.19 (m, 5H), 7.09-7.15 m, 2H), 3.28 (s, 6H).

30 **Example 179** (General procedure 1)

Methyl-phenyl-carbamic acid 4-[(pyridine-3-carbonyl)-amino]-phenyl ester

The title product was prepared from N-(4-Hydroxy-phenyl)-nicotinamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC. (1%, light
35 yellow crystals). HPLC-MS: m/z = 348.1 (M+1); R_t : 2.96 min.

Example 180 (General procedure 11)

4-Pyrrolidin-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

5 The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-(1-pyrrolidinyl)piperidine. The crude product was used without further purification (68%, white solid). HPLC-MS: m/z = 436.2 (M+1); R_t : 2.98 min.

^1H NMR (DMSO- d_6): 10.90 (bs, 1H), 8.57 (s, 1H), 8.24 (dd, 1H), 7.15-7.30 (m, 5H), 4.00-4.40 (m, 2H), 3.45-3.60 (m, 2H), 2.75-3.25 (m, 4H), 2.05-2.25 (d, 2H), 1.80-2.05 (m, 5H),
10 1.60-1.80 (m, 2H).

Example 181 (General procedure 12)

Methyl-o-tolyl-carbamic acid 4-(trifluoromethyl-pyridin-2-yloxy)-phenyl ester

15 The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and N-methyl-o-toluidine. The crude product was subjected to preparative HPLC (51%, colorless oil). HPLC-MS: m/z = 403.2 (M+1); R_t : 4.89 min.

^1H NMR (CDCl_3): 8.40 (s, 1H), 7.87 (dd, 1H), 7.05-7.18 (m, 4H), 6.96 (d, 1H), 3.30 (s, 3H),
2.36 (s, 3H).

20

Example 182 (General procedure 12)

Methyl-pyridin-2-yl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 2-(methylamino)pyridine. The crude product was subjected to preparative HPLC (55%, white
25 solid). HPLC-MS: m/z = 390.1 (M+1); R_t : 4.31 min.

Example 183 (General procedure 12)

Methyl-m-tolyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and N-methyl-m-toluidine. The crude product was subjected to preparative HPLC (51%, colorless oil). HPLC-MS: m/z = 403.2 (M+1); R_t : 4.98 min.

^1H NMR (CDCl_3): 8.42 (s, 1H), 7.88 (dd, 1H), 7.28 (d, 1H), 7.05-7.25 (m, 7H), 6.97 (d, 1H),
35 3.41 (s, 3H), 2.38 (s, 3H).

Example 184 (General procedure 12)

(3-Chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

5 The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 3-chloro-N-methylaniline. The crude product was subjected to preparative HPLC (54%, colorless oil). HPLC-MS: m/z = 423.1 (M+1); R_t : 5.07 min.

^1H NMR (CDCl_3): 8.42 (m, 1H), 7.88 (dd, 1H), 7.39 (m, 1H), 7.33 (t, 1H), 7.22-7.30 (m, 2H), 7.10-7.7.22 (m, 4H), 6.99 (d, 1H), 3.43 (s, 3H).

10

Example 185 (General procedure 12)

Methyl-p-tolyl-carbamic acid 4-(5-trifluoromethyl -pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and N-methyl-p-toluidine. The crude product was subjected to preparative HPLC (54%, white solid).
15 HPLC-MS: m/z = 403.2 (M+1); R_t : 4.99 min.

^1H NMR (CDCl_3): 8.42 (s, 1H), 7.87 (dd, 1H), 7.05-7.30 (m, 8H), 6.98 (d, 1H), 3.40 (s, 3H), 2.36 (s, 3H).

20 **Example 186** (General procedure 14)

Methyl-phenyl-carbamic acid 4-(3-pyridin-3-yl-acryloyl)-phenyl ester

The title product was prepared from 1-(4-hydroxy-phenyl)-3-pyridin-3-yl-prop-2-en-1-one and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative
25 HPLC (56%, off-white solid). HPLC-MS: m/z = 359.0 (M+1); R_t : 3.27 min.

^1H NMR (CDCl_3): 8.87 (d, 1H), 8.65 (dd, 1H), 8.03 (d, 2H), 7.97 (dt, 1H), 7.79 (d, 1H), 7.57 (d, 1H), 7.33-7.47 (m, 5H), 7.27-7.33 (m, 3H), 3.44 (s, 3H), 3.49 (t, 1H), 1.21 (t, 1H).

Example 187 (General procedure 14)30

Methyl-phenyl-carbamic acid 4-[3-(3,4,5-trimethoxy-phenyl)-acryloyl]-phenyl ester

The title product was prepared from 1-(4-hydroxyphenyl)-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (82%, yellow solid). HPLC-MS: m/z = 448.2 (M+1); R_t : 4.61 min.

35

^1H NMR (CDCl_3): 8.01 (d, 2H), 7.70 (d, 1H), 7.33-7.47 (m, 4H), 7.28-7.33 (m, 2H), 6.86 (s,

1H), 3.92 (s, 6H), 3.90 (s, 3H), 3.45 (s, 3H).

Example 188 (General procedure 14)

Methyl-phenyl-carbamic acid 4-diethylcarbamoyl-2-methoxy-phenyl ester

5

The title product was prepared from N,N-diethyl-4-hydroxy-3-methoxy-benzamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (58%, colorless oil). HPLC-MS: m/z = 357.1 (M+1); R_t : 3.64 min.

^1H NMR (CDCl_3): 7.34-7.44 (m, 4H), 7.21-7.28 (m, 1H), 7.03-7.09 (d, 1H), 6.99 (d, 1H), 6.90 (dd, 1H), 3.87 (s, 3H), 3.15-3.65 (bs, 4H), 3.43 (s, 3H), 1.05-1.35 (m, 6H).

Example 189 (General procedure 14)

Methyl-phenyl-carbamic acid 3-phenylcarbamoyl-phenyl ester

15 The title product was prepared from 3-hydroxy-N-phenyl-benzamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (54%, white solid). HPLC-MS: m/z = 347.2 (M+1); R_t : 4.01 min.

^1H NMR (CDCl_3): 7.76 (s, 1H), 7.68 (d, 1H), 7.58-7.65 (m, 3H), 7.27-7.51 (m, 8H), 7.15 (t, 1H), 3.44 (s, 3H).

20

Example 190 (General procedure 14)

Methyl-phenyl-carbamic acid quinolin-7-yl ester

25 The title product was prepared from 7-hydroxyquinoline and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (28%, white solid). HPLC-MS: m/z = 279.1 (M+1); R_t : 2.55 min.

^1H NMR (CDCl_3): 8.93 (dd, 1H), 8.18 (d, 1H), 7.83-7.86 (m, 1H), 7.82 (d, 1H), 7.35-7.50 (m, 6H), 7.27-7.32 (m, 1H), 3.47 (s, 1H).

30 **Example 191** (General procedure 14)

Methyl-phenyl-carbamic acid 4-(4-methyl-piperazine-1-carbonyl)-phenyl ester

The title product was prepared from 1-(4-hydroxybenzoyl)-4-methyl-piperazine and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (22%, colorless oil). HPLC-MS: m/z = 354.1 (M+1); R_t : 2.07 min.

35

^1H NMR (CDCl_3): 7.37-7.49 (m, 4H), 7.28-7.37 (m, 3H), 7.16-7.25 (m, 2H), 4.60-4.00 (bs, 2H), 3.43 (s, 3H), 3.35-3.90 (bs, 4H), 2.82 (s, 3H) 2.55-2.90 (bs, 2H).

Example 192 (General procedure 14)

5 Methyl-phenyl-carbamic acid 3-acetyl-amino-phenyl ester

The title product was prepared from 3-acetamidophenol and N-methyl-N-phenylcarbamo-yl chloride. The crude product was subjected to preparative HPLC (66%, white solid). HPLC-MS: m/z = 285.1 (M+1); R_t : 3.09 min.

10 ^1H NMR (CDCl_3): 7.40-7.49 (m, 1H), 7.31-7.40 (m, 5H), 7.25-7.30 (m, 1H), 7.22 (d, 1H), 7.14 (d, 1H), 6-84 (bd, 1H), 3.42 (s, 3H), 2.10 (s, 3H).

Example 193 (General procedure 14)

15 Methyl-phenyl-carbamic acid 4-benzoyl-phenyl ester

The title product was prepared from (4-hydroxy-phenyl)-phenyl-methanone and N-methyl-N-phenylcarbamo-yl chloride. The crude product was subjected to preparative HPLC (36%, colorless oil). HPLC-MS: m/z = 332.2 (M+1); R_t : 4.42 min.

20 ^1H NMR (CDCl_3): 7.75-7.85 (m, 4H), 7.58 (tt, 1H), 7.33-7.52 (m, 6H), 7.27-7.32 (m, 1H), 7.18-7.25 (m, 2H), 3.44 (s, 3H).

Example 194 (General procedure 14)

Methyl-phenyl-carbamic acid biphenyl-3-yl ester

25 The title product was prepared from 3-phenylphenol and N-methyl-N-phenylcarbamo-yl chloride. The crude product was subjected to preparative HPLC (52%, colorless oil). HPLC-MS: m/z = 304.2 (M+1); R_t : 4.75 min.

^1H NMR (CDCl_3): 7.52-7.67 (m, 2H), 7.22-7.52 (m, 11H), 7.03-7.22 (m, 1H), 3.45 (s, 3H).

30 **Example 195** (General procedure 14)

Methyl-phenyl-carbamic acid 1H-indol-4-yl ester

The title product was prepared from 4-hydroxyindole and N-methyl-N-phenylcarbamo-yl chloride. The crude product was subjected to preparative HPLC (48%, off-white solid). HPLC-

35 MS: m/z = 267.1 (M+1); R_t : 3.57 min.

¹H NMR (CDCl₃): 8.20 (ds, 1H), 7.36-7.50 (m, 4H), 7.17-7.35 (m, 2H), 7.08-7.17 (m, 2H), 6.94 (d, 1H), 6.44 (s, 1H), 3.48 (s, 3H).

Example 196 (General procedure 14)

5 **Methyl-phenyl-carbamic acid 5,6,7,8-tetrahydro-naphthalen-1-yl ester**

The title product was prepared from 5,6,7,8-tetrahydro-1-naphthol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (42%, colorless oil). HPLC-MS: *m/z* = 282.1 (M+1); R_t: 4.77 min.

10 ¹H NMR (CDCl₃): 7.33-7.45 (m, 4H), 7.22-7.32 (m, 1H), 7.02-7.13 (t, 1H), 6.82-7.96 (m, 2H), 3.42 (s, 3H), 2.70-2.82 (m, 2H), 2.50-2.65 (m, 2H), 1.62-1.83 (m, 4H).

Example 197 (General procedure 14)

Methyl-phenyl-carbamic acid 5-oxo-5,6,7,8-tetrahydro-naphthalen-1-yl ester

15

The title product was prepared from 5,6,7,8-tetrahydro-1-naphthol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (60%, colorless oil). HPLC-MS: *m/z* = 296.1 (M+1); R_t: 3.81 min.

¹H NMR (CDCl₃): 7.91 (dd, 1H), 7.21-7.50 (m, 7H), 3.44 (s, 3H), 2.60-2.93 (bs, 2H), 2.62 (t, 20 2H), 2.02-2.20 (m, 2H).

Example 198 (General procedure 14)

Methyl-phenyl-carbamic acid 1,3-dioxo-1,3-dihydro-isobenzofuran-4-yl ester

25 The title product was prepared from 4-hydroxy-isobenzofuran-1,3-dione and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (50%, white solid). HPLC-MS: *m/z* = 298.1 (M+1); R_t: 2.58 min, purity: 85%.

¹H NMR (CDCl₃): 7.90 (d, 1H), 7.48-7.62 (m, 2H), 7.28-7.45 (m, 5H), 3.38 (s, 3H).

30 **Example 199** (General procedure 1)

Methyl-phenyl-carbamic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-chloro-pyridin-2-yloxy)-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to column chromatography
35 (ethyl acetate/heptane (1:5) (85%, white solid). HPLC-MS: *m/z* = 355.1 (M+1); R_t: 4.56 min.

¹H NMR (CDCl₃): 8.10 (d, 1H), 7.62 (dd, 1H), 7.31-7.44 (m, 4H), 7.25-7.30 (m, 1H), 7.05-7.20 (m, 4H), 6.84 (d, 1H), 3.43 (s, 3H).

Example 200 (General procedure 14)

5 (3-Fluoro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenol and 3-fluoro-N-methylaniline. The crude product was subjected to preparative HPLC (31%, white solid). HPLC-MS: *m/z* = 407.0 (M+1); R_t: 4.93 min.

10 ¹H NMR (CDCl₃): 8.42 (s, 1H), 7.88 (dd, 1H), 7.36 (q, 1H), 7.07-7.24 (m, 6H), 7.00 (d, 1H), 7.92-7.05 (m, 1H), 3.44 (s, 3H).

Example 201 (General procedure 11)

4-Benzyl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester

15

The title product was prepared from 4-(5-chloro-pyridin-2-yloxy)-phenol and 1-benzylpiperazine. The title product precipitated from the reaction mixture and was collected by filtration (88%, off-white solid). HPLC-MS: *m/z* = 424.1 (M+1); R_t: 2.91 min.

¹H NMR (CDCl₃): 13.86 (bs, 1H), 8.11 (d, 1H), 7.60-7.70 (m, 3H), 7.45-7.55 (m, 3H), 7.05-7.20 (m, 4H), 6.98 (d, 1H), 4.26-4.40 (m, 2H), 4.15-4.25 (m, 2H), 3.38-3.53 (m, 2H), 3.60-4.15 (m, 2H), 2.72-2.92 (m, 2H).

20

Example 202 (General procedure 11)

4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester

25

The title product was prepared from 4-(5-chloro-pyridin-2-yloxy)-phenol and (3-pyridylmethyl)piperazine. The title product precipitated from the reaction mixture and was collected by filtration and recrystallized from ethanol (35%, off-white solid). HPLC-MS: *m/z* = 425.2 (M+1); R_t: 2.39 min.

30 ¹H NMR (CDCl₃): 13.00-14.50 (bs, 1H), 8.72 (d, 2H), 8.50 (bs, 1H), 8.11 (d, 1H), 7.64 (dd, 1H), 7.56 (bs, 1H), 7.06-7.14 (m, 4H), 6.88 (d, 1H), 3.6-4.4 (m, 6H), 2.70-3.50 (m, 4H).

Example 203 (General procedure 14)

4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester

35

The title product was prepared from 4-(5-chloro-pyridin-2-yloxy)-phenol and 4-hydroxymethyl-piperidine. The crude product was subjected to column chromatography (ethyl acetate/heptane, 1:1) (75%, colorless oil). HPLC-MS: m/z = 363.1 (M+1); R_t : 3.58 min. ^1H NMR (CDCl_3): 8.12 (d, 1H), 7.64 (dd, 1H), 7.05-7.24 (m, 4H), 6.85 (d, 1H), 4.33 (ds, 2H), 3.56 (t, 2H), 2.75-3.10 (m, 2H), 1.69-1.95 (m, 3H), 1.56 (s, 1H), 1.19-1.45 (m, 2H).

Example 204 (General procedure 1)

Methyl-phenyl-carbamic acid 4-morpholin-4-yl-phenyl ester

The title product was prepared from and 4-morpholin-4-yl-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was extracted with dichloromethane from citric acid (5%). The combined organic phases were evaporated and the product crystallized from ethanol (22%, crystals). HPLC-MS: m/z = 313.2 (M+1); R_t : 3.75 min. ^1H NMR ($\text{DMSO}-d_6$): 7.35-7.50 (m, 4H), 7.22-7.32 (m, 1H), 7.00 (d, 2H), 6.92 (d, 2H), 3.72 (t, 4H), 3.32 (s, 3H), 3.05 (t, 4H).

Example 205 (General procedure 1)

1,4-Dioxo-8-aza-spiro[4.5]decane-8-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin)-2-yloxy)-phenol and 4-piperidone ethylene ketal. The crude product was extracted with dichloromethane from citric acid (5%). The combined organic phases were evaporated and the product crystallized from ethanol (61%, crystals). HPLC-MS: m/z = 425.2 (M+1); R_t : 4.45 min.

Example 206 (General procedure 14)

The title product was prepared from 2-(4-hydroxy-phenyl)-indan-1,3-dione and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (1%, oil). HPLC-MS: m/z = 372.1 (M+1); R_t : 4.80 min.

Example 207 (General procedure 14)

Methyl-phenyl-carbamic acid 4-(5,6-dichloro-1,3-dioxo-1,3-dihydro-isoindol-2-yl)-phenyl ester

The title product was prepared from 5,6-dichloro-2-(4-hydroxy-phenyl)-isoindole-1,3-dione

and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (2%,). HPLC-MS: m/z = 441.1 (M+1); R_t : 5.00 min.

Example 208 (General procedure 14)

5 Methyl-phenyl-carbamic acid 4-(2-phenoxy-acetyl-amino)-phenyl ester

The title product was prepared from N-(4-hydroxy-phenyl)-2-phenoxy-acetamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was purified by preparative HPLC (54%, oil). HPLC-MS: m/z = 277.2 (M+1); R_t : 4.19 min.

10

Example 209 (General procedure 14)

Methyl-phenyl-carbamic acid 4-[2-(4-chloro-phenyl)-ethyl]-phenyl ester

The title product was prepared from 4-(4-chlorophenethyl)-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (57%, white crystals). HPLC-MS: m/z = 366.1 (M+1); R_t : 5.58 min.
 ^1H NMR (CDCl_3): 7.31-7.45 (m, 4H), 7.17-7.30 (m, 3H), 6.96-7.12 (m, 6H), 3.41 (s, 3H), 2.85 (s, 4H).

20 **Example 210** (General procedure 14)

Methyl-phenyl-carbamic acid 4-[(pyridine-2-carbonyl)-amino]-phenyl ester

The title product was prepared from pyridine-2-carboxylic acid (4-hydroxy-phenyl)-amide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (41%,). HPLC-MS: m/z = 348.1 (M+1); R_t : 4.00 min.
 ^1H NMR (CDCl_3): 7.32-7.47 (m, 4H), 7.24-7.31 (m, 3H), 7.10-7.23 (m, 4H), 7.79 (d, 2H), 3.41 (m, 6H).

Example 211 (General procedure 14)

30 Methyl-phenyl-carbamic acid 4-[methyl-(thiophene-2-carbonyl)-amino]-phenyl ester

The title product was prepared from thiophene-2-carboxylic acid (4-hydroxy-phenyl)-methyl-amide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (16%, oil). HPLC-MS: m/z = 367.2 (M+1); R_t : 3.97 min.

35

Example 212 (General procedure 14)

Methyl-phenyl-carbamic acid 4-butyrylamino-phenyl ester

5 The title product was prepared from 4'-hydroxybutyranilide and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (60%, white solid). HPLC-MS: m/z = 313.2 (M+1); R_t : 3.58 min.

Example 213 (General procedure 14)

Methyl-phenyl-carbamic acid 4-(4,6-dimethyl-pyrimidin-2-ylsulfanyl)-phenyl ester

10

The title product was prepared from 4-(4,6-dimethylpyrimidin-2-ylsulfanyl)-phenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (59%, white solid). HPLC-MS: m/z = 366.1 (M+1); R_t : 4.50 min.

^1H NMR (CDCl_3): 7.59 (d, 2H), 7.32-7.45 (m, 4H), 7.23-7.31 (m, 1H), 7.16 (d, 2H), 6.68 (s, 15 1H), 3.43 (s, 1H), 2.32.

Example 214 (General procedure 14)

Methyl-phenyl-carbamic acid 4-methanesulfonyl-phenyl ester

20 The title product was prepared from 4-methylsulfonylphenol and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (50%, white solid). HPLC-MS: m/z = 306.1 (M+1); R_t : 3.22 min.

Example 215 (General procedure 14)

25 Methyl-phenyl-carbamic acid 4-[2-(3-oxo-1,2,3,4-tetrahydro-quinoxalin-2-yl)-acetylamino]-phenyl ester

The title product was prepared from N-(4-hydroxyphenyl)-2-(3-oxo-1,2,3,4-tetrahydro-2-quinoxaliny)acetamide and N-methyl-N-phenylcarbamoyl chloride. The crude product was 30 subjected to preparative HPLC (20%, yellow solid). HPLC-MS: m/z = 431.2 (M+1); R_t : 3.55 min.

Example 216 (General procedure 14)

Methyl-phenyl-carbamic acid 4-phenylacetyl-phenyl ester

35

The title product was prepared from benzyl 4-hydroxyphenyl ketone and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (68%, light yellow oil). HPLC-MS: m/z = 431.2 (M+1); R_t : 3.55 min.

5 **Example 217** (General procedure 12)

4-Benzoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 4-benzoylpiperidine (29%, white solid). HPLC-MS: m/z = 471.3 (M+1); R_t : 5.12 min.

10

Example 218 (General procedure 11)

[1,4]Bipiperidiny-1'-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

15

The title product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenol and 4-piperidinopiperidine. The crude product was filtered from the reaction mixture and washed with diethyl ether to give the title product (50%, off-white solid). HPLC-MS: m/z = 450.1 (M+1); R_t : 3.12 min.

Example 219 (General procedure 12)

20

4-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

25

The title product was prepared from 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenol and 1-piperidin-4-yl-1,3-dihydro-benzimidazol-2-one (44%, oil). HPLC-MS: m/z = 499.1 (M+1); R_t : 4.35 min.

^1H NMR (CDCl_3): 10.15 (s, 1H), 8.44 (s, 1H), 7.90 (dd, 1H), 7.23 (t, 1H), 7.13-7.20 (m, 4H), 7.06-7.13 (m, 2H), 7.02 (d, 1H), 4.40-4.70 (m, 3H), 2.95-3.30 (m, 2H), 2.40-2.63 (m, 2H), 1.86 (d, 2H).

30

Example 220 (General procedure 12)

3-Diethylcarbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

35

The title product was prepared from 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenol and N,N-diethylnipecotamide (56%, oil). HPLC-MS: m/z = 466.1 (M+1); R_t : 4.51 min.

Example 221 (General procedure 12)

4-Carbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 5 The title product was prepared from 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenol and piperidine-4-carboxylic acid amide. The crude product was subjected to preparative HPLC (47%, white solid). HPLC-MS: m/z = 410.2 (M+1); R_t : 3.45 min.

Example 222 (General procedure 12)

- 10 3-Carbamoyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenol and nipeco-tamide. The crude product was purified by preparative HPLC (60%, white solid). HPLC-MS: m/z = 410.2 (M+1); R_t : 3.45 min.

15

Example 223 (General procedure 14)

Methyl-phenyl-carbamic acid 4-[[4-(methyl-phenyl-carbamoyloxy)-2-oxo-1,2-dihydro-quinoline-3-carbonyl]-amino]-phenyl ester

- 20 The title product was prepared from 4-hydroxy-2-oxo-1,2-dihydro-quinoline-3-carboxylic acid (4-hydroxyphenyl)-amide and N-methyl-N-phenylcarbonyl chloride. The crude product was subjected to preparative HPLC (6.6 %, oil). HPLC-MS: m/z = 563.2 (M+1); R_t : 4.46 min.
 ^1H NMR (CDCl_3): 8.38 (d, 1H), 7.75 (t, 1H), 7.35-7.45 (m, 4H), 7.22-7.34 (m, 1H), 7.14-7.21 (t, 1H), 6.99-7.15 (bs, 1H), 3.44 (s, 3H).

25

Example 224 (General procedure 14)

Methyl-phenyl-carbamic acid 4-[(4-hydroxy-2-oxo-1,2-dihydro-quinoline-3-carbonyl)-amino]-phenyl ester

- 30 The title product was prepared from 4-hydroxy-2-oxo-1,2-dihydro-quinoline-3-carboxylic acid (4-hydroxy-phenyl)-amide and N-methyl-N-phenylcarbonyl chloride (5%,). HPLC-MS: m/z = 430.1 (M+1); R_t : 4.80 min, purity: 70%.

Example 225 (General procedure 14)

- 35 Methyl-phenyl-carbamic acid 4-(4-hydroxy-benzyl)-phenyl ester

The title compound was prepared from 4,4'-dihydroxydiphenylmethane and N-methyl-N-phenylcarbamoyl chloride. The crude product was subjected to preparative HPLC (21 %, white crystals which turn red after standing). HPLC-MS: m/z = 324.2 (M+1); R_f : 4.21 min.

5 **Example 226** (General procedure 13)

Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-benzylcarbamoyl)-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and 4-trifluoromethyl-benzylamine (94%, white crystals). HPLC-MS: m/z = 429.2 (M+1); R_f : 4.35 min.

Example 227 (General procedure 13)

Methyl-phenyl-carbamic acid 4-(butyl-methyl-carbamoyl)-phenyl ester

15 The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and *n*-butyl-methyl-amine. The crude product was subjected to preparative HPLC (20%, oil). HPLC-MS: m/z = 341.2 (M+1); R_f : 3.96 min.

Example 228 (General procedure 13)

20 Methyl-phenyl-carbamic acid 4-(methyl-phenethyl-carbamoyl)-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and methyl-phenethyl-amine. The crude product was subjected to preparative HPLC (29 %, oil). HPLC-MS: m/z = 389.2 (M+1); R_f : 4.15 min.

25

Example 229 (General procedure 13)

Methyl-phenyl-carbamic acid 4-[(pyridin-2-ylmethyl)-carbamoyl]-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and 2-aminomethylpyridine. The crude product was used without further purification (57%, oil). HPLC-MS: m/z = 362.2 (M+1); R_f : 2.43 min.

30

^1H NMR (MeOH- d_4): 8.50 (d, 1H), 7.92 (d, 2H), 7.83 (dt, 1H), 7.38-7.50 (m, 5H), 7.26-7.37 (m, 2H), 7.15-7.26 (m, 2H), 4.69 (s, 2H), 3.41 (s, 3H).

35 **Example 230** (General procedure 13)

Methyl-phenyl-carbamic acid 4-(2-pyridin-2-yl-ethylcarbamoyl)-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and 2-aminoethylpyridine. The crude product was used without further purification (26%, oil). HPLC-MS: m/z = 376.2 (M+1); R_t : 2.25 min.

^1H NMR (MeOH- d_4): 8.51 (d, 1H), 7.87 (dt, 1H), 7.78 (d, 2H), 7.33-7.50 m, 6H), 7.25-7.33 (m, 1H), 7.18 (d, 2H), 3.74 (t, 2H), 3.41 (bs, 3H), 3.13 (t, 2H).

Example 231 (General procedure 13)**Methyl-phenyl-carbamic acid 4-(2-phenylamino-ethylcarbamoyl)-phenyl ester**

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and N-phenylethylenediamine. The crude product was used without further purification (80%, off-white foam). HPLC-MS: m/z = 390.2 (M+1); R_t : 3.51 min.

^1H NMR (MeOH- d_4): 7.82 (d, 2H), 7.36-7.48 (m, 4H), 7.22-7.34 (m, 1H), 7.15-7.22 (d, 2H), 7.10 (t, 2H), 6.69 (d, 2H), 6.62 (t, 1H), 3.72 (t, 2H), 3.57 (t, 2H), 3.40 (s, 3H), 3.20 (s, 1H).

Example 232 (General procedure 13)**Methyl-phenyl-carbamic acid 4-(3-methyl-butylcarbamoyl)-phenyl ester**

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and *iso*-amylamine. The crude product was used without further purification (95%, oil). HPLC-MS: m/z = 341.2 (M+1); R_t : 3.99 min.

^1H NMR (MeOH- d_4): 7.82 (d, 2H), 7.36-7.50 (m, 4H), 7.26-7.35 (m, 1H), 7.15-7.24 (d, 2H), 3.34-3.46 (m, 5H), 1.67 (sept, 1H), 1.50 (q, 2H), 0.96 (d, 6H).

Example 233 (General procedure 13)**Methyl-phenyl-carbamic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester**

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and 3,3-dimethylbutylamine. The crude product was used without further purification (88%, oil). HPLC-MS: m/z = 355.1 (M+1); R_t : 3.10 min.

^1H NMR (MeOH- d_4): 7.82 (d, 2H), 7.36-7.50 (m, 4H), 7.25-7.35 (m, 1H), 7.15-7.25 (d, 2H), 3.33-3.47 (m, 5H), 1.49-1.57 (m, 2H), 0.97 (s, 9H).

Example 234 (General procedure 13)

Methyl-phenyl-carbamic acid 4-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and C-(tetrahydro-furan-2-yl)methylamine. The crude product was used without further purification (86%, oil). HPLC-MS: $m/z = (M+1)$; R_t : min.

^1H NMR (MeOH- d_4): 7.84 (d, 2H), 7.35-7.50 (m, 4H), 7.25-7.35 (m, 1H), 7.12-7.25 (d, 2H), 4.10 (qui, 1H), 3.87 (q, 1H), 3.68-3.82 (m, 1H), 3.41 (s, 3H), 3.35-3.54 (m, 2H), 1.82-2.10 (m, 3H), 1.58-1.72 (m, 1H).

Example 235 (General procedure 13)

Methyl-phenyl-carbamic acid 4-cyclohexylcarbamoyl-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and cyclohexylamine. The crude product was used without further purification (79%, off-white crystals). HPLC-MS: $m/z = 353.2 (M+1)$; R_t : 3.98 min.

^1H NMR (MeOH- d_4): 8.18 (d, 1H), 7.85 (d, 2H), 7.36-7.52 (m, 4H), 7.25-7.33 (m, 1H), 7.20 (d, 2H), 3.74 (m, 1H), 1.77-1.88 (m, 2H), 1.65-1.77 (m, 2H), 1.60 (d, 1H), 1.24-1.40 (m, 4H), 1.07-1.23 (m, 1H).

Example 236 (General procedure 13)

Methyl-phenyl-carbamic acid 4-cyclopropylcarbamoyl-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and cyclopropylamine. The crude product was used without further purification (97%, oil). HPLC-MS: $m/z = 311.2 (M+1)$; R_t : 3.21 min.

^1H NMR (MeOH- d_4): 7.81 (d, 2H), 7.36-7.50 (m, 4H), 7.25-7.36 (m, 1H), 7.18 (d, 2H), 3.40 (bs, 3H), 2.78-2.87 (m, 1H), 0.75-0.85 (m, 2H), 0.57-0.65 (m, 2H).

Example 237 (General procedure 13)

Methyl-phenyl-carbamic acid 4-(cyclohexylmethyl-carbamoyl)-phenyl ester

The title product was prepared from 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester and C-cyclohexyl-methylamine. The crude product was used without further purification (84%, oil). HPLC-MS: $m/z = 367.3 (M+1)$; R_t : 4.28 min.

¹H NMR (MeOH-d₄): 7.81 (d, 2H), 7.36-7.49 (m, 4H), 7.25-7.32 (m, 1H), 7.19 (d, 2H), 3.40 (bs, 3H), 3.20 (d, 2H), 1.55-1.82 (m, 5H), 1.13-1.35 (m, 4H), 0.90-1.10 (m, 2H).

Example 238

5 Methyl-phenyl-carbamic acid 5-nitro-pyridin-2-yl ester

A solution of 2-hydroxy-5-nitropyridine (1.40 g, 10.0 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (3.43 g, 10.0 mmol) and triethylamine (0.42 ml, 10.0 mmol) in acetonitrile (25 ml) was heated at 50 °C for 5 h. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (25:75)) followed by crystallisation from ethyl acetate:heptane yielding the title compound (1.13 g, 41% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.44 (br.s, 3H), 7.17 (br.d, 1H), 7.27-7.45 (m, 5H), 8.49 (br.d, 1H), 9.19 (br.s, 1H); HPLC-MS (Method A): *m/z* = 296 (M+Na); R_t = 3.45 min.

Example 239

15 Methyl-phenyl-carbamic acid pyrimidin-2-yl ester

A solution of 2-hydroxypyrimidine hydrochloride (0.40 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.83 ml, 6.00 mmol) in acetonitrile (15 ml) was stirred at room temperature overnight. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate) followed by crystallisation from ethyl acetate:heptane yielding the title compound (0.08 g, 12% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.43 (br.s, 3H), 7.14-7.31 (m, 2H), 7.39 (m, 4H), 8.68 (d, 2H); HPLC-MS (Method A): *m/z* = 252 (M+Na); R_t = 2.32 min.

Example 240

30 Methyl-phenyl-carbamic acid 7-chloro-quinolin-4-yl ester

A solution of 7-chloro-4-hydroxyquinoline (0.54 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (40:60)) yielding the title compound (0.87 g, 93% yield) as a colourless

oil which solidified upon standing.

^1H NMR (300MHz, CDCl_3): δ 3.47 (br.s, 3H), 7.28-7.58 (m, 8H), 8.05 (br.s, 1H), 8.85 (d, 1H);
HPLC-MS (Method A): m/z = 313 (M+H); R_t = 3.79 min.

5 Example 241

Methyl-phenyl-carbamic acid quinolin-4-yl ester

A mixture of 4-hydroxyquinoline (0.44 g, 3.00 mmol) , 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00
10 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 , ethyl acetate:heptane (50:50)) yielding the title compound (0.75 g, 90% yield) as a white solid.

^1H NMR (300MHz, CDCl_3): δ 3.49 (br.s, 3H), 7.37 (br.t, 1H), 7.41-7.62 (m, 7H), 7.69 (br.t,
15 1H), 8.08 (br.d, 1H), 8.87 (d, 1H); HPLC-MS (Method A): m/z = 279 (M+H); R_t = 2.56 min.

Example 242

Methyl-phenyl-carbamic acid 5-methyl-isoxazol-3-yl ester

A mixture of 3-hydroxy-5-methylisoxazole (0.30 g, 3.00 mmol) , 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00
20 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 , ethyl acetate:heptane (25:75)) yielding the title compound (0.67 g, 96% yield) as a colourless oil.

^1H NMR (300 MHz, CDCl_3): δ 2.40 (s, 3H), 3.40 (br.s, 3H), 6.14 (br.s, 1H), 7.28-7.44 (m, 5H);
HPLC-MS (Method A): m/z = 255 (M+Na); R_t = 3.31 min.

Example 243

30 Methyl-phenyl-carbamic acid quinoxalin-2-yl ester

A mixture of 2-hydroxyquinoxaline (0.44 g, 3.00 mmol) , 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00
35 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours followed by heating at 40 °C for 24 hours. The solvent was evaporated *in vacuo* and the residue was purified

by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70)) yielding the title compound (0.65 g, 77% yield) as a white solid.

¹H NMR (300 MHz, CDCl₃): δ 3.48 (br.s, 3H), 7.29 (m, 1H), 7.41 (m, 4H), 7.72 (m, 2H), 8.00 (m, 1H), 8.10 (m, 1H), 8.67 (br.s, 1H); HPLC-MS (Method A): m/z = 280 (M+H); R_t = 3.66

5 min.

Example 244

Methyl-phenyl-carbamic acid 4-methyl-quinolin-2-yl ester

- 10 A mixture of 2-hydroxy-4-methylquinoline (0.48 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours followed by heating at 50 °C for 4 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70)) yielding the title compound (0.24 g, 27% yield) as a white solid.

¹H NMR (300 MHz, CDCl₃): δ 2.70 (s, 3H), 3.49 (br.s, 3H), 7.08 (m, 1H), 7.27 (m, 1H), 7.40 (m, 4H), 7.53 (t, 1H), 7.69 (t, 1H), 7.96 (d, 1H), 8.00 (d, 1H); HPLC-MS (Method A): m/z = 293 (M+H); R_t = 3.88 min.

20 **Example 245**

Methyl-phenyl-carbamic acid 3-methyl-quinoxalin-2-yl ester

- A mixture of 2-hydroxy-3-methylquinoxaline (0.48 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3H-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours followed by heating at 50 °C for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (0.59 g, 67% yield) as a white solid.

- ¹H NMR (300 MHz, CDCl₃): δ 2.61 (br.s, 3H), 3.48 (br.s, 3H), 7.30 (m, 1H), 7.42 (m, 4H), 7.69 (m, 2H), 7.99 (m, 2H); HPLC-MS (Method A): m/z = 294 (M+H); R_t = 3.92 min.

Example 246

Methyl-phenyl-carbamic acid 4,6-dimethyl-pyrimidin-2-yl ester

- 35 A solution of 4,6-dimethyl-2-hydroxypyrimidine (0.37 g, 3.00 mmol), 1-methyl-3-(methyl-

phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (0.46 g, 60% yield) as a white solid.

¹H NMR (300 MHz, CDCl₃): δ 2.48 (s, 6H), 3.43 (br.s, 3H), 6.92 (br.s, 1H), 7.22 (m, 1H), 7.37 (m, 4H); HPLC-MS (Method A): m/z = 258 (M+H); R_t = 2.77 min.

Example 247

10 Methyl-phenyl-carbamic acid isoquinolin-6-yl ester

A solution of 6-hydroxyquinoline (0.44 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (0.80 g, 96% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.47 (s, 3H), 7.28 (m, 1H), 7.33-7.54 (m, 6H), 7.59 (s, 1H), 8.08 (d, 2H), 8.86 (m, 1H); HPLC-MS (Method A): m/z = 279 (M+H); R_t = 2.63 min.

20 Example 248

Methyl-phenyl-carbamic acid quinolin-2-yl ester

A solution of 2-hydroxyquinoline (0.44 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. More acetonitrile (60 ml) was added and the solution was heated at 50 °C for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (40:60)) yielding the title compound (0.33 g, 40% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.50 (br.s, 3H), 7.14-7.30 (m, 2H), 7.42 (m, 4H), 7.52 (t, 1H), 7.71 (t, 1H), 7.82 (d, 1H), 8.00 (d, 1H), 8.19 (d, 1H); HPLC-MS (Method A): m/z = 279 (M+H); R_t = 3.91 min.

Example 249

Methyl-phenyl-carbamic acid isoquinolin-3-yl ester

A solution of 3-hydroxyisoquinoline (0.44 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (0.82 g, 99% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.49 (br.s, 3H), 7.24 (m, 1H), 7.33-7.47 (m, 5H), 7.51 (t, 1H), 7.63 (t, 1H), 7.77 (d, 1H), 7.94 (d, 1H), 9.06 (s, 1H); HPLC-MS (Method A): *m/z* = 279 (M+H); R_t = 3.68 min.

Example 250

Methyl-phenyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester

A solution of 4-(trifluoromethyl)-2-pyrimidol (0.49 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70)) yielding the title compound (0.35 g, 39% yield) as a colourless oil.

¹H NMR (300MHz, CDCl₃): δ 3.45 (br.s, 3H), 7.28 (m, 1H), 7.38 (m, 4H), 7.52 (br.s, 1H), 8.93 (br.s, 1H); HPLC-MS (Method A): *m/z* = 320 (M+Na); R_t = 3.58 min.

Example 251

Morpholine-4-carboxylic acid 4-trifluoromethyl-pyrimidin-2-yl ester

A solution of 4-(trifluoromethyl)-2-pyrimidol (0.49 g, 3.00 mmol), 4-morpholinecarbonyl chloride (0.45 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.51 g, 3.00 mmol) in *N,N*-dimethylformamide (15 ml) was stirred at room temperature for 2 hours. Water was added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70 → 50:50)) yielding the title compound (0.66 g, 80% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.57-3.81 (m, 8H), 7.57 (d, 1H), 8.97 (d, 1H); HPLC-MS (Method A): *m/z* = 300 (M+Na); R_t = 2.36 min.

Example 252

Methyl-phenyl-carbamic acid 3-nitro-pyridin-2-yl ester

A solution of 2-hydroxy-3-nitropyridine (0.42 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (40:60)) yielding the title compound (0.41 g, 50% yield) as a yellow solid.

¹H NMR (300MHz, CDCl₃): δ 3.41 + 3.58 (2 x br.s, 3H), 7.30 (m, 1H), 7.42 (m, 5H), 8.45 (br.d, 1H), 8.49 (br.s, 1H); HPLC-MS (Method A): *m/z* = 296 (M+Na); R_t = 3.22 min.

Example 253

Methyl-phenyl-carbamic acid 5-chloro-pyridin-2-yl ester

A solution of 5-chloro-2-pyridol (0.39 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70)) yielding the title compound (0.78 g, 99% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.44 (br.s, 3H), 7.00 (br.s, 1H), 7.27 (m, 1H), 7.39 (m, 4H), 7.69 (d, 1H), 8.30 (d, 1H); HPLC-MS (Method A): *m/z* = 285 (M+Na); R_t = 3.47 min.

Example 254

Methyl-phenyl-carbamic acid 5-(2-nitro-phenyl)-pyrimidin-2-yl ester

A solution of 5-(2-nitrophenyl)-pyrimidin-2-ol (0.35 g, 1.61 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (0.55 g, 1.61 mmol) and triethylamine (0.22 ml, 1.61 mmol) in acetonitrile (15 ml) was heated at 50 °C for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (0.18 g, 32% yield) as a yellow oil.

¹H NMR (300MHz, CDCl₃): δ 3.47 (br.s, 3H), 7.29 (m, 1H), 7.40 (m, 5H), 7.63 (dt, 1H), 7.72 (dt, 1H), 8.10 (d, 1H), 8.61 (br.s, 2H); HPLC-MS (Method A): *m/z* = 351 (M+H), 373 (M+Na), 723 (2M+Na); R_t = 3.65 min.

Example 255

Methyl-phenyl-carbamic acid 5-trifluoromethyl-pyridin-2-yl ester

- A solution of 2-hydroxy-5-(trifluoromethyl)pyridine (0.49 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (15:85)) yielding the title compound (0.59 g, 66% yield) as a white solid.
- ¹H NMR (300MHz, CDCl₃): δ 3.43 (br.s, 3H), 7.23 (br.s, 1H), 7.28 (m, 1H), 7.37 (m, 4H), 7.94 (br.d, 1H), 8.62 (br.s, 1H); HPLC-MS (Method A): *m/z* = 319 (M+Na); R_t = 3.85 min.

Example 256

Methyl-phenyl-carbamic acid 3-chloro-5-trifluoromethyl-pyridin-2-yl ester

- A solution of 3-chloro-5-(trifluoromethyl)-2-pyridinol (0.59 g, 3.00 mmol), 1-methyl-3-(methyl-phenyl-carbamoyl)-3*H*-imidazol-1-ium iodide (1.03 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in acetonitrile (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (15:85)) yielding the title compound (146 mg, 15% yield) as a white solid.
- ¹H NMR (300MHz, CDCl₃): δ 3.43 (br.s, 3H), 7.30 (m, 1H), 7.40 (d, 4H), 8.00 (br.s, 1H), 8.52 (br.s, 1H).; HPLC-MS (Method A): *m/z* = 353 (M+Na); R_t = 4.29 min.

Example 257

Methyl-phenyl-carbamic acid 5-nitro-3-trifluoromethyl-pyridin-2-yl ester

- A solution of 2-hydroxy-5-nitro-3-(trifluoromethyl)pyridine (0.36 g, 1.73 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (0.44 g, 2.59 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.29 g, 2.59 mmol) in tetrahydrofuran (15 ml) was stirred at room temperature for 18 hours. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (15:85)) yielding the title compound (0.55 g, 92% yield) as an orange solid.
- ¹H NMR (300MHz, CDCl₃): δ 3.46 (br.s, 3H), 7.23-7.46 (m, 5H), 8.70 (br.s, 1H), 9.37 (br.s, 1H); HPLC-MS (Method A): *m/z* = 364 (M+H); R_t = 4.08 min.

Example 258

(3-Chloro-phenyl)-methyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester

- 5 At 0 °C diphosgene (0.99 g, 5.00 mmol) was added to a stirred solution of 4-trifluoromethyl-2-hydroxypyrimidine (1.64 g, 10.0 mmol) in tetrahydrofuran (25 ml). The cooling bath was removed and stirring was continued at room temperature for 1 hour. (3-Chlorophenyl)-methylamine (0.35 g, 2.50 mmol) was added to one-fourth of the solution. After stirring overnight at room temperature the solvent was evaporated *in vacuo* and the residue was purified
10 by flash column chromatography (SiO₂, ethyl acetate:heptane (20:80)) followed by preparative HPLC, yielding the title compound (332 mg, 40%) as a colourless oil.
¹H NMR (300MHz, CDCl₃): δ 3.45 (br.s, 3H), 7.23-7.44 (m, 4H), 7.56 (d, 1H), 8.94 (d, 1H);
HPLC-MS (Method A): *m/z* = 354 (M+H); R_t = 4.03 min.

15 **Example 259**

Methyl-*m*-tolyl-carbamic acid 4-trifluoromethyl-pyrimidin-2-yl ester

- At 0 °C diphosgene (0.99 g, 5.00 mmol) was added to a stirred solution of 4-trifluoromethyl-2-hydroxypyrimidine (1.64 g, 10.0 mmol) in tetrahydrofuran (25 ml). The cooling bath was removed and stirring was continued at room temperature for 1 hour. Methyl-*m*-tolyl-amine (0.30
20 g, 2.50 mmol) was added to one-fourth of the solution. After stirring overnight at room temperature the solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (20:80)) followed by preparative HPLC, yielding the title compound (51 mg, 7%) as a colourless oil.
25 ¹H NMR (300MHz, CDCl₃): δ 2.37 (s, 3H), 3.42 (br.s, 3H), 7.07-7.31 (m, 4H), 7.52 (br.s, 1H), 8.92 (br.s, 1H); HPLC-MS (Method A): *m/z* = 334 (M+Na); R_t = 3.92 min.

Example 260

Morpholine-4-carboxylic acid 4-trifluoromethyl-pyrimidin-2-yl ester

- 30 A solution of 4-trifluoromethyl-2-hydroxypyrimidine (0.49 g, 3.00 mmol), 4-morpholinecarbonyl chloride (0.45 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (15 ml) was stirred at room temperature for 1 hour. Water and brine were added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The
35

residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (30:70 → 50:50)) yielding the title compound (0.66 g, 80% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.45 (br.s, 3H), 7.23-7.44 (m, 4H), 7.56 (d, 1H), 8.94 (d, 1H); HPLC-MS (Method A): *m/z* = 354 (M+H); R_t = 4.03 min.

5

Example 261

Methyl-phenyl-carbamic acid 4,5-dichloro-pyridazin-3-yl ester

A solution of 4,5-dichloro-3-hydroxypyridazine (0.49 g, 3.00 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and triethylamine (0.42 ml, 3.00 mmol) in tetrahydrofuran (15 ml) was stirred at room temperature for 3 days. The solvent was evaporated *in vacuo* and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (20:80)) yielding the title compound (0.12 g, 14% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 3.50 (s, 3H), 7.18-7.32 (m, 5H), 7.63 (s, 1H); HPLC-MS (Method A): *m/z* = 320 (M+Na); R_t = 2.91 min.

15

Example 262

Methyl-phenyl-carbamic acid 5-benzoylamino-pyridin-2-yl ester

A solution of *N*-(6-hydroxy-pyridin-3-yl)-benzamide (0.64 g, 3.00 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (15 ml) was stirred at room temperature for 1 hour. Water was added and the precipitates were collected by suction. The solids were dissolved in dichloromethane and the solution was dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was dissolved in ethyl acetate and filtered over a short pad of silica. Evaporation of the solvent *in vacuo* yielded the title compound (0.70 g, 68% yield) as a thick oil.

¹H NMR (300MHz, CDCl₃): δ 3.40 (br.s, 3H), 6.90 (br.s, 1H), 7.26 (m, 1H), 7.31-7.44 (m, 6H), 7.50 (m, 1H), 7.88 (d, 2H), 8.08 (dd, 1H), 8.37 (d, 1H), 8.79 (br.s, 1H); HPLC-MS (Method A): *m/z* = 348 (M+H); R_t = 3.49 min.

25

30

Example 263

Methyl-phenyl-carbamic acid 5-(cyclohexanecarbonyl-amino)-pyridin-2-yl ester

A solution of cyclohexanecarboxylic acid (6-hydroxy-pyridin-3-yl)-amide (0.66 g, 3.00 mmol),

35

N-methyl-*N*-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (20 ml) was stirred at room temperature for 18 hours. Water was added and the precipitates were collected by suction. The solids were dissolved in dichloromethane and the solution was dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was crystallised from ethyl acetate:heptane yielding the title compound (0.75 g, 71% yield) as a slightly coloured solid.

¹H NMR (300MHz, CDCl₃): δ 1.18-1.33 (m, 3H), 1.42-1.59 (m, 2H), 1.60 (m, 1H), 1.77-1.94 (m, 4H), 2.20 (m, 1H), 3.45 (br.s, 3H), 6.91 (br.s, 1H), 7.28 (m, 1H), 7.39 (m, 4H), 7.94 (br.s, 1H), 8.00 (dd, 1H), 8.20 (d, 1H); HPLC-MS (Method A): *m/z* = 354 (M+H); R_t = 3.74 min.

Example 264

Methyl-phenyl-carbamic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2*H*-[1,3']bipyridinyl-6'-yl ester

A solution of 6'-hydroxy-4,4-dimethyl-4,5-dihydro-3*H*-[1,3']bipyridinyl-2,6-dione (0.70 g, 3.00 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (20 ml) was stirred at room temperature for 18 hours. Water was added and the precipitates were collected by suction and subsequently dried in a vacuum oven yielding the title compound (0.75 g, 71% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ 1.20 (s, 6H), 2.68 (s, 4H), 3.44 (br.s, 3H), 7.14 (br.s, 1H), 7.25 (m, 1H), 7.37 (m, 4H), 7.48 (br.d, 1H), 8.08 (d, 1H); HPLC-MS (Method A): *m/z* = 368 (M+H); R_t = 3.41 min.

Example 265

Methyl-phenyl-carbamic acid 5-(2,2-dimethyl-propionylamino)-pyridin-2-yl ester

A solution of *N*-(6-hydroxy-pyridin-3-yl)-2,2-dimethyl-propionamide (0.58 g, 3.00 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (20 ml) was stirred at room temperature for 1 hour. Water was added and a thick oil was being formed. The water was decanted and the residue was dissolved in dichloromethane. The solution was dried over sodium sulphate, filtered and evaporated *in vacuo* yielding the title compound (0.55 g, 56% yield) as a brown oil that solidified upon standing.

¹H NMR (300MHz, CDCl₃): δ 1.29 (s, 9H), 3.43 (br.s, 3H), 6.97 (br.s, 1H), 7.26 (m, 1H), 7.38

(m, 4H), 7.64 (br.s, 1H), 8.10 (dd, 1H), 8.28 (br.s, 1H); HPLC-MS (Method A): m/z = 348 (M+H); R_t = 3.49 min.

Example 266

5 Methyl-phenyl-carbamic acid 5-(2-cyclohexyl-acetylamino)-pyridin-2-yl ester

A solution of 2-cyclohexyl-*N*-(6-hydroxy-pyridin-3-yl)-acetamide (0.70 g, 3.00 mmol), *N*-methyl-*N*-phenylcarbonyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.34 g, 3.00 mmol) in dimethylformamide (15 ml) was stirred at room temperature for 1 hour.
10 Water was added and the precipitates were collected by suction. The solids were dissolved in dichloromethane and the solution was dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was crystallised from ethyl acetate:heptane yielding the title compound (0.79 g, 72% yield) as a brown solid.

^1H NMR (300MHz, CDCl_3): δ 0.86-1.01 (m, 2H), 1.05-1.37 (m, 3H), 1.60-1.78 (m, 5H), 1.83
15 (m, 1H), 2.13 (d, 2H), 3.46 (br.s, 3H), 6.90 (br.s, 1H), 7.27 (m, 1H), 7.39 (m, 4H), 7.98 (d, 1H), 8.12 (s + br.s, 2H, CH + NH); HPLC-MS (Method A): m/z = 368 (M+H); R_t = 4.04 min.

Example 267

20 Methyl-phenyl-carbamic acid 5-(4-methoxy-phenoxy)-pyrimidin-2-yl ester

A solution of 5-(4-methoxy-phenoxy)-pyrimidin-2-ol (0.44 g, 2.00 mmol), *N*-methyl-*N*-phenylcarbonyl chloride (0.34 g, 2.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.22 g, 2.00 mmol) in dimethylformamide (15 ml) was stirred at room temperature for 1 hour. Water was added and the precipitates were collected by suction. The solids were dissolved in di-
25 chloromethane and the solution was dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was crystallised from ethyl acetate:heptane yielding the title compound (0.55 g, 79% yield) as an off-white solid.

^1H NMR (300MHz, CDCl_3): δ 3.43 (br.s, 3H), 3.82 (s, 3H), 6.91 + 7.00 (AB-system, 2 x 2H),
7.26 (m, 1H), 6.39 (m, 4H), 8.33 (s, 2H); HPLC-MS (Method A): m/z = 352 (M+H); R_t = 4.02
30 min.

Example 268

Methyl-phenyl-carbamic acid 5-(3,4-dichloro-phenoxy)-pyrimidin-2-yl ester

35 A solution of 5-(3,4-dichloro-phenoxy)-pyrimidin-2-ol (0.51 g, 2.00 mmol), *N*-methyl-*N*-

phenylcarbamoyl chloride (0.34 g, 2.00 mmol) and 1,4-diazabicyclo[2.2.2]octane (0.22 g, 2.00 mmol) in dimethylformamide (15 ml) was stirred at room temperature for 1 hour. Water was added and the precipitates were collected by suction. The solids were dissolved in dichloromethane and the solution was dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was crystallised from ethyl acetate:heptane yielding the title compound (0.51 g, 65% yield) as an off-white solid.

¹H NMR (300MHz, CDCl₃): δ 3.44 (br.s, 3H), 6.89 (dd, 1H), 7.14 (d, 1H), 7.27 (m, 1H), 7.39 (m, 4H), 7.44 (d, 1H), 8.42 (s, 2H); HPLC-MS (Method A): *m/z* = 390 (M+H); *R*_t = 4.66 min.

10 Example 269

Methyl-phenyl-carbamic acid 6-pyridin-2-ylmethyl-pyridazin-3-yl ester

A solution of 6-(2-pyridinylmethyl)-3-pyridazinol (100 mg, 0.53 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (91 mg, 0.53 mmol) and 1,4-diazabicyclo[2.2.2]octane (60 mg, 0.53 mmol) in dimethylformamide (10 ml) was stirred at room temperature for 2 hours. Water was added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was purified by flash column chromatography (SiO₂, ethyl acetate) yielding the title compound (70 mg, 41% yield) as a yellow oil.

¹H NMR (300MHz, CDCl₃): δ 3.42 (br.s, 3H), 4.50 (s, 2H), 7.11-7.33 (m, 4H), 7.39 (d, 4H), 7.60 (m, 2H), 8.52 (d, 1H); HPLC-MS (Method A): *m/z* = 321 (M+H); *R*_t = 1.98 min.

Example 270

Methyl-phenyl-carbamic acid 6-(4-methoxy-benzyl)-pyridazin-3-yl ester

25

A solution of 6-(2-pyridinylmethyl)-3-pyridazinol (97 mg, 0.45 mmol), *N*-methyl-*N*-phenylcarbamoyl chloride (76 mg, 0.45 mmol) and 1,4-diazabicyclo[2.2.2]octane (50 mg, 0.45 mmol) in dimethylformamide (10 ml) was stirred at room temperature for 2 hours. Water was added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (50:50)) yielding the title compound (117 mg, 41% yield) as a white solid.

30

¹H NMR (300MHz, CDCl₃): δ 3.43 (br.s, 3H), 3.79 (s, 3H), 4.28 (s, 2H), 6.83 (d, 2H), 7.17 (d, 2H), 7.27 (m, 3H), 7.40 (d, 4H); HPLC-MS (Method A): *m/z* = 350 (M+H); *R*_t = 3.60 min.

35

Example 271

Methyl-phenyl-carbamic acid 6-(2,4-dichloro-benzyl)-pyridazin-3-yl ester

A solution of 6-(2,4-dichlorobenzyl)-3-pyridazinol (98 mg, 0.38 mmol), *N*-methyl-*N*-phenylcarbonyl chloride (65 mg, 0.38 mmol) and 1,4-diazabicyclo[2.2.2]octane (43 mg, 0.38 mmol) in dimethylformamide (10 ml) was stirred at room temperature for 2 hours. Water was added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated *in vacuo*. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane (40:60)) yielding the title compound (119 mg, 80% yield) as a yellow oil.

¹H NMR (300MHz, CDCl₃): δ 3.43 (br.s, 3H), 4.42 (s, 2H), 7.17-7.44 (m, 10H); HPLC-MS (Method A): *m/z* = 388 (M+H); R_t = 4.44 min.

Example 272 (General procedure 15)

4-Pyridin-2-yl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl chloroformate and 1-pyridin-2-yl-piperazine. White crystals, yield 87 %; m.p. 247 - 248 °C; HPLC-MS: *m/z* = 445 (M+H); IR (KBr): ν 1713 (C=O) cm⁻¹.

Example 273 (General procedure 15)

4-(1,3-Benzodioxol-5-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(1,3-benzodioxol-5-yl)-piperazine. The crude product was partitioned between dichloromethane and 1 M aqueous sodium carbonate. The organic layer was washed with water, dried and evaporated. The residue was triturated with ethyl acetate - heptane (1:4) and the precipitate was collected by filtration and dried to give the title compound. Yield 39 %; m.p. 146 - 147 °C; ¹H NMR (DMSO-*d*₆): δ 8.60 - 8.56 (br, 1H), 8.29 - 8.20 (dd-like, 1H), 7.31 - 7.20 (m, 5H), 6.84 - 6.72 (d-like, 1H), 6.76 - 6.72 (d-like, 1H), 6.45 - 6.36 (dd-like, 1H), 3.81 - 3.47 (br m, 4H), 3.16 - 3.00 (br,m, 4H); HPLC-MS: *m/z* = 488 (M+H); IR (KBr): ν 1719 (C=O) cm⁻¹.

Example 274 (General procedure 15)

4-[2-(2-Hydroxyethoxy)ethyl]-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 2-(2-hydroxyethoxy)ethyl-piperazine. Yield 13 %; ¹H NMR (DMSO-*d*₆): δ 10.8 (br), 8.61 - 8.54 (br, 1H), 8.30 - 8.21 (dd-like, 1H), 7.32 - 7.19 (m, 5H), 4.3 - 3.9 (br, 2H), 3.9 - 3.0 (br m, alicyclics and aliphatics + water); HPLC-MS: m/z = 456(M+H); IR (KBr): ν 1724 (C=O) cm⁻¹.

10 **Example 275** (General procedure 15)

4-(Diphenylmethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(diphenylmethyl)piperazine, white crystals, yield 74 %; m.p. 168 - 169 °C;

¹H NMR (DMSO-*d*₆): δ 12.4 (br, 1H), 8.60 - 8.54 (d-like m, 1H), 8.28 - 8.20 (dd-like m, 1H), 7.98 - 7.82 (br, 2H), 7.55 - 7.15 (br m, 13 H), 5.6 (br, 1H), 4.35 - 3.48 (br, 3H), 3.35 - 3.0 (br, 5H); IR (KBr): ν 1723 (C=O) cm⁻¹.

20

Example 276 (General procedure 15)

4-(4-tert-Butylbenzyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

25 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-tert-butylbenzyl)diphenylmethyl)piperazine, white crystals, yield 56 %; m.p. 274 - 275 °C; HPLC-MS: m/z = 514 (M+H); IR (KBr): ν 1721 (C=O) cm⁻¹.

30 **Example 277** (General procedure 15)

4-(4-Fluorobenzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-fluorobenzyl)piperazine, white crystals, yield 69 %;

35

m.p. 240 - 243 °C; HPLC-MS: m/z = 476 (M+H), 498 (M+Na); IR (KBr): ν 1720 (C=O) cm^{-1} .

Example 278 (General procedure 15)

4-(2-Thienylethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-thienylethyl)piperazine, white crystals, yield 62 %; m.p. 236 - 237 °C; ^1H NMR ($\text{DMSO}-d_6$): δ 11.51 (br s, 1H), 8.61 - 8.54 (br m, 1H), 7.48 - 7.17 (m, 6H), 7.06 - 6.89 (m, 2H), 4.4 - 3.9 (br, 2H), 3.9 - 2.6 (br m, 16.5 H ~ 12H + water); HPLC-MS: m/z = 478 (M+H); IR (KBr): ν 1714 (C=O) cm^{-1} .

Example 279 (General procedure 15)

4-(1-Phenylethyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The crude hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(1-phenylethyl)piperazine. Trituration with water, filtering and drying of the residue gave white crystals, yield 31 %; m.p. 242 °C; ^1H NMR ($\text{DMSO}-d_6$): δ 11.56 (br s, 1H), 8.61 - 8.52 (br m, 1H), 8.30 - 8.19 (dd-like m, 1H), 7.77 - 7.31 (br m, 5H), 7.31 - 7.13 (m, 5H), 4.60 - 3.27 (br m, 6H + water), 3.27 - 2.57 (br, 3H), 1.73 (br d, 3H); IR (KBr): ν 1712 (C=O) cm^{-1} .

Example 280 (General procedure 15)

4-Octylpiperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The crude hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(1-phenylethyl)piperazine. Trituration with water, filtering and drying of the residue gave white crystals, yield 31 %; m.p. 244-245 °C; ^1H NMR ($\text{DMSO}-d_6$): δ 11.16 (br s, 1H), 8.61 - 8.57 (br m, 1H), 8.30 - 8.20 (dd-like m, 1H), 7.32 - 7.18 (m, 5H), 4.39 - 3.96 (br, 2H), 3.77 - 3.38 (br, 4H), 3.25 - 2.88 (br, 4H), 1.84 - 1.58 (br, 2H), 1.42 - 1.12 (br s, 10H), 0.87 (br t, 3H); IR (KBr): ν 1731, 1713 (C=O) cm^{-1} .

Example 281 (General procedure 15)

4-(3-Dimethylamino-propyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-

phenyl ester

The crude hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-dimethylaminopropyl)piperazine. A suspension of the product in ether was stirred with an excess of HCl in ether and the precipitate was washed with ether and dried to give the dihydrochloride of the title compound as white crystals, m.p. 292 - 293 °C; ¹H NMR (DMSO-*d*₆): δ 11.35 (br s, 1H), 10.46 (br s, 1H), 8.61 - 8.52 (br m, 1H), 8.31 - 8.17 (m, 1H), 7.35 - 7.16 (m, 5H), 4.45 - 4.00 (br, 2H), 3.80 - 3.45 (br, 4H), 3.30 - 3.01 (br, 6H), 2.78 (br s, 6H), 2.31 - 2.07 (br, 2H); IR (KBr): ν 1731, 1713 (C=O) cm⁻¹.

Example 282 (General procedure 15)

4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(pyrimidin-2-yl)piperazine. The crude product was dried in vacuo at 50 °C for 80 min and extracted with ether. The ether phase was evaporated and the residue was purified by flash chromatography on silica eluted with ethyl acetate - heptane 1:1 to give the title compound as white needles. Yield 14 %; m.p. 120-121 °C; ¹H NMR (DMSO-*d*₆): δ 8.61 - 8.56 (br, 1H), 8.41 (d, J = 4.8 Hz, 2H), 8.29 - 8.20 (dd-like, 1H), 7.31 - 7.20 (m, 5H), 6.69 (t-like m, J ~ 4.8 Hz, 1H), 3.94 - 3.78 (br s, 4H), 3.78 - 3.45 (br d, 4H); HPLC-MS: m/z = 446 (M+H); IR (KBr): ν 1719 (C=O) cm⁻¹.

Example 283 (General procedure 15)

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-cyclopropylpiperazine, yield 62 %. Recrystallisation from 0.2 M HCl gave white crystals, m.p. 238 - 239 °C; ¹H NMR (DMSO-*d*₆): δ 11.51 (br s, 1H), 8.61 - 8.55 (m, 1H), 8.30 - 8.20 (m, dd, 1H), 7.32 - 7.19 (m, d+s, 5H), 4.44 - 4.00 (br, 2H), 3.80 - 3.40 (br m, 4H), 3.29 - 2.93 (br m, 4H), 1.28 - 1.05 (br m, 1H), 0.75 - 0.58 (m, 2H), 0.50 - 0.35 (m, 2H). IR (KBr): ν 1730, 1713 (C=O) cm⁻¹.

Example 284 (General procedure 15)

4-Methyl-1,4-diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-methylhomopiperazine; white crystals, m.p. 210 - 211 °C; ¹H NMR (DMSO-*d*₆): δ 11.27 (br s, 1H), 8.61 - 8.54 (m, 1H), 8.30 - 8.20 (dd-like m, 1H),
5 7.34 - 7.18 (m, 5H), 4.13 - 3.08 (br, 11H, 8H + water), 2.80 (br s, 3H), 2.47 - 1.98 (br m, 2H); IR (KBr): ν 1723, 1711 (C=O) cm⁻¹.

Example 285 (General procedure 15)

4-Phenethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-phenethylpiperazine, yield 54 %. Recrystallisation from 99% EtOH gave white crystals, m.p. 245 - 247 °C; ¹H NMR (DMSO-*d*₆): δ 11.72 (br, 1H),
8.63 - 8.53 (br, 1H), 8.31 - 8.19 (dd-like m, 1H), 7.44 - 7.16 (m, 10H), 4.44 - 4.01 (br, 2H),
15 3.83 - 3.45 (br, 4H), 3.45 - 2.95 (br, ~8H, 6H + water); IR (KBr): ν 1713 (C=O) cm⁻¹.

Example 286 (General procedure 15)

4-Pyridin-2-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

20

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-pyridin-2-ylmethyl-piperazine. White crystals, yield 64 %; m.p. 189 - 190 °C; ¹H NMR (DMSO-*d*₆): δ 8.72 - 8.63 (m, 1H), 8.60 - 8.55 (br, 1H), 8.29 -
8.20 (dd-like, 1H), 8.02 - 7.90 (m, 1H), 7.80 - 7.65 (m, 1H), 7.56 - 7.45 (m, 1H), 7.31 - 7.22
25 (m, 5H), 4.52 (br s, 2H), 4.06 - 3.68 (br s, 4H), 3.68 - 2.93 (br, 4H + NH + water); HPLC-MS: m/z = 459 (M+H); IR (KBr): ν 1717 (C=O) cm⁻¹.

Example 287 (General procedure 15)

4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-pyridin-3-ylmethyl-piperazine. Trituration with water, filtering and drying of the residue gave white crystals. ¹H NMR (DMSO-*d*₆): δ 8.78 - 8.51 (m, 3H),
35 8.30 - 8.18 (dd-like, 1H), 8.12 - 8.00 (br d, 1H), 7.57 - 7.46 (m, 1H), 7.32 - 7.17 (m, 5H), 4.65

- 4.11 (br, 2H), 4.11 - 2.78 (br m, 6H + water); HPLC-MS: $m/z = 459$ (M+H); IR (KBr): ν 1723 (C=O) cm^{-1} .

Example 288 (General procedure 15)

5 4-(3-Phenylpropyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-phenylpropyl)piperazine, yield 68 %. m.p. 235 - 238
10 °C; ^1H NMR (DMSO- d_6): δ 11.51 (br, 1H), 8.61 - 8.55 (br, 1H), 8.29 - 8.20 (dd-like m, 1H), 7.38 - 7.16 (m, 10H), 4.38 - 3.96 (br, 2H), 3.83 - 3.40 (br, 4H), 3.30 - 2.91 (br, 4H), 2.75 - 2.57 (t-like m, 2H), 2.20 - 1.94 (m, 2H); IR (KBr): ν 1715 (C=O) cm^{-1} .

Example 289 (General procedure 15)

15 4-(4-Phenylbutyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-phenylbutyl)piperazine, yield 71 %. m.p. 232 - 234 °C;
20 ^1H NMR (DMSO- d_6): δ 11.32 (br s, 1H), 8.61 - 8.55 (br, 1H), 8.29 - 8.20 (dd-like m, 1H), 7.36 - 7.13 (m, 10H), 4.40 - 3.97 (br, 2H), 3.81 - 3.39 (br m, 4H), 3.26 - 2.91 (br, 4H), 2.71 - 2.55 (t-like m, 2H), 1.88 - 1.51 (br m, 4H); IR (KBr): ν 1728, 1713 (C=O) cm^{-1} .

25 **Example 290** (General procedure 15)

4-Benzyl-1,4-diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-benzylhomopiperazine, crude yield 70 %. 0.20 g Of the
30 crude product was heated with 3 ml of water, cooled at 0 °C, the precipitate filtered off and dried; m.p. 231 - 233 °C; ^1H NMR (DMSO- d_6): δ 11.38 (br s, 1H), 8.62 - 8.55 (br, 1H), 8.30 - 8.20 (dd-like m, 1H), 7.76 - 7.60 (br, 2H), 7.52 - 7.41 (br m, 3H), 7.32 - 7.19 (m, 5H), 4.38 (br s, 2H), 4.18 - 3.01 (br m, 8H + water), 2.6 - 2.0 (br, 2H + DMSO); IR (KBr): ν 1726, 1710 (C=O) cm^{-1} .

35

Example 291 (General procedure 15)

4-(3,4-Dichlorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

- 5 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3,4-dichlorophenyl)piperazine. The crude product was partitioned between dichloromethane and 2 M aqueous sodium carbonate. The organic layer was washed with water, dried and evaporated. The residue was triturated with ethyl acetate - heptane (1:4) and the precipitate was collected by filtration and dried to give the title
10 compound as white crystals. Yield 28 %; m.p. 115 - 116 °C; ¹H NMR (DMSO-*d*₆): δ 8.61 - 8.55 (br, 1H), 8.30 - 8.20 (dd-like, 1H), 7.48 - 7.40 (d-like, 1H), 7.31 - 7.16 (m, 6H), 7.04 - 6.94 (dd-like, 1H), 3.83 - 3.47 (br m, 4H), 3.36 - 3.24 (br s, 4H + water); HPLC-MS: m/z = 512 (M+H); IR (KBr): ν 1724, 1706 cm⁻¹.

15 **Example 292** (General procedure 15)

4-(4-Fluorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

- The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-fluorophenyl)piperazine. White crystals, yield 20 %;
20 m.p. 131-132 °C; ¹H NMR (DMSO-*d*₆): δ 8.61 - 8.53 (br, 1H), 8.30 - 8.19 (dd-like, 1H), 7.33 - 7.17 (m, 5H), 7.17 - 6.95 (m, 4H), 3.85 - 3.47 (br d, 4H), 3.25 - 3.07 (br m, 4H); HPLC-MS: m/z = 462 (M+H); IR (KBr): ν 1739, 1714 cm⁻¹.

25 **Example 293** (General procedure 15)

4-(2-Chlorophenyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

- The the hydrochloride of title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-chlorophenyl)piperazine. Drying *in vacuo* at 50 °C for
30 3½ h gave the title compound; ¹H NMR (DMSO-*d*₆): δ 8.61 - 8.55 (br, 1H), 8.29 - 8.20 (dd-like, 1H), 7.53 - 7.41 (m, 1H), 7.39 - 7.17 (m, 7H), 7.14 - 7.03 (m, 1H), 4.58 (br s, NH + water), 3.85 - 3.55 (br m, 4H), 3.12 - 2.99 (br m, 4H); HPLC-MS: m/z = 478 (M+H); IR (KBr): ν 1733 (C=O) cm⁻¹.

Example 294 (General procedure 15)

(2-Dimethylamino-ethyl)methylcarbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

5 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and N,N,N'-trimethylethylenediamine; m.p. 139 - 140 °C; ¹H NMR (MeOH-d₆): δ 8.45 - 8.38 (br, 1H), 8.15 - 8.05 (dd-like m, 1H), 7.31 - 7.11 (m, 5H), 3.84 - 3.73 (br m, 2H), 3.54 - 3.38 (br m, 3H), 3.19 (br s, 2H), 2.99 (br s, 6H); traces of impurities at 3.9 (br) and 3.1 (br); IR (KBr): ν 2695, 1705 cm⁻¹ (C=O) cm⁻¹.

10

Example 295 (General procedure 15)

4-Methylpiperazine-1-carboxylic acid 4-chlorophenyl ester

15 The hydrochloride of the title compound was prepared from 4-chlorophenyl chloroformate and 1-methylpiperazine, yield 81 %. White crystals, m.p. 237 - 240 °C; ¹H NMR (DMSO-d₆): δ 11.67 (br s, 1H), 7.50, 7.45, 7.24, 7.20 (AB-system, d = 7.47 and 7.22; J = 8.84 Hz, 4H), 4.40 - 3.91 (br, 2H), 3.77 - 2.92 (br m, 6H + water), 2.77 (s, 3H); IR (KBr): ν 1717 (C=O) cm⁻¹.

Example 296 (General procedure 15)

20 4-(4-Phenylbutyl)piperazine-1-carboxylic acid 4-chlorophenyl ester

The hydrochloride of the title compound was prepared from 4-chlorophenyl chloroformate and 1-(4-phenylbutyl)piperazine, yield 86 %. White crystals, m.p. 230 - 232 °C; ¹H NMR (DMSO-d₆): δ 11.43 (br, 1H), 7.51 - 7.43 (d-like m, 2H), 7.33 - 7.15 (m, 7H), 4.33 - 3.95 (br, max at 4.21 and 4.10 ppm; 2H), 3.72 - 3.36 (br m, 4H), 3.22 - 2.96 (br, max at 3.10 ppm, 4H), 2.62 (t, J = 7.54 Hz, 2H), 1.84 - 1.69 (m, 2H), 1.69 - 1.54 (m, 2H) ppm; IR (KBr): ν 1736, 1720 (C=O) cm⁻¹.

25

Example 297 (General procedure 15)

30 4-[2-(2-Hydroxyethoxy)ethyl]piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 2-(2-hydroxyethoxy)ethyl-piperazine. Pale powder, m.p. 176 - 178 °C; ¹H NMR (MeOH-d₄): δ Two AB-systems: 7.68 - 7.58 (d-like, 2H)

35

and 7.29 - 7.04 (m, 6H); 4.74 - 3.18 (complex, 16 H, partly overlapping with MeOH- d_4); HPLC-MS m/z = 455 (M+H), 477 (M+Na), R_t = 3.08 min.; IR (KBr): ν 1718 (C=O) cm^{-1} .

Example 298 (General procedure 15)

5 4-(1-Ethylpropyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-(1-ethylpropyl)piperazine, yield 74 %. White crystals, ^1H NMR (DMSO- d_6): δ 10.59 (br s, 1H), 2 AB-systems: 7.81 - 7.70 (d-like, 2H) and 7.31 - 7.09 (m, 6H); 4.38 - 3.99 (br s, 2H), 3.90 - 3.38 (br, 4H), 3.33 - 2.99 (br, 3H), 10 2.01 - 1.77 (m, 2H), 1.77 - 1.49 (m, 2H), 0.98 (t, 6H); HPLC-MS m/z = 437 (M+H).

Example 299 (General procedure 15)

15 4-Cycloheptylpiperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-cycloheptylpiperazine. The crude product was partitioned between dichloromethane and aqueous sodium carbonate. The organic layer was washed with water, dried and evaporated. The residue was triturated with ethyl acetate - heptane (1:4) and the precipitate was collected by filtration and dried to give the title compound. White crystals, ^1H NMR (MeOH- d_4): δ Two AB-systems: 7.68 - 7.58 (d-like, 2H) and 7.23 - 7.04 (m, 6H); 3.86 - 3.50 (br d, 4H), 2.98 - 2.66 (br, 5H), 2.04 - 1.37 (m, 12H); 20 HPLC-MS m/z = 463 (M+H); IR (KBr): ν 1730, 1707 cm^{-1} .

25 **Example 300** (General procedure 15)

4-Cyclohexylpiperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-cyclohexylpiperazine, yield 80 %. White crystals, m.p. 290 - 291 $^{\circ}\text{C}$, ^1H NMR (DMSO- d_6): δ 10.82 (br s, 1H), 2 AB-systems: 7.83 - 7.69 (d-like, 2H) and 7.32 - 7.10 (m, 6H); 4.40 - 4.02 (br, 2H), 3.75 - 3.39 (br, 4H), 3.31 - 2.98 (br, 3H), 2.23 - 2.02 (m, 2H), 1.92 - 1.74 (m, 2H), 1.70 - 0.97 (m, 6H); HPLC-MS m/z = 449 (M+H); IR (KBr): ν 1717 (C=O) cm^{-1} . 30

35 **Example 301** (General procedure 15)

4-(4-Chlorobenzyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-(4-chlorobenzyl)piperazine, yield 86 %.

- 5 White crystals, m.p. 232 - 234 °C; ¹H NMR (DMSO-*d*₆): δ 11.92 (br s, 1H), 3 AB-systems: 7.83 - 7.62 (t-like, 4H) and 7.62 - 7.48 (d-like, 2H), and 7.32 - 7.07 (m, 6H); 4.51 - 3.95 (br s at 4.37 ppm overlapping with br signal at 4.2 ppm, 4H), 3.95 - 2.95 (br m, 9H: 6H + water); IR (KBr): ν 1717 (C=O) cm⁻¹.

10 **Example 302** (General procedure 15)

4-(4-Methylbenzyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

- The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-(4-methylbenzyl)piperazine, yield 96 %.
- 15 White crystals, m.p. 250-252°C; HPLC-MS m/z = 472 (M+H); IR (KBr): ν 1720 (C=O) cm⁻¹.

Example 303 (General procedure 15)

- 4-(4-Methoxybenzyl)piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester
- 20 ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-methoxybenzyl)piperazine, yield 78 %. White crystals, m.p. 237 - 238 °C; HPLC-MS m/z = 488 (M+H); IR (KBr): ν 1719 (C=O) cm⁻¹.

25

Example 304 (General procedure 15)

4-(2-Chloro-6-fluoro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 30 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-chloro-6-fluorobenzyl)piperazine. White crystals, m.p. 204 - 205 °C (from ethanol); HPLC-MS m/z = 510 (M+H); IR (KBr): ν 1726 (C=O) cm⁻¹;

Example 305 (General procedure 15)

- 35 4-(3-Methoxyphenyl)piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethylphenoxy)phenyl chloroformate and 1-(3-methoxyphenyl)piperazine. White crystals, m.p. 168-171°C (sinters at 160 °C); HPLC-MS m/z = 473(M+1); IR (KBr): ν 1739, 1716 (C=O) cm^{-1} .

Example 306 (General procedure 15)

4-Benzyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)phenyl ester

The hydrochloride of the title compound was prepared from 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-benzyl-piperazine, yield 94 %. White crystals; m.p. 111 - 113 °C (resolidifies) and 114 - 115 °C; HPLC-MS m/z = 492 (M+H).

Example 307 (General procedure 8)Methyl-phenyl-carbamic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (98%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.44 (bs, 3H), 7.30-7.48 (m, 7H).; HPLC-MS : m/z = 343.9 (M+1); R_t = 4.12 min.

Example 308 (General procedure 8)Methyl-phenyl-carbamic acid benzotriazol-1-yl ester

The title compound was prepared from 1-hydroxybenzotriazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (98%, crystallizes slowly):

^1H NMR (300MHz; CDCl_3): δ 3.50 (bs, 3H), 7.37-7.57 (m, 8H), 8.04 (d, 1H).; HPLC-MS : m/z = 269.0 (M+1); R_t = 3.69 min.

Example 309 (General procedure 8)Methyl-phenyl-carbamic acid [1,2,3]triazolo[4,5-b]pyridin-3-yl ester

The title compound was prepared from [1,2,3]Triazolo[4,5-*b*]pyridin-3-ol and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified

by flash chromatography (Quad flash 12, EtOAc-heptane) (99%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.50 (bs, 3H), 7.30-7.60 (m, 6H), 8.40 (d, 1H), 8.75 (d, 1H);

HPLC-MS : m/z = 270.0 (M+1); R_t = 3.18 min.

5 **Example 310** (General procedure 8) Methyl-phenyl-carbamic acid 3-(2-nitro-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3-(2-nitrophenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified

10 by flash chromatography (Quad flash 12, EtOAc-heptane) (94%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.45 (bs, 3H), 6.44 (bs, 1H), 7.30-7.50 (m, 7H), 7.58 (dt, 1H), 7.72-7.78 (m, 2H); HPLC-MS : m/z = 339.1 (M+1); R_t = 4.15 min.

15 **Example 311** (General procedure 8) Methyl-phenyl-carbamic acid 3-(4-nitro-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3-(4-nitrophenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (99%, yellow crystals).

20 ^1H NMR (300MHz; CDCl_3): δ 3.47 (bs, 3H), 6.70 (bd, 1H), 7.32-7.50 (m, 6H), 7.94 (d, 2H), 8.26 (d, 2H); HPLC-MS : m/z = 339.1 (M+1); R_t = 4.41 min.

Example 312 (General procedure 8) Methyl-phenyl-carbamic acid 3-pyridin-2-yl-pyrazol-1-yl ester

25

The title compound was prepared from 1-hydroxy-3-(2-pyridyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (90%, oil).

30 ^1H NMR (300MHz; CDCl_3): δ 3.48 (bs, 3H), 6.95 (d, 1H), 7.20 (dd, 1H), 7.30-7.48 (m, 6H), 7.70 (dt, 1H), 7.93 (d, 1H), 8.61 (d, 1H); HPLC-MS : m/z = 295.1 (M+1); R_t = 2.75 min.

Example 313 (General procedure 8) Methyl-phenyl-carbamic acid 3-thiophen-2-yl-pyrazol-1-yl ester

35 The title compound was prepared from 1-hydroxy-3-(2-thienyl)pyrazole and N-methyl-N-

phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (66%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.46 (bs, 3H), 6.48 (bd, 1H), 7.03 (dd, 1H), 7.25 (dd, 1H), 7.30-7.48 (m, 7H); HPLC-MS : m/z = 300.1 (M+1); R_t = 4.16 min.

5

Example 314 (General procedure 8)Methyl-phenyl-carbamic acid 3-(2-fluoro-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3-(2-fluorophenyl)pyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (97%, oil).

10

^1H NMR (300MHz; CDCl_3): δ 3.48 (bs, 3H), 6.75 (bt, 1H), 7.07-7.47 (m, 9H), 7.97 (dt, 1H); HPLC-MS : m/z = 312.1 (M+1); R_t = 4.45 min.

15 **Example 315** (General procedure 8)Methyl-phenyl-carbamic acid 3-bromo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3-bromopyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (63%, oil).

20 ^1H NMR (300MHz; CDCl_3): δ 3.43 (bs, 3H), 6.31 (d, 1H), 7.26-7.48 (m, 6H).; HPLC-MS : m/z = 298.0 (M+1); R_t = 3.97 min.

Example 316 (General procedure 8)Methyl-phenyl-carbamic acid 5-iodo-pyrazol-1-yl ester

25 The title compound was prepared from 1-hydroxy-5-iodopyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (64%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.48 (bs, 3H), 6.42 (d, 1H), 7.28-7.47 (m, 6H).; HPLC-MS : m/z = 343.9 (M+1); R_t = 3.81 min.

30

Example 317 (General procedure 8)Methyl-phenyl-carbamic acid 2-chloro-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-chloroimidazole, hydrochloride and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product

35

was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (77%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.45 (bs, 3H), 6.90 (bs, 1H), 7.07 (bs, 1H), 7.35-7.40 (m, 3H), 7.46 (bt, 2H); HPLC-MS : m/z = 251.9 ($\text{M}+1$); R_t = 3.29 min.

5 **Example 318** (General procedure 8)Methyl-phenyl-carbamic acid 4-(4-methoxy-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-(4-methoxyphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product

10 was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (6%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.45 (bs, 3H), 3.82 (s, 3H), 6.90 (d, 2H), 7.30-7.48 (m, 7H), 7.54 (bs, 2H); HPLC-MS : m/z = 346.1 ($\text{M}+23$); R_t = 4.16 min.

15 **Example 319** (General procedure 8)Methyl-phenyl-carbamic acid 5-benzoyl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-5-benzoylpyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (30%, crystals).

20 ^1H NMR (300MHz; CDCl_3): δ 3.48 (bs, 3H), 6.69 (bs, 1H), 7.27-7.52 (m, 8H), 7.63 (t, 1H), 7.79 (d, 2H); HPLC-MS : m/z = 344.0 ($\text{M}+23$); R_t = 4.41 min.

Example 320 (General procedure 8)Methyl-phenyl-carbamic acid 5-(4-methoxy-phenyl)-pyrazol-1-yl ester

25

The title compound was prepared from 1-hydroxy-5-(4-methoxyphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (38%, yellow crystals).

30 ^1H NMR (300MHz; CDCl_3): δ 3.33 (bs, 3H), 3.86 (s, 3H), 6.34 (d, 1H), 6.95 (d, 2H), 7.25-7.45 (m, 8H); HPLC-MS : m/z = 324.1 ($\text{M}+1$); R_t = 4.27 min.

Example 321 (General procedure 8)Methyl-phenyl-carbamic acid 5-(4-dimethylamino-phenyl)-pyrazol-1-yl ester

35

The title compound was prepared from 1-hydroxy-5-(4-dimethylaminophenyl)-pyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (27%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.02 (s, 6H), 3.35 (bs, 3H), 6.30 (d, 1H), 6.72 (d, 2H), 7.30-7.46 (m, 8H); HPLC-MS : *m/z* = 337.1 (M+1); R_t = 4.11 min.

Example 322 (General procedure 8) Methyl-phenyl-carbamic acid 4,5-diiodo-pyrazol-1-yl ester

10 The title compound was prepared from 1-hydroxy-4,5-diiodopyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (76%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.45 (bs, 3H), 7.30-7.50 (m, 6H); HPLC-MS : *m/z* = 369.9 (M+1); R_t = 4.56 min.

15 **Example 323** (General procedure 8) Methyl-phenyl-carbamic acid 5-thiophen-2-yl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-5-(2-thienyl)pyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (51%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.44 (bs, 3H), 6.45 (bs, 1H), 7.09 (dd, 1H), 7.26 (bs, 1H), 7.28-7.49 (m, 7H); HPLC-MS : *m/z* = 300.1 (M+1); R_t = 4.18 min.

25 **Example 324** (General procedure 8) Methyl-phenyl-carbamic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-(4-methoxyphenyl)imidazole, hydrochloride and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (89%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.33 (bs, 3H), 3.86 (s, 3H), 6.92 (d, 2H), 7.04 (d, 1H), 7.10 (bs, 1H), 7.28 (d, 2H), 7.34-7.50 (m, 3H), 7.60 (bd, 2H); HPLC-MS : *m/z* = 324.1 (M+1); R_t = 2.87 min.

Example 325 (General procedure 8) Methyl-phenyl-carbamic acid 2-methylsulfanyl-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-methylsulfanyl-imidazole hydrochloride and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (97%, oil).
¹H NMR (300MHz; CDCl₃): δ 2.56 (s, 3H), 3.44 (bs, 3H), 7.00 (bs, 1H), 7.08 (bs, 1H), 7.32-7.49 (m, 5H); HPLC-MS : *m/z* = 264.1 (M+1); R_t = 2.99 min.

Example 326 (General procedure 8) Methyl-phenyl-carbamic acid 3,5-bis-(4-methoxy-phenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3,5-bis-(4-methoxyphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (29%, beige crystals).
¹H NMR (300MHz; CDCl₃): δ 3.35 (bs, 3H), 3.84 (s, 3H), 3.88 (s, 3H), 6.57 (s, 1H), 6.92 (d, 2H), 6.97 (d, 2H), 7.25-7.48 (m, 7H), 7.74 (d, 2H); HPLC-MS : *m/z* = 881.2 (2M+23); R_t = 5.26 min.

Example 327 (General procedure 8) Methyl-phenyl-carbamic acid 4-(4-fluoro-phenyl)-5-(4-methoxy-phenyl)-3-(4-methylphenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-(4-fluorophenyl)-5-(4-methoxyphenyl)-3-(4-methylphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (11%, crystals).
¹H NMR (300MHz; CDCl₃): δ 2.32 (s, 3H), 3.32 (bs, 3H), 3.83 (s, 3H), 6.87 (d, 2H), 6.93 (d, 2H), 7.12-7.48 (m, 13H); HPLC-MS : *m/z* = 530.2 (M+23); R_t = 6.04 min.

Example 328 (General procedure 8) Methyl-phenyl-carbamic acid 4-benzyl-5-(4-methoxy-phenyl)-3-(methylphenyl)-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-benzyl-5-(4-methoxyphenyl)-3-(4-methylphenyl)pyrazole and N-methyl-N-phenylcarbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-

heptane) (14%, oil).

¹H NMR (300MHz; CDCl₃): δ 2.31 (s, 3H), 3.31 (bs, 3H), 3.83 (s, 3H), 3.95 (s, 2H), 6.89 (d, 2H), 7.08-7.39 (m, 13H), 7.43 (d, 2H); HPLC-MS : *m/z* = 504.2 (M+1); R_t = 6.11 min.

5 **Example 329** (General procedure 8)Methyl-phenyl-carbamic acid 4-acetyl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-acetylpyrazole and N-methyl-N-phenylcarbamoil chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (89%, oil).

10 ¹H NMR (300MHz; CDCl₃): δ 2.44 (s, 3H), 3.44 (bs, 3H), 7.32-7.48 (m, 5H), 7.78 (bs, 1H), 7.85 (bs, 1H).

Example 330 (General procedure 8)Methyl-phenyl-carbamic acid 2-(4-nitro-phenyl)-imidazol-1-yl ester

15

The title compound was prepared from 1-hydroxy-2-(4-nitrophenyl)imidazole, hydrochloride and N-methyl-N-phenylcarbamoil chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (90%, yellow crystals).

20 ¹H NMR (300MHz; CDCl₃): δ 3.38 (bs, 3H), 7.16 (s, 1H), 7.20 (s, 1H), 7.31 (d, 2H), 7.38-7.61 (m, 3H), 7.86 (bs, 2H), 8.25 (d, 2H).

Example 331 (General procedure 8)Methyl-phenyl-carbamic acid 2-chloro-5-(4-methylphenyl)-imidazol-1-yl ester

25

The title compound was prepared from 1-hydroxy-2-chloro-5-(4-methylphenyl)imidazole, hydrochloride and N-methyl-N-phenylcarbamoil chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (80%, oil).

30 ¹H NMR (300MHz; CDCl₃): δ 2.40 (s, 3H), 3.33 (bs, 3H), 7.00 (s, 1H), 7.19-7.29 (m, 6H), 7.35-7.50 (m, 3H).

Example 332 (General procedure 8)Methyl-phenyl-carbamic acid 4-formyl-pyrazol-1-yl ester

35 The title compound was prepared from 1-hydroxy-4-formylpyrazole and N-methyl-N-

phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (73%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.45 (bs, 3H), 7.29-7.50 (m, 5H), 7.84 (bs, 1H), 7.90 (bs, 1H), 9.83 (s, 1H).

5

Example 333 (General procedure 8)Methyl-phenyl-carbamic acid 4-hydroxymethyl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-hydroxymethylpyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (62%, oil).

^1H NMR (300MHz; CDCl_3): δ 2.33 (bs, 1H), 3.40 (bs, 3H), 4.50 (s, 2H), 7.28-7.46 (m, 7H).

Example 334 (General procedure 8)Methyl-phenyl-carbamic acid 4-phenylethynyl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-phenylethynylpyrazole and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (40%, yellow crystals).

^1H NMR (300MHz; CDCl_3): δ 3.42 (bs, 3H), 7.30-7.58 (m, 12H).

Example 335 (General procedure 8)Methyl-phenyl-carbamic acid 2-bromo-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-bromoimidazole, hydrochloride and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (63%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.45 (bs, 3H), 6.98 (bd, 1H), 7.11 (bs, 1H), 7.33-7.50 (m, 5H); HPLC-MS : m/z = 296.0 (M+1); R_t = 2.90 min.

30

Example 336 (General procedure 8)Methyl-phenyl-carbamic acid 2-phenylsulfanyl-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2- phenylsulfanylimidazole hydrochloride and N-methyl-N-phenylcarbamoyl chloride applying the general procedure 8. The crude

35

product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (61%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.35 (s, 3H), 7.12 (bd, 1H), 7.17-7.39 (m, 11H); HPLC-MS : m/z = 326.0 (M+1); R_t = 3.65 min.

5 **Example 337** (General procedure 8)Morpholine-4-carboxylic acid imidazol-1-yl ester

The title compound was prepared from 1-hydroxyimidazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by preparative HPLC (water-acetonitrile-0.1% TFA) (36%, crystals).

10 ^1H NMR (300MHz; CDCl_3): δ 3.57 (bs, 2H), 3.66 (bs, 2H), 3.78 (t, 4H), 7.12 (bs, 1H), 7.15 (bt, 1H), 7.90 (s, 1H); HPLC-MS : m/z = 198.1 (M+1); R_t = 0.36 min.

Example 338 (General procedure 8)Morpholine-4-carboxylic acid 2-bromo-imidazol-1-yl ester

15

The title compound was prepared from 1-hydroxy-2-bromoimidazole, hydrochloride and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc) (98%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.60 (bs, 2H), 3.71 (bs, 2H), 3.80 (t, 4H), 7.02 (d, 1H), 7.18 (d, 20 1H); HPLC-MS : m/z = 276.0 (M+1); R_t = 1.73 min.

Example 339 (General procedure 8)Morpholine-4-carboxylic acid 2-chloro-imidazol-1-yl ester

25 The title compound was prepared from 1-hydroxy-2-chloroimidazole, hydrochloride and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc) (54%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.58 (bs, 2H), 3.68 (bs, 2H), 3.69 (t, 4H), 6.94 (d, 1H), 7.10 (d, 1H); HPLC-MS : m/z = 232.0 (M+1); R_t = 1.69 min.

30

Example 340 (General procedure 8)Morpholine-4-carboxylic acid 2-phenylsulfanyl-imidazol-1-yl ester

35 The title compound was prepared from 1-hydroxy-2-phenylsulfanylimidazole, hydrochloride and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product

was purified by flash chromatography (Quad flash 12, EtOAc) (98%, oil).

^1H NMR (300MHz; CDCl_3): δ 3.46 (bs, 4H), 3.66 (bs, 4H), 7.15 (d, 1H), 7.18-7.31 (m, 6H);

HPLC-MS : m/z = 306.1 (M+1); R_t = 2.75 min.

5 **Example 341** (General procedure 8)Morpholine-4-carboxylic acid 2-(4-methoxy-phenyl)-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-(4-methoxyphenyl)imidazole, hydrochloride and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product
10 was purified by flash chromatography (Quad flash 12, EtOAc) (49%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.51 (bs, 2H), 3.62-3.76 (m, 6H), 3.86 (s, 3H), 6.96 (d, 2H), 7.08 (d, 1H), 7.11 (d, 1H), 7.70 (d, 2H); HPLC-MS : m/z = 304.1 (M+1); R_t = 1.81 min.

15 **Example 342** (General procedure 8)Morpholine-4-carboxylic acid 4-bromo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-bromopyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (85%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.57 (bs, 2H), 3.68 (bs, 2H), 3.79 (t, 4H), 7.35 (d, 1H), 7.43 (d, 1H); HPLC-MS : m/z = 298.0 (M+23); R_t = 2.46 min.
20

Example 343 (General procedure 8)Morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash
25 chromatography (Quad flash 12, EtOAc-heptane) (99%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.56 (bs, 2H), 3.66 (bs, 2H), 3.77 (t, 4H), 7.41 (d, 1H), 7.44 (d, 1H); HPLC-MS : m/z = 324.0 (M+1); R_t = 2.65 min.

30 **Example 344** (General procedure 8)Morpholine-4-carboxylic acid 3,4,5-tribromo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-3,4,5-tribromopyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash
35 chromatography (Quad flash 12, EtOAc-heptane) (90%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.59 (bs, 2H), 3.69 (bs, 2H), 3.79 (t, 4H); HPLC-MS : m/z = 455.6 (M+23); R_t = 3.91 min.

Example 345 (General procedure 8)

5 **Morpholine-4-carboxylic acid 3-(4-methoxy-phenyl)-pyrazol-1-yl ester**

The title compound was prepared from 1-hydroxy-3-(4-methoxyphenyl)pyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (62%, crystals).

10 ¹H NMR (300MHz; CDCl₃): δ 3.58 (bs, 2H), 3.71 (bs, 2H), 3.79 (t, 4H), 3.84 (s, 3H), 6.53 (d, 1H), 6.92 (d, 2H), 7.40 (d, 1H), 7.71 (d, 2H); HPLC-MS : m/z = 326.0 (M+23); R_t = 3.21 min.

Example 346 (General procedure 8)Morpholine-4-carboxylic acid 3-thiophen-2-yl-pyrazol-1-yl ester

15

The title compound was prepared from 1-hydroxy-3-(2-thienyl)pyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (75%, crystals).

20 ¹H NMR (300MHz; CDCl₃): δ 3.57 (bs, 2H), 3.69 (bs, 2H), 3.79 (t, 4H), 6.51 (d, 1H), 7.04 (dd, 1H), 7.26 (dd, 1H), 7.34 (dd, 1H), 7.40 (d, 1H); HPLC-MS : m/z = 280.0 (M+1); R_t = 3.12 min.

Example 347 (General procedure 8)Morpholine-4-carboxylic acid pyrazol-1-yl ester

25 The title compound was prepared from 1-hydroxypyrazole and 4-morpholine carbonyl chloride applying the general procedure 8. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (94%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.57 (bs, 2H), 3.69 (bs, 2H), 3.79 (t, 4H), 6.32 (t, 1H), 7.38 (dd, 1H), 7.41 (dd, 1H); HPLC-MS : m/z = 198.0 (M+1); R_t = 1.18 min.

30 **Example 348** (General procedure 16)4-Methyl-piperazine-1-carboxylic acid pyrazol-1-yl ester

35 The title compound was prepared from 1-hydroxypyrazole and N-methylpiperazine applying the general procedure 16. The crude product was purified by preparative HPLC (water-acetonitrile-0.1% TFA) (17%, salt with TFA).

¹H NMR (300MHz; CDCl₃): δ 2.36 (s, 3H), 2.50 (bt, 4H), 3.59 (bs, 2H), 3.71 (bs, 2H), 6.31 (t, 1H), 7.38 (dd, 1H), 7.40 (dd, 1H); HPLC-MS : *m/z* = 211.0 (M+1); R_t = 0.40 min.

Example 349 (General procedure 16) 4-Cyclopentyl-piperazine-1-carboxylic acid pyrazol-1-yl ester

The title compound was prepared from 1-hydroxypyrazole and N-cyclopentylpiperazine applying the general procedure 16. The crude product was purified by preparative HPLC (water-acetonitrile-0.1% TFA) (34%, salt with TFA).

¹H NMR (300MHz; CDCl₃): δ 1.45-2.01 (m, 8H), 2.70 (bs, 4H), 2.92 (bs, 1H), 3.55 (bt, 1H), 3.63 (bs, 2H), 3.77 (bs, 2H), 6.31 (t, 1H), 7.38 (dd, 1H), 7.41 (dd, 1H); HPLC-MS : *m/z* = 265.1 (M+1); R_t = 0.54 min.

Example 350 (General procedure 16) 4-Phenyl-piperazine-1-carboxylic acid pyrazol-1-yl ester

The title compound was prepared from 1-hydroxypyrazole and N-phenylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (48%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.25 (bt, 4H), 3.72 (bs, 2H), 3.85 (bs, 2H), 6.33 (t, 1H), 6.92-6.98 (m, 3H), 7.27-7.33 (m, 2H), 7.38 (dd, 1H), 7.41 (dd, 1H); HPLC-MS : *m/z* = 273.1 (M+1); R_t = 3.06 min.

Example 351 (General procedure 16) 4-Pyridin-2-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester

The title compound was prepared from 1-hydroxypyrazole and 1-(2-pyridyl)piperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (59%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.62-3.85 (m, 8H), 6.32 (t, 1H), 6.66-6.72 (m, 2H), 7.39 (dd, 1H), 7.42 (dd, 1H), 7.53 (dt, 1H), 8.21 (d, 1H); HPLC-MS : *m/z* = 274.1 (M+1); R_t = 0.63 min.

Example 352 (General procedure 16) 4-Pyrimidin-2-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester

The title compound was prepared from 1-hydroxypyrazole and 1-(2-pyrimidyl)piperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (57%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.65 (bs, 2H), 3.78 (bs, 2H), 3.95 (bs, 4H), 6.32 (t, 1H), 6.58 (t, 2H), 7.38 (dd, 1H), 7.42 (dd, 1H), 8.35 (d, 1H); HPLC-MS : *m/z* = 275.2 (M+1); R_t = 2.14 min.

Example 353 (General procedure 16)4-Benzo[1,3]dioxol-5-yl-piperazine-1-carboxylic acid pyrazol-1-yl ester

10 The title compound was prepared from 1-hydroxypyrazole and 1-Benzo[1,3]dioxol-5-ylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (92%, crystals).

¹H NMR (300MHz; CDCl₃): δ 3.11 (bt, 4H), 3.71 (bs, 2H), 3.83 (bs, 2H), 5.93 (s, 2H), 6.32 (t, 1H), 6.39 (dd, 1H), 6.57 (d, 1H), 6.74 (d, 1H), 7.37 (dd, 1H), 7.41 (dd, 1H); HPLC-MS : *m/z* = 15 317.2 (M+1); R_t = 2.96 min.

Example 354 (General procedure 16)4-Benzyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

20 The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-benzylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (32%, oil).

¹H NMR (300MHz; CDCl₃): δ 2.53 (bs, 4H), 3.57 (bs, 4H), 3.67 (bs, 2H), 7.28-7.38 (m, 5H), 7.40 (d, 1H), 7.43 (d, 1H); HPLC-MS : *m/z* = 413.0 (M+1); R_t = 1.75 min.

25

Example 355 (General procedure 16)4-Cyclopentyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and N-cyclopentylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane-3% Et₃N) (61%, crystals).

30 ¹H NMR (300MHz; CDCl₃): δ 1.33-1.94 (m, 8H), 2.46-2.61 (m, 5H), 3.57 (bt, 2H), 3.67 (bt, 2H), 3.63 (bs, 2H), 3.77 (bs, 2H), 7.40 (d, 1H), 7.44 (d, 1H); HPLC-MS : *m/z* = 391.1 (M+1); R_t = 1.31 min.

35

Example 356 (General procedure 16) 4-(4-Fluoro-benzyl)-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-(4-fluorobenzyl)piperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (48%, crystals).

^1H NMR (300MHz; CDCl_3): δ 2.50 (t, 4H), 3.51 (s, 2H), 3.57 (bt, 2H), 3.66 (bt, 2H), 7.02 (t, 2H), 7.29 (dd, 2H), 7.40 (d, 1H), 7.43 (d, 1H); HPLC-MS : m/z = 431.0 (M+1); R_t = 1.79 min.

Example 357 (General procedure 16) 4-Phenyl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and N-phenylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (52%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.25 (t, 4H), 3.72 (bs, 2H), 3.81 (bs, 2H), 6.90-6.97 (m, 3H), 7.30 (t, 2H), 7.41 (d, 1H), 7.47 (d, 1H); HPLC-MS : m/z = 399.1 (M+1); R_t = 3.91 min.

Example 358 (General procedure 16) 4-Pyridin-2-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-(2-pyridyl)piperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (45%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.62-3.82 (m, 8H), 6.66-6.72 (m, 2H), 7.42 (d, 1H), 7.47 (d, 1H), 7.54 (dt, 1H), 8.22 (d, 1H); HPLC-MS : m/z = 400.0 (M+1); R_t = 1.47 min.

Example 359 (General procedure 16) 4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-(2-pyrimidyl)piperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (10%, crystals).

^1H NMR (300MHz; CDCl_3): δ 3.62 (bs, 2H), 3.72 (bs, 2H), 3.95 (bs, 4H), 6.58 (t, 2H), 7.41 (d, 1H), 7.46 (d, 1H), 8.36 (d, 1H); HPLC-MS : m/z = 401.0 (M+1); R_t = 3.09 min.

Example 360 (General procedure 16) 4-Benzo[1,3]dioxol-5-yl-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

- 5 The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-Benzo[1,3]dioxol-5-ylpiperazine applying the general procedure 16. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (59%, crystals).
1H NMR (300MHz; CDCl₃): δ 3.11 (t, 4H), 3.69 (bs, 2H), 3.80 (bs, 2H), 5.92 (s, 2H), 6.38 (dd, 1H), 6.56 (d, 1H), 6.73 (d, 1H), 7.41 (d, 1H), 7.45 (d, 1H); HPLC-MS : m/z = 443.0 (M+1); R_t
10 = 3.79 min.

Example 361

4-(1-Ethylpropyl)piperazine-1-carboxylic acid 3-trifluoromethylphenyl ester hydrochloride salt

- 15 To a stirred mixture of 1-(1-ethylpropyl)piperazine (175 μ l, 1.0 mmol) and dry DCM (10 ml) was added 3-trifluoromethylphenyl chloroformate (250 mg, 1.1 mmol). The mixture was stirred overnight at room temperature and then diluted with DCM (50 ml). The reaction mixture was washed with 1 N NaOH (3 x 25 ml) and water (2 x 25 ml). The organic solution was concentrated and the residue was dissolved in a 0.5 N HCl solution (15 ml) and a small portion of acetonitrile. The
20 acidic solution was concentrated and stirred with ethyl acetate (15 ml). The solid was isolated and dried to give 330 mg (86 %) of the title compound as a solid.

M.p. 260-261°C.

- 1H NMR (400 MHz, DMSO-d₆): δ 0.98 (t, 6H), 1.63 (hept, 2H), 1.86-1.98 (m, 2H), 3.03-3.12 (m, 1H), 3.12-3.31 (m, 2H), 3.41-3.49 (m, 2H), 3.52-3.85 (m, 2H), 4.05-4.35 (m, 2H), 7.47-7.70 (m,
25 4H), 11.0 (brs, 1H).

Example 362

4-(1-Ethylpropyl)piperazine-1-carboxylic acid naphthalen-1-yl ester hydrochloride salt

- 30 To a stirred mixture of 1-(1-ethylpropyl)piperazine (175 μ l, 1.0 mmol) and dry DCM (10 ml) was added 1-naphthalenyl chloroformate (225 mg, 1.1 mmol). The mixture was stirred overnight at room temperature and then diluted with DCM (50 ml). The reaction mixture was washed with 1 N NaOH (3 x 25 ml) and water (2 x 25 ml). The organic solution was concentrated and the residue was dissolved in a 0.5 N HCl solution (15 ml) and a small portion of acetonitrile. The acidic
35 solution was concentrated and stirred with ethyl acetate (15 ml). The solid was isolated and

dried to give 310 mg (85 %) of the title compound as a solid.

M.p. 288-290°C.

¹H NMR (400 MHz, DMSO-d₆): δ 1.00 (t, 6H), 1.63 (hept, 2H), 1.86-2.02 (m, 2H), 3.07-3.18 (m, 1H), 3.18-3.42 (m, 2H), 3.42-3.55 (m, 2H), 3.55-3.73 (m, 1H), 3.78-3.95 (m, 1H), 4.05-4.25 (m, 1H), 4.35-4.55 (m, 1H), 7.35 (d, 1H), 7.53 (t, 1H), 7.56-7.7.61 (m, 2H), 7.85 (d, 1H), 7.90-8.05 (m, 2H), 10.75 (brs, 1H).

Example 363

4-(1-Ethylpropyl)piperazine-1-carboxylic acid 4-fluorophenyl ester hydrochloride salt

To a stirred mixture of 1-(1-ethylpropyl)piperazine (350 µl, 2.0 mmol) and dry DCM (15 ml) was added 4-fluorophenyl chloroformate (350 mg, 2.0 mmol). The mixture was stirred overnight at room temperature and then diluted with DCM (50 ml). The reaction mixture was washed with 1 N NaOH (3 x 25 ml) and water (2 x 25 ml). The organic solution was concentrated and the residue was re-evaporated twice with acetonitrile to give 590 mg of the free base. The hydrochloride salt was prepared from 465 mg free base by addition of a 0.5 N HCl solution (15 ml) and a small portion of acetonitrile. The acidic solution was concentrated and stirred with ethyl acetate (15 ml). The solid was isolated and dried to give 470 mg (90 %) of the title compound as a solid.

M.p. 275-277°C.

¹H NMR (400 MHz, DMSO-d₆): δ 0.98 (t, 6H), 1.64 (hept, 2H), 1.85-1.95 (m, 2H), 3.02-3.11 (m, 1H), 3.11-3.28 (m, 2H), 3.38-3.46 (m, 2H), 3.50-3.80 (m, 2H), 4.00-4.30 (m, 2H), 7.18-7.26 (m, 4H), 10.85 (brs, 1H).

Example 364

4-(1-Ethylpropyl)piperazine-1-carboxylic acid 2-nitrophenyl ester hydrochloride salt

To a stirred mixture of 1-(1-ethylpropyl)piperazine (175 µl, 1.0 mmol) and dry DCM (10 ml) was added 2-nitrophenyl chloroformate (201 mg, 1.0 mmol). The mixture was stirred overnight at room temperature and then diluted with DCM (50 ml). The reaction mixture was washed with 1 N NaOH (3 x 25 ml) and water (2 x 25 ml). The organic solution was concentrated and the residue was dissolved in a 0.5 N HCl solution (15 ml). The acidic solution was concentrated and

stirred with ethyl acetate (15 ml). The solid was isolated and dried to give 310 mg (86 %) of the title compound as a solid.

M.p. 251-253°C.

5 ¹H NMR (400 MHz, DMSO-d₆): δ 0.96 (t, 6H), 1.65 (hept, 2H), 1.83-1.95 (m, 2H), 3.06-3.25 (m, 3H), 3.42-3.53 (m, 2H), 3.53-3.83 (m, 2H), 4.02-4.13 (m, 1H), 4.20-4.34 (m, 1H), 7.50-7.55 (m, 2H), 7.83 (t, 1H), 8.13 (d, 1H), 10.9 (brs, 1H).

10 **Example 365** 4-(1-Ethylpropyl)piperazine-1-carboxylic acid 4-methoxycarbonylphenyl ester hydrochloride salt

To a stirred mixture of 1-(1-ethylpropyl)piperazine (350 μl, 2.0 mmol) and dry DCM (15 ml) was
15 added 4-methoxycarbonylphenyl chloroformate (430 mg, 2.0 mmol). The mixture was stirred overnight at room temperature and then diluted with DCM (50 ml). The reaction mixture was washed with 1 N NaOH (3 x 25 ml) and water (2 x 25 ml). The organic solution was concentrated and re-evaporated twice with acetonitrile. The residue was dissolved in a 0.5 N HCl solution (15 ml) and a small portion of acetonitrile. The acidic solution was concentrated and stirred
20 with ethyl acetate (15 ml). The solid was isolated and dried to give 670 mg (90 %) of the title compound as a solid.

M.p. 248°C decomp.

25 ¹H NMR (400 MHz, DMSO-d₆): δ 0.97 (t, 6H), 1.63 (hept, 2H), 1.85-1.95 (m, 2H), 3.02-3.11 (m, 1H), 3.11-3.30 (m, 2H), 3.40-3.48 (m, 2H), 3.50-3.80 (m, 2H), 3.85 (s, 3H), 4.05-4.33 (m, 2H), 7.33 (d, 2H), 8.00 (d, 2H), 10.7 (brs, 1H).

Example 366

Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-butyrylamino)-pyridin-2-yl ester

30

A solution of N-(6-hydroxy-pyridin-3-yl)-3,3-dimethyl-butyramide (0.42 g, 2.00 mmol), N-methyl-N-phenylcarbonyl chloride (0.37 g, 2.20 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.25 g, 2.20 mmol) in tetrahydrofuran (20 mL) was stirred at room temperature for 2.5 hours. Water was added and the solution was extracted three times with dichloromethane. The
35 combined organic layers were dried over sodium sulphate, filtered and evaporated in vacuo.

The residue was redissolved in ethyl acetate and the solution was filtered over a short pad of silicagel. Evaporation of the solvent and crystallisation of the solid from ethyl acetate:heptane yielded the title compound (0.54 g, 79% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 1.03 (s, 9H), 2.15 (s, 2H), 3.47 (br.s, 3H), 6.91 (br.s, 1H), 7.27 (m, 1H), 7.38 (m, 4H), 7.99 (br.s + dd, 2H), 8.13 (d, 1H); HPLC-MS (Method A): m/z = 342 (M+H)⁺; Rt = 3.67 min.

Example 367

Methyl-phenyl-carbamic acid 5-[(pyridine-2-carbonyl)-amino]-pyridin-2-yl ester

10

A solution of pyridine-2-carboxylic acid (6-hydroxy-pyridin-3-yl)-amide hydrochloride (0.50 g, 1.99 mmol), N-methyl-N-phenylcarbonyl chloride (0.44 g, 2.59 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.54 g, 4.81 mmol) in dimethylformamide (15 mL) was stirred at room temperature for 1 hour. Water was added and the solids were isolated by suction, re-

15 dissolved in dichloromethane. The solution was dried over sodium sulphate, filtered and evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50) yielding the title compound (0.38 g, 55% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 3.45 (br.s, 3H), 7.11 (br.s, 1H), 7.26 (m, 1H), 7.40 (d, 4H), 7.51 (m, 1H), 7.92 (dt, 1H), 8.28 (d, 1H), 8.45 (dd, 1H), 8.60 (m, 2H), 10.10 (s, 1H); HPLC-MS (Method A): m/z = 349 (M+H)⁺; Rt = 3.31 min.

20

Example 368

Methyl-phenyl-carbamic acid 2-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-pyrimidin-5-yl ester

25 A solution of 1-(5-hydroxy-pyrimidin-2-yl)-4,4-dimethyl-piperidine-2,6-dione (0.60 g, 2.55 mmol), N-methyl-N-phenylcarbonyl chloride (0.48 g, 2.81 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.31 g, 2.81 mmol) in dichloromethane (25 mL) was stirred at room temperature for 15 minutes. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50), yielding the title compound as a white solid.

30

¹H NMR (300MHz, CDCl₃): δ = 1.22 (s, 6H), 2.65 (s, 4H), 3.44 (br.s, 3H), 7.28-7.48 (m, 5H), 8.68 (br.s, 2H); HPLC-MS (Method A): m/z = 369 (M+H)⁺; Rt = 3.45 min.

Example 369

35 Methyl-phenyl-carbamic acid 5-bromo-pyrimidin-2-yl ester

A mixture of 4-bromo-2-hydroxypyrimidine (0.96 g, 5.49 mmol), N-methyl-N-phenylcarbamoyl chloride (1.02 g, 6.04 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.68 g, 6.04 mmol) in dry tetrahydrofuran (15mL) was stirred at room temperature for 1 hour. Dichloromethane was added and the solution was extracted twice with water. The organic layer was dried over sodium sulphate, filtered and evaporated in vacuo yielding the title compound (1.70 g, 100% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 3.43 (br.s, 3H), 7.27 (m, 1H), 7.38 (m, 4H), 8.68 (br.s, 2H); HPLC-MS (Method A): m/z = 330 and 332 (M+H)⁺; Rt = 3.33 min.

Example 370

Methyl-phenyl-carbamic acid 5-[(6-chloro-pyridine-3-carbonyl)-amino]-pyridin-2-yl ester

A mixture of 4-chloro-N-(6-hydroxy-pyridin-3-yl)-nicotinamide (0.56 g, 2.25 mmol), N-methyl-N-phenylcarbamoyl chloride (0.51 g, 3.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.67 g, 6.00 mmol) in dry tetrahydrofuran (15mL) was stirred at room temperature for 1 hour. Dichloromethane was added and the solution was extracted twice with water. The organic layer was dried over sodium sulphate, filtered and evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 70:30) followed by crystallised from ethyl acetate:heptane, yielding the title compound (44 mg, 5% yield) HPLC-MS (Method A): m/z = 383 (M+H)⁺; Rt = 3.40 min.

Example 371

Methyl-phenyl-carbamic acid 5-(2,2-dimethyl-propylcarbamoyl)-pyridin-2-yl ester

A solution of N-(2,2-dimethyl-propyl)-6-hydroxy-nicotinamide (0.50 g, 2.40 mmol), N-methyl-N-phenylcarbamoyl chloride (0.41 g, 2.40 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.27 g, 2.40 mmol) in tetrahydrofuran (15 mL) was stirred at room temperature for 3 hours. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (first column: SiO₂, dichloromethane:ethyl acetate 95:5, second column: SiO₂, ethyl acetate), yielding the title compound (0.43 g, 52% yield).

¹H NMR (300MHz, CDCl₃): δ = 0.97 (s, 9H), 3.26 (d, 2H), 3.44 (br.s, 3H), 6.26 (br.s, 1H), 7.08 (br.s, 1H), 7.29 (m, 1H), 7.39 (m, 4H), 8.13 (br.d, 1H), 8.70 (br.s, 1H); HPLC-MS (Method A): m/z = 383 (M+H)⁺; Rt = 3.40 min.

Example 372

[Methyl-phenyl-carbamic acid 6-(3,4-dichloro-phenoxy)-pyridazin-3-yl ester

A solution of 6-(3,4-dichloro-phenoxy)-pyridazin-3-ol (0.51 g, 2.00 mmol), N-methyl-N-phenylcarbonyl chloride (0.36g, 2.10 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.24 g, 2.10 mmol) in tetrahydrofuran (15 mL) was stirred at room temperature for 2 hours. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 30:70), yielding the title compound (0.34 g, 44% yield).

¹H NMR (300MHz, CDCl₃): δ = 3.43 (br.s, 3H), 7.10 (dd, 1H), 7.20-7.51 (m, 9H); HPLC-MS (Method A): m/z = 389 (M+H)⁺; Rt = 4.56 min.

Example 373

4-(tert-Butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester

A solution of N-(6-hydroxy-pyridin-3-yl)-benzamide (214 mg, 1.00 mmol), 3-[4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium iodide (451 mg, 1.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (112 mg, 1.00 mmol) in dimethylformamide (10 mL) was stirred for 18 hours at room temperature followed by heating for 3 days at 40 °C. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 40:60), yielding the title compound (364 mg, 77% yield)

¹H NMR (300MHz, CDCl₃): δ = 0.08 (s, 6H), 0.91 (s, 9H), 1.59 (m, 2H), 1.78 (m, 2H), 3.50 (m, 1H), 3.62 (m, 2H), 3.76 (m, 1H), 4.00 (m, 1H), 6.87 (d, 1H), 7.41 (m, 2H), 7.50 (m, 1H), 7.90 (d, 2H), 8.01 (dd, 1H), 8.36 (d, 1H), 9.03 (s, 1H, NH); HPLC-MS (Method A): m/z = 456 (M+H)⁺; Rt = 5.23 min.

Example 374

4-Hydroxy-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester

Hydrofluoric acid (min. 40% in water, 0.50 mL) was added to a stirred solution of 4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester (364 mg, 0.77 mmol) in acetonitrile. After stirring overnight at room temperature the solvent was evaporated in vacuo. The residue was redissolved in dichloromethane. After addition of triethylamine (1 mL) the solution was filtered over a short pad of silicagel and washed with ethyl acetate:acetone 50:50. Evaporation of the solvent yielded the title compound (180 mg,

68% yield) as a white solid.

¹H NMR (400MHz, CDCl₃ + DMSO-d₆): δ = 1.60 (m, 2H), 1.91 (m, 2H), 3.25 (m, 2H), 4.02 (m, 1H), 4.41 (d, 1H), 7.07 (d, 1H), 7.44-7.61 (m, 3H), 7.99 (m, 2H), 8.35 (dd, 1H), 8.72 (d, 1H), 10.19 (s, 1H, NH); HPLC-MS (Method A): m/z = 342 (M+H)⁺; Rt = 2.37 min.

5

Example 375

4-Hydroxy-piperidine-1-carboxylic acid 5-trifluoromethyl-pyridin-2-yl ester

A solution of 2-hydroxy-5-trifluoromethylpyridine (0.32 g, 2.00 mmol), 4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester (0.90 g, 2.00 mmol) and triethylamine (0.20 g, 2.00 mmol) in acetonitrile (10 mL) was stirred at room temperature for three days. Hydrofluoric acid (min. 40% in water, 0.50 mL) was added and stirring was continued for 18 hours. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 75:25), followed by crystallisation from ethyl acetate:heptane, yielding the title compound (0.27 g, 47% yield) as a white solid. ¹H NMR (300MHz, CDCl₃): δ = 1.64 (m, 3H, 2 x CH + OH), 1.96 (m, 2H), 3.33 (m, 1H), 3.45 (m, 1H), 4.00 (m, 3H), 7.27 (d, 1H), 8.00 (dd, 1H), 8.64 (d, 1H); HPLC-MS (Method A): m/z = 313 (M+Na)⁺; Rt = 2.45 min.

20

Example 376

4-Hydroxy-piperidine-1-carboxylic acid 5-(4-chloro-benzoylamino)-pyridin-2-yl ester

A solution of 4-chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide (0.50 g, 2.00 mmol), 4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester (0.90 g, 2.00 mmol) and triethylamine (0.20 g, 2.00 mmol) in dimethylformamide (10 mL) was stirred overnight at room temperature. The solvent was evaporated in vacuo. The residue was dissolved in dichloromethane, filtered over a short pad of silicagel and washed with ethyl acetate:heptane 50:50. The solvent was evaporated in vacuo and the residue was redissolved in acetonitrile and hydrofluoric acid (min. 40% in water, 0.50 mL) was added. After stirring overnight at room temperature the solvent was evaporated in vacuo. Dichloromethane and triethylamine (1 mL) were added and the solution was extracted twice with water, dried over sodium sulphate, filtered and evaporated in vacuo. Crystallisation from ethyl acetate:heptane yielded the title compound (321 mg, 43% yield).

¹H NMR (300MHz, DMSO-d₆): δ = 1.50 (m, 2H), 1.80 (m, 2H), 3.16 (m, 1H), 3.31 (m, 1H), 3.75 (m, 2H), 3.88 (m, 1H), 4.81 (d, 1H), 7.18 (d, 1H), 7.62 (d, 2H), 8.01 (d, 2H), 8.24 (dd,

35

1H), 8.66 (d, 1H), 10.58 (s, 1H).; HPLC-MS (Method A): m/z = 376 (M+H)⁺; Rt = 2.81 min.

Example 377

4-Hydroxy-piperidine-1-carboxylic acid 5-(3-methoxy-benzoylamino)-pyridin-2-yl ester

5

Starting from N-(6-hydroxy-pyridin-3-yl)-3-methoxy-benzamide (0.49 g, 2.00 mmol) and using the procedure as described in Example 376 yielded the title compound (347 mg, 47% yield).

¹H NMR (300MHz, CDCl₃ + DMSO-d₆): δ = 1.51 (m, 2H), 1.82 (m, 2H), 3.16 (m, 1H), 3.30 (m, 1H), 3.79 (s, 3H + m, 2H), 3.92 (m, 1H), 4.28 (br.s, 1H), 6.98 (m, 2H), 7.30 (t, 1H), 7.47 (m, 10 2H), 8.24 (dd, 1H), 8.64 (d, 1H), 10.04 (s, 1H); HPLC-MS (Method A): m/z = 372 (M+H)⁺; Rt = 2.58 min.

Example 378

4-Hydroxy-piperidine-1-carboxylic acid 5-(4-methoxy-benzoylamino)-pyridin-2-yl ester

15

Starting from N-(6-hydroxy-pyridin-3-yl)-4-methoxy-benzamide (0.49 g, 2.00 mmol) and using the procedure as described in Example 376 yielded the title compound (297 mg, 40% yield).

¹H NMR (300MHz, CDCl₃): δ = 1.61 (m, 2H), 1.92 (m, 2H), 3.25 (m, 1H), 3.39 (m, 1H), 3.89 (s, 3H + m, 3H), 4.03 (m, 1H), 6.96 (d, 2H), 7.06 (d, 1H), 7.98 (d, 2H), 8.34 (dd, 1H), 8.67 (d, 20 1H), 9.72 (s, 1H); HPLC-MS (Method A): m/z = 372 (M+H)⁺; Rt = 2.53 min.

Example 379

4-Hydroxy-piperidine-1-carboxylic acid 5-(2,4-dichloro-benzoylamino)-pyridin-2-yl ester

25 Starting from 2,4-dichloro-N-(6-hydroxy-pyridin-3-yl)-benzamide (0.49 g, 2.00 mmol) and using the procedure as described in Example 376 yielded the title compound (367 mg, 45% yield).

¹H NMR (300MHz, CDCl₃): δ = 1.62 (m, 2H), 1.90 (m, 2H), 3.24 (m, 1H), 3.39 (m, 1H), 3.88 (m, 3H), 4.01 (m, 1H), 7.03 (d, 1H), 7.32 (d, 1H), 7.47 (s, 1H), 7.53 (d, 1H), 8.28 (dd, 1H), 30 8.58 (d, 1H), 10.1 (s, 1H); HPLC-MS (Method A): m/z = 410 and 412 (M+H)⁺; Rt = 2.99 min.

Example 380

4-Hydroxy-piperidine-1-carboxylic acid 5-(4-trifluoromethyl-benzoylamino)-pyridin-2-yl ester

35 Starting from N-(6-hydroxy-pyridin-3-yl)-4-trifluoromethyl-benzamide (0.56 g, 2.00 mmol) and

using the procedure as described in Example 376 yielded the title compound (367 mg, 45% yield).

¹H NMR (300MHz, CDCl₃): δ = 1.61 (m, 2H), 1.93 (m, 2H), 3.27 (m, 1H), 3.41 (m, 1H), 3.67 (br.s, 1H), 3.91 (m, 2H), 4.05 (m, 1H), 7.08 (d, 1H), 7.74 (d, 2H), 8.14 (d, 2H), 8.37 (dd, 1H), 8.67 (d, 1H), 10.11 (s, 1H); HPLC-MS (Method A): m/z = 410 (M+H)⁺; Rt = 3.18 min.

Example 381

4-Hydroxy-piperidine-1-carboxylic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3]bipyridinyl-6'-yl ester

A solution of 6'-hydroxy-4,4-dimethyl-4,5-dihydro-3H-[1,3]bipyridinyl-2,6-dione (468 mg, 2.00 mmol), 4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carboxylic acid 5-benzoylamino-pyridin-2-yl ester (0.90 g, 2.00 mmol) and triethylamine (0.20 g, 2.00 mmol) in dimethylformamide (10 mL) was stirred at room temperature for 18 hours. The solvent was evaporated in vacuo. The residue was dissolved in ethyl acetate, filtered over a short pad of silicagel and washed with ethyl acetate. The solvent was evaporated in vacuo and the residue was dissolved in 1N HCl in ethyl acetate (10 mL, 10.0 mmol). After stirring for 1.5 hours at room temperature the solvent was evaporated in vacuo. The white solid was washed with a small amount of ethyl acetate and diethyl ether and dissolved in a few millilitres of dichloromethane and triethylamine (1 mL). Purification by flash column chromatography (SiO₂, ethyl acetate:acetone 90:10) yielded the title compound (182 mg, 25% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 1.22 (s, 6H), 1.61 (m, 2H), 1.92 (m, 2H), 2.70 (s, 4H), 3.28 (m, 1H), 3.40 (m, 1H), 3.63 (d, 1H), 3.84-4.08 (m, 3H), 7.21 (d, 1H), 7.51 (dd, 1H), 8.07 (d, 1H); HPLC-MS (Method A): m/z = 362 (M+H)⁺; Rt = 2.25 min.

Example 382

Methyl-phenyl-carbamic acid 2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3]bipyridinyl-6'-yl ester

A solution of N-methyl-N-phenyl carbamoyl chloride (170 mg, 1.00 mmol), 6'-hydroxy-4,5-dihydro-3H-[1,3]bipyridinyl-2,6-dione (206 mg, 1.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (112 mg, 1.00 mmol) in tetrahydrofuran (10 mL) was stirred at room temperature for 18 hours. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, first ethyl acetate:heptane 75:25 followed by pure ethyl acetate). Evaporation of the solvent yielded the title compound (260 mg, 77% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 2.10 (quintet, 2H), 2.81 (t, 4H), 3.53 (br.s, 3H), 7.12 (br.s,

1H), 7.27 (m, 1H), 7.38 (m, 4H), 7.50 (d, 1H), 8.09 (s, 1H); HPLC-MS (Method A): m/z = 340 (M+H)⁺; Rt = 2.89 min.

Example 383

5 Methyl-phenyl-carbamic acid 5-(2,5-dioxo-pyrrolidin-1-yl)-pyridin-2-yl ester

A solution of N-methyl-N-phenyl carbamoyl chloride (170 mg, 1.00 mmol), 1-(6-hydroxy-pyridin-3-yl)-pyrrolidine-2,5-dione (192 mg, 1.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (112 mg, 1.00 mmol) in tetrahydrofuran (10 mL) was stirred at room temperature for 4 hours.

10 The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 80:20) yielding the title compound (275 mg, 85% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 2.89 (s, 4H), 3.44 (br.s, 3H), 7.14 (br.s, 1H), 7.27 (m, 1H), 7.39 (m, 4H), 7.74 (br.d, 1H), 8.38 (s, 1H); HPLC-MS (Method A): m/z = 326 (M+H)⁺; Rt =
15 2.78 min.

Example 384

Methyl-phenyl-carbamic acid 5-(4-trifluoromethyl-benzoylamino)-pyridin-2-yl ester

20 A solution of N-(6-hydroxy-pyridin-3-yl)-4-trifluoromethyl-benzamide (1.41 g, 5.00 mmol), N-methyl-N-phenylcarbamoyl chloride (0.85 g, 5.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.56 g, 5.00 mmol) in dimethylformamide (10 mL) was stirred at room temperature overnight. The solvent was evaporated in vacuo and the residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50) yielding the title compound (1.34 g, 64% yield)
25 as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 3.35 + 3.51 (2 x br.s, 3H), 6.83 (br.s, 1H), 7.24-7.42 (m, 5H), 7.60 (d, 2H), 7.98 (d, 3H), 8.33 (s, 1H), 9.03 + 9.18 (2 x br.s, 1H, NH); HPLC-MS (Method A): m/z = 416 (M+H)⁺; Rt = 4.14 min.

30 Example 385

Methyl-phenyl-carbamic acid quinolin-6-yl ester

A solution of 6-hydroxyquinoline (1.00 g, 6.89 mmol), N-methyl-N-phenylcarbamoyl chloride (1.17 g, 6.89 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.77 g, 6.89 mmol) in dichloro-
35 methane (20 mL) was stirred at room temperature for 1.25 hours. More dichloromethane was

added and the solution was extracted with water. The organic layer was dried over sodium sulphate, filtered and evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50). Evaporation of the solvent and recrystallised from ethylacetate/heptane yielded the title compound (1.33 g, 69% yield) as a white solid.

5 ¹H NMR (300MHz, CDCl₃): δ = 3.48 (s, 3H), 7.22-7.64 (m, 8H), 8.09 (d, 2H), 8.87 (m, 1H); HPLC-MS (Method A): m/z = 279 (M+H)⁺; Rt = 2.56 min.

Example 386

4-Hydroxy-piperidine-1-carboxylic acid 5-(5-trifluoromethyl-pyridin-2-yloxy)-pyridin-2-yl ester

10

A solution of 5-(5-trifluoromethyl-pyridin-2-yloxy)-pyridin-2-ol (180 mg, 0.70 mmol), 3-[4-(tert-butyl-dimethyl-silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium iodide (317 mg, 0.70 mmol) and triethylamine (98 µL) in acetonitrile (10 mL) was stirred at room temperature overnight. The solvent was evaporated in vacuo. The residue was redissolved in some dichloromethane, filtered over a short pad of silicagel and washed with ethyl acetate:heptane
15 50:50. The solvent was evaporated in vacuo and the residue was dissolved in 3.2N HCl in ether (10 mL). After stirring for 1 hour at room temperature the solvent was evaporated in vacuo and the residue was dissolved in dichloromethane. Triethylamine (0.5 mL) was added and the solution was extracted with water. The organic layer was dried over sodium sulphate,
20 filtered and evaporated in vacuo to yield the title compound (174 mg, 65% yield) as a thick oil.

¹H NMR (300MHz, CDCl₃): δ 1.63 (m, 2H), 1.79 (br.s, 1H, OH), 1.97 (m, 2H), 3.31 (m, 1H), 3.43 (m, 1H), 3.89-4.12 (m, 3H), 7.09 (d, 1H), 7.18 (d, 1H), 7.62 (dd, 1H), 7.94 (dd, 1H), 8.23 (d, 1H), 8.39 (d, 1H); HPLC-MS (Method A): m/z = 384 (M+H)⁺; Rt = 3.00 min.

25

Example 387

4-Hydroxy-piperidine-1-carboxylic acid 5-(3,5-dichloro-pyridin-2-yloxy)-pyridin-2-yl ester

Starting from 5-(3,5-dichloro-pyridin-2-yloxy)-pyridin-2-ol (265 mg, 1.03 mmol) and using the
30 procedure as described in Example 386 yielded the title compound (121 mg, 54% yield) as a thick oil.

¹H NMR (300MHz, CDCl₃): δ = 1.62 (m, 2H), 1.94 (m, 2H), 3.06 (br.s, 1H), 3.28 (m, 1H), 3.41 (m, 1H), 3.90 (m, 2H), 4.01 (m, 1H), 7.18 (d, 1H), 7.62 (dd, 1H), 7.79 (d, 1H), 7.94 (d, 1H), 8.22 (d, 1H); HPLC-MS (Method A): m/z = 384 (M+H)⁺; Rt = 3.35 min.

35

Example 388

Methyl-phenyl-carbamic acid 5-(4-chloro-benzoylamino)-pyridin-2-yl ester

A solution of 4-chloro-N-(6-hydroxy-pyridin-3-yl)-benzamide (0.50 g, 2.01 mmol), N-methyl-N-phenylcarbonyl chloride (0.34 g, 2.01 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.22 g, 0.34 mmol) in dimethylformamide (10 mL) was stirred at room temperature for 2 hours. Water (100 mL) was added and a thick oil was being formed. The water was decanted and the residue dissolved in dichloromethane, dried over sodium sulphate, filtered and evaporated in vacuo. Ethyl acetate was added and the solution was heated briefly (some of the compound does not dissolve). The solvent was decanted and heptane was added to it. After standing overnight, the crystals were isolated by suction, washed with heptane and dried in a vacuum oven at 45 °C. Further purification by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50), yielded the title compound (0.41 g, 53% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 3.38 (br.s, 3H), 6.84 (br.s, 1H), 7.20-7.42 (m, 7H), 7.80 (d, 2H), 7.97 (dd, 1H), 8.32 (d, 1H), 9.09 (br.s, 1H); HPLC-MS (Method A): m/z = 382 (M+H)⁺; Rt = 3.63 min.

Example 389

4 Methyl-phenyl-carbamic acid 5-(4-methoxy-benzoylamino)-pyridin-2-yl ester

A solution of N-(6-hydroxy-pyridin-3-yl)-4-methoxy-benzamide (1.22 g, 5.00 mmol), N-methyl-N-phenylcarbonyl chloride (0.85 g, 5.00 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.56 g, 5.00 mmol) in dimethylformamide (20 mL) was stirred at room temperature for 2 hours. Water (100 mL) was added. The solids were isolated by suction and washed with water. Crystallisation from ethyl acetate:heptane yielded the title compound (1.08 g, 57% yield).

¹H NMR (300MHz, CDCl₃): δ = 3.42 (br.s, 3H), 3.86 (s, 3H), 6.89 (d, 2H + br.s, 1H), 7.27 (m, 1H), 7.39 (m, 4H), 7.84 (d, 2H), 8.11 (dd, 1H), 8.34 (d, 1H), 8.51 (br.s, 1H); HPLC-MS (Method A): m/z = 378 (M+H)⁺; Rt = 3.55 min.

Example 390

Methyl-phenyl-carbamic acid 4,4-dimethyl-3,4,5,6-tetrahydro-2H-[1,3]bipyridinyl-6'-yl ester

A solution of 4,4-dimethyl-3,4,5,6-tetrahydro-2H-[1,3]bipyridinyl-6'-ol (0.86 g, 4.17 mmol), N-methyl-N-phenylcarbonyl chloride (0.71 g, 4.17 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.47 g, 4.17 mmol) in dichloromethane (10 mL) was stirred at room temperature for 2 hours.

Extra dichloromethane was added and the solution was washed with water, dried over sodium sulphate, filtered and evaporated in vacuo. Crystallisation from ethyl acetate:heptane yielded the title compound (0.58 g, 41% yield)

¹H NMR (300MHz, CDCl₃): δ = 0.99 (s, 6H), 1.52 (m, 4H), 3.16 (m, 4H), 3.43 (br.s, 3H), 6.92 (br.d, 1H), 7.20-7.41 (m, 6H), 7.98 (d, 1H); HPLC-MS (Method A): m/z = 340 (M+H)⁺; Rt = 4.21 min.

Example 391

Methyl-phenyl-carbamic acid 2-methyl-quinolin-6-yl ester

10

A solution of 2-methylquinolin-6-ol (1.00 g, 6.28 mmol), N-methyl-N-phenylcarbamoyle chloride (1.07 g, 6.28 mmol) and 1,4-diazabicyclo[2,2,2]octane (0.70 g, 6.28 mmol) in dichloromethane (10 mL) was stirred at room temperature for 18 hours. Extra dichloromethane was added and the solution was extracted twice with water, dried over sodium sulphate and filtered. Some ethyl acetate and heptane were added and the solution was slowly evaporated

15

in vacuo yielding the title compound (1.64 g, 89% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 2.71 (s, 3H), 3.45 (br.s, 3H), 7.25 (m, 2H), 7.40 (m, 5H), 7.54 (s, 1H), 7.98 (t, 2H); HPLC-MS (Method A): m/z = 293 (M+H)⁺; Rt = 2.16 min.

20

Example 392 (General procedure 17){2-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-ethyl}-carbamic acid tert-butyl ester

Tyramin was N-Boc protected as described in J. Org. Chem, 49, 1984, 1016. To a solution of the N-Boc protected tyramin (10 mmol) in CH₂Cl₂ (50 mL) was added N-methyl-N-phenylcarbamoyle chloride (15 mmol) and DABCO (15 mmol) at room temperature. The reaction mixture was stirred for 16 hours at rt, added CH₂Cl₂ (20 mL) and washed with aqueous citric acid (5%) and brine. The organic phase was separated, dried (MgSO₄) and evaporated to give the crude product which was purified by FC (Quad flash 40 MeOH-CH₂Cl₂ 5:95) to give 3.45 g (93%) of the title compound as colorless crystals.

25

30 HPLC-MS : m/z = 393.4 (M+Na); Rt = 4.44 min.

Example 393 (General procedure 17)Methyl-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester

35

{2-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-ethyl}-carbamic acid tertbutyl ester(3.7 g; 10

mmol) from above was dissolved in CH_2Cl_2 (90 mL). Addition of TFA (6 mL) and stirring for 4 h. The reaction mixture was evaporated to dryness and dried in vacuo at 50 °C overnight producing the title compound as a TFA salt in quantitative yield as yellow hygroscopic crystals.

5 HPLC-MS : m/z = 271.1 ($M+1$); R_t = 2.17 min.

Example 394 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester

10 The title compound was prepared in 39% yield as a clear oil using toluenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 425.2 ($M+1$); R_t = 4.33 min.

Example 395 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(5-dimethylamino-naphthalene-1-sulfonylamino)-ethyl]-phenyl ester

The title compound was prepared in 29% yield as a yellow fluorescent oil using 5-dimethylamino-naphthalene-1-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 505.1 ($M+1$); R_t = 4.58 min.

20

Example 396 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(3,4-difluorobenzenesulfonylamino)-ethyl]-phenyl ester

25 The title compound was prepared in 40% yield as a clear oil using 3,4-difluorobenzenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 447.1 ($M+1$); R_t = 4.47 min.

Example 397 (General procedure 17) 2-[2-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-ethylsulfamoyl]-benzoic acid methyl ester

30

The title compound was prepared in 30% yield as an oil using 2-chlorosulfonyl-benzoic acid methyl ester as the aryl sulfonyl chloride.

HPLC-MS : m/z = 469.1 ($M+1$); R_t = 4.37 min.

35 **Example 398** (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(2,5-dichloro-

thiophene-3-sulfonylamino)-ethyl]-phenyl ester

The title compound was prepared in 39% yield as an oil using 2,5-dichloro-thiophene-3-sulfonyl chloride as the aryl sulfonyl chloride.

5 HPLC-MS : m/z = 487.0 (M+1); R_t = 4.80 min.

Example 399 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-phenyl ester

10 The title compound was prepared in 12% yield as crystals using 5-pyridin-2-yl-thiophene-2-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 494.0 (M+1); R_t = 4.42 min.

Example 400 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(1-methyl-1H-

15 imidazole-4-sulfonylamino)-ethyl]-phenyl ester

The title compound was prepared in 42% yield as a yellow oil using 1-methyl-1H-imidazole-4-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 415.1 (M+1); R_t = 3.31 min.

20

Example 401 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonylamino)-ethyl]-phenyl

The title compound was prepared in 22% yield as an oil using 5-chloro-1,3-dimethyl-1H-

25 pyrazole-4-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 463.1 (M+1); R_t = 3.91 min.

Example 402 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(4-nitro-benzenesulfonylamino)-ethyl]-phenyl ester

30

The title compound was prepared in 25% yield as a yellow oil using 4-nitro-benzenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 456.0 (M+1); R_t = 4.19 min.

35 **Example 403** (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(6-chloro-

imidazo[2,1-b]thiazole-5-sulfonylamino)-ethyl]-phenyl ester

The title compound was prepared in 18% yield as an oil using 6-chloro-imidazo[2,1-b]thiazole-5-sulfonyl chloride as the aryl sulfonyl chloride.

5 HPLC-MS : m/z = 490.9 (M+1); R_t = 3.90 min.

Example 404 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(2-trifluoromethoxy-benzenesulfonylamino)-ethyl]-phenyl ester

10 The title compound was prepared in 38% yield as an oil using 2-trifluoromethoxy-benzenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 495.1 (M+1); R_t = 4.49 min.

Example 405 (General procedure 17)

15 Methyl-phenyl-carbamic acid 4-(2-dimethylaminosulfonylamino-ethyl)-phenyl ester

The title compound was prepared in 32% yield as an oil using dimethylaminosulfamoyl chloride as the sulfonyl chloride.

HPLC-MS : m/z = 378.1 (M+1); R_t = 3.62 min.

20

Example 406 (General procedure 17) Methyl-phenyl-carbamic acid 4-(2-methanesulfonylamino-ethyl)-phenyl ester

25 The title compound was prepared in 22% yield as an oil using methanesulfonyl chloride as the sulfonyl chloride.

HPLC-MS : m/z = 349.0 (M+1); R_t = 3.26 min.

Example 407 (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(6-morpholin-4-yl-pyridine-3-sulfonylamino)-ethyl]-phenyl ester

30

The title compound was prepared in 55% yield as crystals using 6-morpholin-4-yl-pyridine-3-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 497.0 (M+1); R_t = 3.90 min.

35 **Example 408** (General procedure 17) Methyl-phenyl-carbamic acid 4-[2-(6-phenoxy-pyridine-3-sulfonylamino)-ethyl]-phenyl ester

The title compound was prepared in 54% yield as crystals using 6-phenoxy-pyridine-3-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : $m/z = 504.4$ ($M+1$); $R_t = 4.45$ min.

5 **Example 409** Methyl-phenyl-carbamic acid 4-[2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonylamino]-ethyl]-phenyl ester

To a stirred solution of 1-(4-bromophenyl)-4-methylpiperazine (2.05 g, 8.0 mmol) in THF (10 mL) was added dropwise 1.57 M solution in hexanes *n*-BuLi (4.6 mL, 7.2 mmol) over a 5-min
10 period at -78°C . The mixture was stirred at -78°C for 15 min. Then gaseous sulphur dioxide (ca. 5 g) was added causing an immediate precipitation. The mixture was allowed to warm to room temperature and stirred for 1 h. The precipitated lithium; 4-(4-methyl-piperazin-1-yl)-benzenesulfinate was isolated by filtration under N_2 (g), washed with THF (20 mL) and dried
15 *in vacuo* providing 1.83 g (96%) of the lithium sulfinate as a solid. This lithium sulfinate (83 mg, 0.34 mmol) was suspended in CH_2Cl_2 (1 mL) and stirred with NCS (45 mg, 0.34 mmol) for 10 min at rt. A solution of N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.26 mmol) in CH_2Cl_2 (1.5 mL) was added together with DIPEA (0.90 mmol). The mixture was stirred at rt for 16 h and quenched with HOAc (2 mL) and water (2 mL). Ex-
20 traction with CH_2Cl_2 and drying of the combined organic extracts gave the crude product which was purified by preparative HPLC (Gilson). This gave 54 mg (33%) of the title compound as its TFA salt as colorless crystals.

HPLC-MS : $m/z = 509.0$ ($M+1$); $R_t = 2.88$ min.

Example 410 Methyl-phenyl-carbamic acid 4-[2-(4-dimethylamino-benzenesulfonylamino)-ethyl]-phenyl ester

25

To a stirred solution of (4-bromo-phenyl)-dimethylamine (4.02 g, 20 mmol) in THF (25 mL) was added dropwise 1.57 M solution in hexanes *n*-BuLi (11.5 mL, 18 mmol) over a 5-min pe-
30 riod at -78°C . The mixture was stirred at -78°C for 15 min. Then gaseous sulphur dioxide (ca. 5 g) was added causing an immediate precipitation. The mixture was allowed to warm to room temperature and stirred for 1 h. The precipitated lithium sulfinate was isolated by filtra-
35 tion under N_2 (g), washed with THF (20 mL) and dried *in vacuo* providing 2.99 g (87%) of lithium; 4-dimethylamino-benzenesulfinate as a bluegreen solid. This lithium sulfinate (65 mg, 0.34 mmol) was suspended in CH_2Cl_2 (1 mL) and stirred with NCS (45 mg, 0.34 mmol) for 10 min at rt. A solution of N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its
TFA salt (0.26 mmol) in CH_2Cl_2 (1.5 mL) was added together with DIPEA (0.90 mmol). The

mixture was stirred at rt for 16 h and quenched with HOAc (2 mL) and water (2 mL). Extraction with CH₂Cl₂ and drying of the combined organic extracts gave the crude product which was purified by preparative HPLC (Gilson). This gave 37 mg (31%) of the title compound as an oil.

5 HPLC-MS : m/z = 454.5 (M+1); R_t = 4.16 min.

Example 411

Methyl-phenyl-carbamic acid 4-{2-[4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfonylamino]-ethyl}-phenyl ester

10 To a stirred solution of 1-[2-(4-bromo-phenoxy)-ethyl]-pyrrolidine (6.72 g, 25 mmol) in THF (45 mL) was added dropwise 1.6 M solution in hexanes *n*-BuLi (14 mL, 22.4 mmol) over a 5-min period at -78°C. The mixture was stirred at -78°C for 15 min. Then gaseous sulphur dioxide (ca. 6 g) was added causing an immediate precipitation. The mixture was allowed to warm to room temperature and stirred for 1 h. The precipitated lithium sulfinate was isolated
15 by filtration under N₂ (g), washed with THF (40 mL) and dried *in vacuo* providing 5.04 g (78%) of lithium; 4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfinate as a solid. This lithium sulfinate (179 mg, 0.69 mmol) was suspended in CH₂Cl₂ (2 mL) and stirred with NCS (80 mg, 0.60 mmol) for 10 min at rt. A solution of N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.55 mmol) in CH₂Cl₂ (3 mL) was added together with
20 DIPEA (2.0 mmol). The mixture was stirred at rt for 16 h and evaporated to dryness. It was then redissolved in MeCN and purified by preparative HPLC (Gilson). This gave 89 mg (25%) of the title compound as its TFA salt as a crystals.

HPLC-MS : m/z = 524.5 (M+1); R_t = 3.04 min.

25 Example 412 (General procedure 15)

4-(Tetrahydrofuran-2-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(tetrahydrofuran-2-ylmethyl)-piperazine, crude yield 0.28 g
30 (89%). The crude product was stirred with a mixture of ethyl acetate (10 ml) and a solution of sodium bicarbonate (0.05 g) in water (5 ml). The aqueous layer was extracted with ethyl acetate (10 + 5 ml), the combined organic phases were washed with brine (5 ml), dried over sodium sulfate, filtered and evaporated. The residue was triturated with heptane (3 ml), filtered
35 and dried to give the title compound, White crystals, m.p. 98 - 100 °C; ¹H NMR (CDCl₃) δ

8.52 - 8.37 (br, 1H), 7.98 - 7.83 (dd, 1H), 7.25 - 7.06 (m, 4H), 7.00 (d, J = 8.5 Hz), 4.18 - 3.47 (m, 7H), 2.76 - 2.36 (m, 6H), 2.13 - 1.74 (m, 3H), 1.64 - 1.40 (m, 1H); IR (KBr): ν 1715 (C=O) cm^{-1} .

5 **Example 413** (General procedure 15)

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10 The hydrochloride of the title compound was prepared from 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-cyclopropylmethyl-piperazine, yield 83%. White crystals, m.p. 254-255°C; IR (KBr): ν 1728 (C=O) cm^{-1} .

Example 414 (General procedure 15)

15 4-(Tetrahydro-furan-2-ylmethyl)-piperazine-1-carboxylic acid 4-(4-trifluoromethylphenoxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenyl chloroformate and 1-(tetrahydro-furan-2-ylmethyl)-piperazine, yield 93%. White crystals, m.p. 216-217°C; IR (KBr): ν 1730 (C=O) cm^{-1} .

20

Example 415 (General procedure 15)

4-Cyclohexylmethyl-piperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester

25 The hydrochloride of the title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenyl chloroformate and 1-cyclohexylmethyl-piperazine, yield 93%. White crystals, m.p. 256-258 °C; IR (KBr): ν 1715 (C=O) cm^{-1} .

Example 416 (General procedure 15)

30 4-Cyclohexylmethyl-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-cyclohexylmethyl-piperazine, yield 76%. White crystals, m.p. 265-266 °C; IR (KBr): ν 1732 (C=O) cm^{-1} .

35

Example 417 (General procedure 15)

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(4-trifluoromethyl-phenoxy)-phenyl ester

5 The hydrochloride of the title compound was prepared from 4-(4-trifluoromethyl-phenoxy)-phenyl chloroformate and 1-cyclopropylmethyl-piperazine, yield 43%. White crystals, m.p. 238-239 °C; IR (KBr): ν 1725 (C=O) cm^{-1} .

Example 418 (General procedure 15)

4-(Tetrahydrofuran-2-ylmethyl)-piperazine-1-carboxylic acid 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10

The hydrochloride of the title compound was prepared from 4-(3-chloro-5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(tetrahydrofuran-2-ylmethyl)-piperazine, yield 23%. White crystals, m.p. 98-100 °C; IR (KBr): ν 1731 (C=O) cm^{-1} .

15 **Example 419** (General procedure 15)

4-Naphthalen-1-ylmethyl-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

20 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-naphthalen-1-ylmethyl-piperazine, yield 71%. White crystals, m.p. 218-219 °C; IR (KBr): ν 1713 (C=O) cm^{-1} .

Example 420 (General procedure 15)

25 4-(2-Cyclohexyl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-cyclohexyl-ethyl)-piperazine, yield 90%. White crystals, m.p. 274-276 °C; IR (KBr): ν 1715 (C=O) cm^{-1} .

30

Example 421 (General procedure 15)

4-(3-Methoxy-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

35 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate

mate and 1-(3-methoxy-phenyl)-piperazine, yield 96%. White solid, m.p. 109-111 °C; IR (KBr): ν 1721 (C=O) cm^{-1} .

Example 422 (General procedure 15)

5 4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-phenyl ester

The title compound was prepared from 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-phenyl chloro-
formate and 1-cyclopropylmethyl-piperazine, yield 18%. White crystals, m.p. 225-226 °C; IR
10 (KBr): ν 1710 (ester C=O), 1661 (amide C=O) cm^{-1} .

Example 423 (General procedure 15)

4-(Tetrahydro-furan-2-ylmethyl)-piperazine-1-carboxylic acid 4-[2-(4-chloro-phenyl)-
ethylcarbamoyl]-phenyl ester

15

The title compound was prepared from 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-phenyl chloro-
formate and 1-(tetrahydrofuran-2-ylmethyl)-piperazine, yield 12%. White crystals, m.p. 220-
221 °C; IR (KBr): ν 1708 (ester C=O), 1660 (amide C=O) cm^{-1} .

20 **Example 424** (General procedure 15)

4-(3,4-Dichloro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl chloroformate and 1-(3,4-dichloro-benzyl)-piperazine, yield 86%. White crys-
25 tals, m.p. 229-230 °C; IR (KBr): ν 1717 (C=O) cm^{-1} .

Example 425 (General procedure 15)

4-Cyclopropylmethyl-[1,4]diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-
30 phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-
yloxy)-phenyl chloroformate and 1-cyclopropylmethyl-[1,4]diazepane, yield 61%. White crys-
tals, m.p. 208-209 °C; IR (KBr): ν 1711 (C=O) cm^{-1} .

35

Example 426 (General procedure 15)

4-(2-Pyridin-2-yl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 5 The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-pyridin-2-yl-ethyl)-piperazine, yield 81%. White crystals, m.p. 281-282 °C; IR (KBr): ν 1713 (C=O) cm^{-1} .

Example 427 (General procedure 18)

- 10 4-(Pyrazin-2-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(pyrazin-2-yl)-piperazine. White crystals, m.p. 160-161 °C; HPLC-MS: m/z = 446 (M+1) at R_t = 4.0 min.; ^1H NMR (DMSO- d_6): δ 8.60 - 8.57 (m, 1H), 8.39 - 8.36 (m, 1H), 8.27 - 8.22 (dd-like, 1H), 8.14 - 8.11 (m, 1H), 7.91 - 7.87 (d-like, 1H), 7.29 - 7.21 (m, 5H), 3.80 - 3.64 (br, 6H), 3.64 - 3.50 (br, 2H); IR (KBr): ν 1737, 1714 (C=O) cm^{-1} .

Example 428 (General procedure 15)

- 20 4-(Benzo-isothiazol-3-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(benzo-isothiazol-3-yl)-piperazine. Drying on a rotary evaporator of the crude product gave the title compound as the free base. Purification by flash chromatography (silica, ethyl acetate - heptane 1:4) gave white crystals, m.p. 132-133 °C; IR (KBr): ν 1726 (C=O) cm^{-1} .

Example 429 (General procedure 15)

- 30 4-(5-Chloro-thiophen-2-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(5-chloro-thiophen-2-ylmethyl)-piperazine, yield 48 %. White crystals, m.p. 225 - 226 °C (from EtOH); ^1H NMR (DMSO- d_6): δ 12.00 (br, 1H), 8.60 - 8.55 (1H), 8.27 - 8.21 (dd-like, 1H), 7.33 - 7.20 (m, 6H), 7.20 - 7.15 (d, 1H), 4.55 (br s, 2H),

4.40 - 4.00 (br, 2H), 3.74 - 3.27 (br, 4H + H₂O); 3.27 - 2.97 (br, 2H); IR (KBr): ν 1723 (C=O) cm⁻¹.

Example 430 (General procedure 15)

5 4-(3-Trifluoromethyl-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-trifluoromethyl-phenyl)-piperazine. The crude product
10 was converted to the free base, yield 56%. White crystals, m.p. 87-88 °C; IR (KBr): ν 1719 (C=O) cm⁻¹.

Example 431 (General procedure 15)

15 4-(5-Chloro-2-methyl-phenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(5-chloro-2-methyl-phenyl)-piperazine. The crude product was converted to the free base, yield 26%. White crystals; HPLC-MS:m/z = 492(M+H) at Rt =
20 5.5 min.; IR (KBr): ν 1722 (C=O) cm⁻¹.

Example 432 (General procedure 15)

25 4-(1-Methyl-piperidin-4-ylmethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The dihydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(1-methyl-piperidin-4-ylmethyl)-piperazine, yield 6%. White crystals, m.p. 305-306 °C; IR (KBr): ν 1713 (C=O) cm⁻¹.

30 **Example 433** (General procedure 15)

4-Biphenyl-4-ylmethyl-[1,4]diazepane-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-biphenyl-4-ylmethyl-[1,4]diazepane. The crude product
35

was converted to the free base, yield 17%. White crystals, m.p. 143 °C; HPLC-MS: m/z = 548(M+H) at R_t = 3.6 min.; IR (KBr): ν 1711 (C=O) cm^{-1} .

Example 434 (General procedure XX)

5 4-(5-Dimethylamino-naphthalene-1-sulfonyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperazine and purified by flash chromatography (ethyl acetate - heptane 1:4), yield 32%. Pale crystals, HPLC-MS: m/z: 601(M+1) at R_t = 5.2 min.; m.p. °C; IR (KBr): ν 1723 (C=O) cm^{-1} .

Example 435 (General procedure 15)

4-(3-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-methoxy-benzyl)-piperazine, yield 99%. White crystals, m.p. 214-215 °C; HPLC-MS: m/z: 489(M+1) at R_t = 2.8 min.; IR (KBr): ν 1712 (C=O) cm^{-1} .

Example 436 (General procedure 15)

4-(3-Fluoro-benzyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-fluoro-benzyl)-piperazine, yield 77%. White crystals, m.p. 138-139 °C; HPLC-MS: m/z: 476 (M+1) at R_t = 3.0 min.; IR (KBr): ν 1714 (C=O) cm^{-1} .

Example 437 (General procedure 18)

4-(3-Trifluoromethyl-pyridin-2-yl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(3-trifluoromethyl-pyridin-2-yl)-piperazine, yield 25%. White crystals, m.p. 131-132 °C; HPLC-MS: m/z: 513 (M+1) at R_t = 5.0 min.; IR (KBr): ν 1722 (C=O) cm^{-1} .

Example 438 (General procedure 15)

4-(3-Fluorobenzyl)-piperazine-1-carboxylic acid 4-(4,6-dimethyl-pyrimidin-2-ylsulfanyl)-phenyl ester

5

The hydrochloride of the title compound was prepared from 4-(4,6-dimethyl-pyrimidin-2-ylsulfanyl)-phenyl chloroformate and 1-(3-fluorobenzyl)-piperazine. The crude product was converted to the free base, yield 48%. White crystals, m.p. 117-118 °C; HPLC-MS: m/z: 453 (M+1) at Rt = 2.6 min.; IR (KBr): ν 1714 (C=O) cm^{-1} .

10

Example 439 (General procedure 15)

5-(4-Trifluoromethoxybenzyl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl ester

15

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 2-(4-trifluoromethoxybenzyl)-2,5-diazabicyclo[2.2.1]heptane. The crude product was converted to the free base, yield 20%. Oil; HPLC-MS: m/z: 554 (M+1) at Rt = 3.1 min.; IR (KBr): ν 1723 (C=O) cm^{-1} .

20

Example 440 (General procedure 18)

1-Oxo-1 λ^4 -thiomorpholine-4-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and thiomorpholine 1-oxide and purified by flash chromatography (ethyl acetate), yield 37%. White crystals, m.p. 156 - 157 °C; ^1H NMR (CDCl_3) δ 8.47 - 8.40 (s-like, 1H), 7.94 - 7.87 (dd-like, 1H), 7.24 - 7.13 (m, 4H), 7.06 - 6.99 (d-like), 4.31 - 3.91 (m, 4H), 2.97 - 2.87 - 2.75 (m, 2H + 2H); IR (KBr): ν 1710 (C=O) cm^{-1} .

25

Example 441 (General procedure 18)

4-(2,4-Dimethoxyphenyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2,4-dimethoxyphenyl)-piperazine. Purified by flash chromatography (ethyl acetate - heptane 1:4), yield 69%. White crystals, m.p. 98-99 °C; HPLC-MS: m/z = 504 (M+1) at

35

Rt = 4.0 min.; IR (KBr): ν 1733, 1712 (C=O) cm^{-1} .

Example 442 (General procedure 15)

5-Benzyl-2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl ester

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 2-benzyl-2,5-diazabicyclo[2.2.1]heptane. The crude product was converted to the free base and further purified by flash chromatography (ethyl acetate - heptane 1:4), yield 16 %. White crystals, m.p. 114-115 °C; HPLC-MS: m/z: 470 (M+1) at R_t = 2.6 min.; IR (KBr): ν 1718 (C=O) cm^{-1} .

Example 443 (General procedure 10)

4-Aminomethyl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 4-aminomethylpiperidine. The crude product was purified by preparative HPLC (Method C) (32%, off-white crystals). HPLC-MS m/z = 396.1 (M+1), R_t : 2.65 min. purity: 86%.

Example 444 (General procedure 10)

4-Pyrimidin-2-yl-piperazine-1-carboxylic acid 4-(5-chloro-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(2-pyrimidinyl)-piperazine, two equivalent of diisopropylethylamine was added. The crude product was obtained by filtration of the reaction mixture. The crude product was washed with diethyl ether and subjected to column chromatography, ethyl acetate/heptane (1:3) (18%, white crystals). HPLC-MS m/z = 412.1 (M+1), R_t : 4.12 min.

Example 445 (General procedure 12)

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester

The title compound was prepared from 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl chloroformate and 1-cyclopropylmethyl-piperazine, preparative HPLC (Method C) (45%, off-white crystals). HPLC-MS m/z = 400.3 (M+1), R_t : 1.83 min.

Example 446 (General procedure 12)

4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester

5

The title compound was prepared from 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl chloroformate and 1-(4-methoxybenzyl)-piperazine, preparative HPLC (Method C) (54%, white crystals). HPLC-MS m/z = 466.3 (M+1), Rt: 2.26 min.

10 **Example 447** (General procedure 12)

4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl ester

15 The title compound was prepared from 4-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-phenyl chloroformate and 1-pyridin-3-ylmethyl-piperazine, hydrochloride, preparative HPLC (Method C) (75%, colourless oil). HPLC-MS m/z = 437.2 (M+1), Rt: 1.70 min.

Example 448 (General procedure 12)

4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester

20

The title compound was prepared from 4-(2-cyclohexyl-acetylamino)-phenyl chloroformate and 1-(4-methoxybenzyl)-piperazine, preparative HPLC (Method C) (49%, white crystals). HPLC-MS m/z = 466.3 (M+1), Rt: 2.85 min.

25

Example 449 (General procedure 12)

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester

30 The title compound was prepared from 4-(2-cyclohexyl-acetylamino)-phenyl chloroformate and 1-cyclopropylmethyl-piperazine, preparative HPLC (methode C (60%, off-white crystals). HPLC-MS m/z = 400.3 (M+1), Rt: 2.42 min.

Example 450 (General procedure 12)

4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester

35

The title compound was prepared from 4-(2-cyclohexyl-acetylamino)-phenyl chloroformate and 1-pyridin-3-ylmethyl-piperazine, hydrochloride, preparative HPLC (methode C) (64%, white crystals). HPLC-MS m/z = 437.4 (M+1), Rt: 2.38 min.

5 **Example 451** (General procedure 14)

Methyl-phenyl-carbamic acid 4-(2-cyclohexyl-ethylsulfamoyl)-phenyl ester

The title compound was prepared from N-(2-cyclohexyl-ethyl)-4-hydroxy-benzenesulfonamide and N-methyl-N-phenyl carbamoylchloride, preparative HPLC (Method
10 C) (14%, light yellow oil). HPLC-MS m/z = 417.3 (M+1), Rt: 4.89 min.

Example 452 (General procedure 14)

Methyl-phenyl-carbamic acid 4-(3-methyl-butylsulfamoyl)-phenyl ester

15 The title compound was prepared from 4-hydroxy-N-(3-methyl-butyl)-benzenesulfonamide and N-methyl-N-phenyl carbamoylchloride, preparative HPLC (Method C) (11%, light yellow oil). HPLC-MS m/z = 377.2 (M+1), Rt: 4.32 min.

Example 453 (General procedure 12)

20 (6-Methoxy-pyridin-2-yl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The crude product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 2-methoxy-6-methylaminopyridine. The reaction mixture was evaporated to
25 dryness, dissolved in dichloromethane (100 ml) and extracted with a aqueous phosphate buffer, pH 7. The aqueous phase was extracted with dichloromethane (100 ml \times 2) and the combined organic phases were dried and evaporated to dryness. The product was subjected to flash chromatography, ethyl acetate/heptane (1:7) to give the titlew product (86%, white crystals). HPLC-MS m/z = 420.4 (M+1), Rt: 5.23 min.

30

Example 454 (General procedure 12)

4-Benzimidazol-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

35 The crude product was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate

mate and 1-piperidine-4-yl-1H-benzimidazole, 5 equivalent of diisopropylamine was added, preparative HPLC (method C) (23%, colourless crystals). HPLC-MS m/z = 483.3 (M+1), Rt: 3.14 min.

5 **Example 455** (General procedure 12)

4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(2-cyclohexyl-acetylamino)-phenyl ester

The title compound was prepared from 4-(2-cyclohexyl-acetylamino)-phenyl chloroformate and 4-hydroxymethyl-piperidine, preparative HPLC (Method C) (37%, off-white crystals).

10 HPLC-MS m/z = 375.2 (M+1), Rt: 3.41 min.

Example 456 (General procedure 12)

4-(4-Amino-phenyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

15

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 4-(4-aminophenyl)piperidine, hydrochloride, preparative HPLC (Method C) (30%, light yellow crystals). HPLC-MS m/z = 458.2 (M+1), Rt: 3.42 min.

20 **Example 457** (General procedure 12)

4-(Methyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and methyl-piperidin-4-yl-pyridin-3-ylmethyl-amine, hydrochloride, preparative HPLC (Method C) (31%, orange oil). HPLC-MS m/z = 487.1 (M+1), Rt: 2.64 min.

25

Example 458 (General procedure 12)

4-(2-Oxo-pyrrolidin-1-yl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-piperidin-4-yl-pyrrolidine-2-one, hydrochloride, preparative HPLC (Method C) (55%, semi-crystalline oil). HPLC-MS m/z = 450.1 (M+1), Rt: 3.81 min.

35

Example 459 (General procedure 12)

4-(Methyl-phenethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 5 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and methyl-phenethyl-piperidin-4-yl-amine, hydrochloride, preparative HPLC (Method C) (37%, colourless oil). HPLC-MS m/z = 500.1 (M+1), Rt: 3.26 min.

Example 460

- 10 4-[(Benzyl-ethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

Benzyl-ethyl-piperidin-4-ylmethyl-amine (1.42 mmol) was dissolved in dichloromethane. 4-(5-Trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate (1.42 mmol, 0.451 g) (prepared from
15 the corresponding phenol by conventional methods) was added at room temperature. The reaction mixture was stirred overnight and evaporated to dryness. The crude product was subjected to column chromatography, ethyl acetate/heptane (1:3) → Triethylamine/ethyl acetate 1:9). The fractions containing the title product, was evaporated to dryness and HCl (g) in ethyl acetate was added and stirred for 16 hours. The solution was evaporated to dryness
20 and dried in vacuum for 16 h. to give the title product. (58%, light yellow crystals). HPLC-MS m/z = 514.1 (M+1), Rt: 3.27 min.

Example 461 (General procedure 12)

- 25 4-[Methyl-phenethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and methyl-phenethyl-piperidin-4-ylmethyl-amine, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (46%, colourless crystals). HPLC-
30 MS m/z = 514.1 (M+1), Rt: 3.31 min.

Example 462 (General procedure 12)

4-[(Cyclohexyl-methyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and cyclohexyl-methyl-piperidine-4-ylmethyl-amine, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (43%, white crystals). HPLC-MS m/z = 492.3 (M+1), Rt: 3.22 min.

5

Example 463 (General procedure 12)

4-[(Ethyl-pyridin-4-ylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 10 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and ethyl-piperidin-4-ylmethyl-pyridin-4-ylmethyl-amine, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (25%, off-white crystals). HPLC-MS m/z = 515.2 (M+1), Rt: 2.77 min.

- 15 **Example 464** (General procedure 12)

4-[(Benzyl-methyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 20 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and benzyl-methyl-piperidin-4-ylmethyl-amine, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (56%, colourless crystals). HPLC-MS m/z = 500.2 (M+1), Rt: 3.18 min.

Example 465 (General procedure 12)

- 25 4-[(Methyl-pyridin-3-ylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 30 The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and methyl-piperidine-4-ylmethyl-pyridine-4-ylmethyl-amine, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (36%, colourless crystals). HPLC-MS m/z = 501.1 (M+1), Rt: 2.74 min.

Example 466 (General procedure 12)

- 35 4-(1,3-Dihydro-isoindol-2-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 2-piperidin-4-ylmethyl-2,3-dihydro-1H-isoindole, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (63%, colourless crystals).

5 HPLC-MS m/z = 498.1 ($M+1$), Rt: 3.07 min.

Example 467

4-Benzotriazol-1-yl-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-piperidin-4-yl-1H-benzotriazole, hydrochloride, 5 equivalent of diisopropylamine was added, preparative HPLC (Method C) (7%, colourless crystals). HPLC-MS m/z = 506.2 ($M+1$), Rt: 4.56 min.

15

Example 468 (General procedure 12)

4-[(Cyclopropylmethyl-amino)-methyl]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

20

Cyclopropylmethyl-piperidine-4-ylmethyl-amine (0.65 mmol, 110 mg) was dissolved in dichloromethane. 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate (0.65 mmol, 0.208 g) (prepared from the corresponding phenol by conventional methods) was added at -15°C for 15 min and stirred at room temperature overnight and evaporated to dryness. The crude product was purified by preparative HPLC (Method C) (16%, colourless crystals). HPLC-MS m/z = 450.1 ($M+1$), Rt: 2.98 min.

25

Example 469 (General procedure 12)

4-[Methyl-(2-pyridin-2-yl-ethyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The title product was prepared from methyl-piperidin-4-yl-(2-pyridin-2-yl-ethyl)-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 5 equivalent of diisopropylamine was added, preparative HPLC (method C) (48%, colourless oil). HPLC-MS m/z = 501.1 ($M+1$), Rt: 2.77 min.

35

Example 470 (General procedure 12)

4-(Cyclohexyl-methyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 5 The title product was prepared from cyclohexyl-methyl-piperidin-4-yl-amine, dihydrochloride and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 2 equivalent of diisopropylamine; dimethylformamide/ tetrahydrofuran (2:1). The reaction mixture was evaporated to dryness and the crude product extracted with ethyl acetate from an aqueous HCl solution saturated with sodium chloride, pH 1-2. The title product was crystallized from ethyl acetate
10 during evaporation of the solvent, filtered and dried in vacuum (45%, white crystals). HPLC-MS m/z = 478.2 (M+1), Rt: 3.10 min.

Example 471 (General procedure 12)

- 15 4-(Isopropyl-methyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from isopropyl-methyl-piperidin-4-yl-amine, dihydrochloride and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate; dimethylformamide as solvent, preparative HPLC (method C) (77%, light yellow oil). HPLC-MS m/z = 438.3 (M+1), Rt: 2.77
20 min.

Example 472 (General procedure 12)

(4-Methoxy-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

- 25 The title product was prepared from N-methyl-p-anisidine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (61%, off-white crystals). HPLC-MS m/z = 419.2 (M+1), Rt: 4.67 min.

Example 473 (General procedure 12)

- 30 (2-Methoxy-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 2-methoxy-N-methylamine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (63%, colourless oil). HPLC-MS m/z = 419.2 (M+1), Rt: 4.75 min.

Example 474 (General procedure 12)

(2-Carbamoyl-4-chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester.

The title product was prepared from 5-chloro-2-(methylamino)-benzamide and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (31%,
5 light yellow oil). HPLC-MS m/z = 466.1 ($M+1$), R_t : 3.99 min.

Example 475 (General procedure 12)

(2-Carbamoyl-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

10

The title product was prepared from 2-(methylamino)benzamide and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (57%, light yellow oil).
HPLC-MS m/z = 454.2 ($M+Na$), R_t : 3.66 min.

15 **Example 476** (General procedure 12)

(2-Chloro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 2-chlor-N-methylaniline and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (48%, light yellow oil). HPLC-
20 MS m/z = 423.1 ($M+1$), R_t : 4.98 min.

Example 477 (General procedure 12)

(2,4-Difluoro-phenyl)-methyl-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

25 The title product was prepared from 2,4-difluoro-N-methylaniline and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (37%, white crystals).
HPLC-MS m/z = 425.1 ($M+1$), R_t : 4.90 min.

Example 478 (General procedure 12)

30 Methyl-(2-trifluoromethoxy-phenyl)-carbamic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from N-methyl-2-(trifluoromethoxy)-aniline and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (49%,
35 colourless oil). HPLC-MS m/z = 473.2 ($M+1$), R_t : 5.18 min.

Example 479 (General procedure 13)

Methyl-phenyl-carbamic acid 4-(1,1,3,3-tetramethyl-butylcarbamoyl)-phenyl ester

- 5 The title product was prepared from tert-octylamine and 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester, preparative HPLC (method C) (26%, white crystals). HPLC-MS m/z = 383.5 (M+1), Rt: 4.67 min.

Example 480 (General procedure 13)

- 10 Methyl-phenyl-carbamic acid 4-[(2-dimethylamino-ethyl)-methyl-carbamoyl]-phenyl ester

The title product was prepared from N,N,N'-trimethylethylendiamine and 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester, preparative HPLC (method C) (2%, colourless oil). HPLC-MS m/z = 356.2 (M+1), Rt: 2.17 min.

15

Example 481 (General procedure 13)

Methyl-phenyl-carbamic acid 4-(cyclopropylmethyl-carbamoyl)-phenyl ester

- 20 The title product was prepared from cyclopropylmethylamine and 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester. The crude product was used without further purification (100%, semi-crystal oil). HPLC-MS m/z = 325.1 (M+1), Rt: 3.40 min.

Example 482 (General procedure 13)

- 25 Methyl-phenyl-carbamic acid 4-(methyl-pyridin-3-ylmethyl-carbamoyl)-phenyl ester

- The title product was prepared from methyl-pyridin-3-ylmethyl-amine and 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester. The crude product was subjected to preparative HPLC (method C) (48%, clear colourless oil). HPLC-MS m/z = 376.2 (M+1),
30 Rt: 2.37 min.

Example 483 (General procedure 13)

Methyl-phenyl-carbamic acid 4-[(1H-benzimidazol-2-yl)methyl]-carbamoyl]-phenyl ester

- 35 The title product was prepared from C-(1H-benzimidazol-2-yl)-methylamine and 4-(methyl-

phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester. The crude product was subjected to preparative HPLC (method C) (8%, off-white crystals). HPLC-MS m/z = 401.1 ($M+1$), Rt: 2.56 min.

5 **Example 484** (General procedure 13)

Methyl-phenyl-carbamic acid 4-[2-(4-chloro-phenyl)-ethylcarbamoyl]-phenyl ester

The title product was prepared from 2-(4-chloro-phenyl)-ethylamine and 4-(methyl-phenyl-carbamoyloxy)-benzoic acid 2,5-dioxo-pyrrolidin-1-yl ester. The crude product was subjected
10 to preparative HPLC (method C) (39%, off-white crystals). HPLC-MS m/z = 409.2 ($M+1$), Rt: 4.27 min.

Example 485

4-Cyclopropylmethyl-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl
15 ester

The title product was prepared from 1-cyclopropylmethylpiperazine (0.35 mmol) dissolved in dichloromethane (5 ml). 4-(3,3-dimethyl-butylcarbamoyl)-phenyl chloroformate (0.35 mmol) was added at room temperature. The reaction mixture was stirred for 16 h and evaporated to
20 dryness and subjected to preparative HPLC (method C) (5%, white crystals). HPLC-MS m/z = 388.2 ($M+1$), Rt: 2.43 min.

Example 486

4-Hydroxymethyl-piperidine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester
25 ester

The title product was prepared from 4-hydroxymethylpiperidine (0.35 mmol) was dissolved in dichloromethane (5 ml). Diisopropylethylamine (0.35 mmol) was added together with 4-(3,3-dimethyl-butylcarbamoyl)-phenyl chloroformate (0.35 mmol) at room temperature. The reac-
30 tion mixture was stirred for 16 h and evaporated to dryness and subjected to preparative HPLC (method C). (25%, white crystals). HPLC-MS m/z = 363.3 ($M+1$), Rt: 3.27 min.

Example 487

4-Pyridin-3-ylmethyl-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl es-
35 ter

The title product was prepared from 1-pyridin-3-ylmethyl-piperazine was dissolved in dichloromethane (5 ml). 4-(3,3-Dimethyl-butylcarbamoyl)-phenyl chloroformate (0.35 mmol) was added at room temperature. The reaction mixture was stirred for 16 h and evaporated to dryness and subjected to preparative HPLC (metod C) (88%, light yellow oil). HPLC-MS m/z = 425.4 (M+1), Rt: 2.33 min.

Example 488

4-(4-Methoxy-benzyl)-piperazine-1-carboxylic acid 4-(3,3-dimethyl-butylcarbamoyl)-phenyl ester

The title product was prepared from 1-(4-methoxy-benzyl)-piperazine (0.35 mmol) was dissolved in dichloromethane (5 ml). 4-(3,3-Dimethyl-butylcarbamoyl)-phenyl chloroformate (0.35 mmol) was added at room temperature. The reaction mixture was stirred for 16 h and evaporated to dryness and subjected to preparative HPLC (metod C). (54%, colourless semi-crystalline oil). HPLC-MS m/z = 454.3 (M+1), Rt: 2.78 min.

Example 489

4-(2-Pyridin-2-yl-acetyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

a) Piperazine-1,4-dicarboxylic acid tert-butyl ester 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester (General procedure 18)

The title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-tert-butoxycarbonyl-piperazin, yield 59% (recrystallized from EtOAc - heptane 1:1). White crystals, m.p. 165 - 166 °C; ¹H NMR (CDCl₃) δ 8.45 - 8.42 (m, 1H), 7.92 - 7.87 (m, dd-like, 1H), 7.21 - 7.12 (AB-system, 4H), 7.03 - 6.98 (m, d-like, 1H), 3.72 - 3.45 (br m, 8H), 1.49 (s, 9H); ¹³C-NMR (CDCl₃) δ (ref. CDCl₃ 77.00 ppm): 165.71, 154.56, 153.51, 150.21, 148.34, 145.43 (q, J = 4.4 Hz), 136.701 (q, J = 3 Hz), 123.65 (q, J = 271.5 Hz), 122.86, 122.24, 121.62 (q, J = 33 Hz), 11.20, 80.32, 44.38, 43.78, 44.6 - 42.3 (br), 28.36 ppm; IR (KBr) δ 1722 (C=O), 1691(C=O) cm⁻¹.

b) Piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester hydrochloride

Piperazine-1,4-dicarboxylic acid tert-butyl ester 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

(0.12 g, 0.26 mmol) was stirred at 100 °C for 10 min in a mixture of ethanol (3 ml) and concentrated hydrochloric acid (1 ml). The solvent was removed in vacuum. The residue was stirred with toluene (10 ml) and evaporated, then stirred with ethanol and evaporated to dryness to give white crystals (0.10 g). Recrystallization from absolute ethanol gave the title compound. White
5 crystals, 0.068 g (65%); m.p. 279 - 282 °C (decomp.); ¹H-NMR (DMSO) δ 9.62 (br, 2H, NH₂+), 8.61 - 8.55 (m, 1H), 8.29 - 8.21 (m, dd-like, 1H), 7.30 - 7.20 (m, 5H), 3.86 and 3.70 (br s, 4H), 3.19 (br s, 4H); ¹³C-NMR (DMSO) δ (ref DMSO 39.50 ppm): 165.51, 152.71, 149.98, 148.04, 145.20 (J = 4.4 Hz), 137.62 (J = 3.7 Hz), 123.87 (J = 271.5 Hz), 123.11, 123.39, 120.39 (J = 32.2Hz), 111.80, 42.15 and a broad signal partly overlapping with the DMSO signal. IR (KBr) δ
10 1731, 1709 (C=O) cm⁻¹.

c) 4-(2-Pyridin-2-yl-acetyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

15 Triethylamine (0.026 ml) was added to a stirred solution of (pyridin-2-yl)acetic acid hydrochloride (0.033 g, 0.19 mmol) in DMF (2.5 ml) at 0 °C. 1-Hydroxy-benzotriazole containing 8% of water (0.034 g) and then 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.044 g) was added and the mixture was stirred for 50 min. Piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester hydrochloride (0.070 g) in a mixture of DMF (1 ml)
20 and triethylamine (0.032 ml) was added and stirring was continued over night at room temperature. The solvent was removed in vacuum and the residue was partitioned between dichloromethane (10 ml) and water (10 ml). The organic phase was washed with water (10 ml), dried over sodium sulfate, filtered and evaporated. The residue was triturated with ether-petroleumsether 1:1 and the (2 + 1 ml). The residue was dried to give the title compound. White
25 solid, m.p. 143 - 146 °C; HPLC-MS m/z: IR (KBr) δ 1732 (C=O), 1643 (C=O).

Example 490 (General procedure 15)

4-(2-Pyridin-4-yl-ethyl)-piperazine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

30

The hydrochloride of the title compound was prepared from 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate and 1-(4-pyridin-2-yl-ethyl)-piperazine. White crystals, m.p. 268 - 271 °C (decomp); IR (KBr) δ 2680, 2579, 2456 (N⁺-H), 1731, 1713 (C=O) cm⁻¹; HPLC-MS m/z: 473 (M+H) Rt = 2.3 min.

35

Example 491**Methyl-phenyl-carbamic acid 5-amino-pyridin-2-yl ester**

A solution of methyl-phenyl-carbamic acid 5-nitro-pyridin-2-yl ester (10.41 g, 38.1 mmol) in tetrahydrofuran (100 mL) was hydrogenated in a Parr apparatus with wet 5% palladium on carbon and 20 psi hydrogen pressure. After 2 hours the solution was filtered over a short pad of Celite, washed thoroughly with ethyl acetate and evaporated in vacuo, yielding title compound (9.51 g, 103% yield) as a thick oil, which solidified upon standing.

¹H NMR (300MHz, CDCl₃): δ = 3.40 (br.s, 3H), 3.70 (br.s, 2H), 6.80 (br.d, 1H), 6.97 (dd, 1H), 7.22 (m, 1H), 7.36 (m, 4H), 7.72 (d, 1H); HPLC-MS (Method A): m/z = 244 (M+H)⁺; R_t = 2.28 min.

Example 492**Methyl-phenyl-carbamic acid 5-benzenesulfonylamino-pyridin-2-yl ester**

Benzenesulphonyl chloride (0.18 g, 1.00 mmol), dissolved in a small amount of dichloromethane, was added dropwise to a stirred solution of methyl-phenyl-carbamic acid 5-amino-pyridin-2-yl ester (0.24 g, 1.00 mmol) and triethylamine (0.10 g, 1.00 mmol) in dichloromethane (10 mL). After stirring for 4 hours the solution was extracted with water and evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 40:60 followed by 50:50) yielding the title compound (108 mg, 23% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 3.46 (br.s, 3H), 6.82 (br.s, 1H), 7.24-7.52 (m, 9H), 7.63 (m, 3H), 7.84 (br.s, 1H); HPLC-MS (Method A): m/z = 384 (M+H)⁺; R_t = 3.26 min.

Example 493**3,3-Dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbamoyl]-butyric acid**

A solution of methyl-phenyl-carbamic acid 5-amino-pyridin-2-yl ester (243 mg, 1.00 mmol) and 3,3-dimethylglutaric anhydride (142 mg, 1.00 mmol) in dichloromethane (10 mL) was stirred at room temperature for 48 hours. Evaporation of the solvent yielded the title compound as a thick oil.

HPLC-MS (Method A): m/z = 386 (M+H)⁺; R_t = 2.76 min.

Example 494**Piperidine-1-carboxylic acid 4,4-dimethyl-2,6-dioxo-3,4,5,6-tetrahydro-2H-[1,3']bipyridinyl-6'-yl**

1-Piperidinecarbonyl chloride (63 microL, 0.50 mmol) was added to a solution of 6'-hydroxy-4,4-dimethyl-4,5-dihydro-3H-[1,3']bipyridinyl-2,6-dione (117 mg, 0.50 mmol) and DABCO (56 mg, 0.50 mmol) in dimethylformamide (10 mL). After stirring for 1 hour at room temperature water was added and the solution was extracted twice with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and evaporated in vacuo. The residue was crystallised from ethyl acetate:heptane yielding the title compound (0.12 g, 69% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 1.21 (s, 6H), 1.64 (s, 6H), 2.69 (s, 4H), 3.53 (m, 2H), 3.63 (m, 2H), 7.22 (d, 1H), 7.49 (dd, 1H), 8.09 (d, 1H); HPLC-MS (Method A): m/z = 346 (M+H)⁺; R_t = 3.12 min.

Example 495

2,2-Dimethyl-N-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-yl]-succinamic acid

A solution of methyl-phenyl-carbamic acid 5-amino-pyridin-2-yl ester (0.49 g, 2.00 mmol) and 2,2-dimethylsuccinic anhydride (0.26 g, 2.00 mmol) in dichloromethane (10 mL) was stirred at room temperature overnight. Evaporation of the solvent in vacuo yielded the title compound. HPLC-MS (Method A): m/z = 372 (M+H)⁺; R_t = 2.84 min.

Example 496

Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-2,5-dioxo-pyrrolidin-1-yl)-pyridin-2-yl ester

A mixture of thionyl chloride (0.726 mL, 10.00 mmol) and 2,2-dimethyl-N-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-yl]-succinamic acid (0.74 g, 2.00 mmol) in dichloromethane (25 mL) was stirred at room temperature for 18 hours. The solvent and excess thionyl chloride were evaporated in vacuo. The crude product was redissolved in dichloromethane (25 mL) and pyridine (316 mg, 4.00 mmol) was added. The solution was extracted with water, dried over sodium sulphate, filtered and evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, ethyl acetate:heptane 50:50), yielding the title compound (490 mg, 69% yield) as a white solid.

¹H NMR (300MHz, CDCl₃): δ = 1.43 (s, 6H), 2.74 (s, 2H), 3.45 (br.s, 3H), 7.14 (br.s, 1H), 7.26 (m, 1H), 7.38 (m, 4H), 7.76 (br.d, 1H), 8.39 (s, 1H); HPLC-MS (Method A): m/z = 354 (M+H)⁺; R_t = 3.35 min.

Example 497

Methyl-phenyl-carbamic acid 5-[3,3-dimethyl-5-(4-methyl-piperazin-1-yl)-5-oxo-pentanoylamino]-pyridin-2-yl ester

- 5 Thionyl chloride (56 microL, 0.78 mmol) was added to a stirred solution of 3,3-dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbonyl]-butyric acid (0.15 g, 0.39 mmol) in dichloromethane (5 mL) and 2 drops of dimethylformamide. After stirring for 10 minutes N-methylpiperazine (1 mL) was added and stirring was continued for 2 hours. The solvent was evaporated in vacuo. The residue was redissolved in dichloromethane and extracted with water,
10 dried over sodium sulphate, filtered and evaporated in vacuo. The crude product was purified by filtration over a short column (SiO₂, ethyl acetate followed by acetone) yielding the title compound (93 mg, 51% yield) as a thick oil.
- ¹H NMR (300MHz, CDCl₃): δ = 1.11 (s, 6H), 2.32 (s, 3H), 2.43 (m, 8H), 3.42 (br.s, 3H), 3.61 (m, 2H), 3.74 (m, 2H), 7.00 (br.s, 1H), 7.38 (d, 4H), 8.21 (dd, 1H), 8.47 (d, 1H), 10.90 (s, 1H);
15 HPLC-MS (Method A): m/z = 468 (M+H)⁺; R_t = 2.20 min.

Example 498

Methyl-phenyl-carbamic acid 5-[3,3-dimethyl-4-(pyridin-3-ylcarbonyl)-butyrylamino]-pyridin-2-yl ester

- 20 Thionylchloride (237 microL, 3.27 mmol) was added to a stirred solution of 3,3-dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbonyl]-butyric acid (0.63 g, 1.63 mmol) in dichloromethane (10 mL). After stirring for 10 minutes the solution was divided into 4 equal portions of acid chloride. To one portion was added 2-aminopyridine (0.5 mL) and after stirring for 2
25 hours at room temperature the crude product was purified by flash column chromatography (SiO₂, ethyl acetate followed by ethyl acetate:acetone 90:10), yielding the title compound (105 mg, 56% yield).
- ¹H NMR (300MHz, CDCl₃): δ = 1.09 (s, 6H), 2.36 (s, 2H), 2.38 (s, 2H), 3.44 (br.s, 3H), 6.95 (br.s, 1H), 7.26 (m, 2H), 7.38 (m, 4H), 8.05 (dd, 1H), 8.10 (dt, 1H), 8.30 (m, 2H), 8.67 (d, 1H);
30 HPLC-MS (Method A): m/z = 462 (M+H)⁺; R_t = 2.52 min.

Example 499

Methyl-phenyl-carbamic acid 5-(3,3-dimethyl-5-morpholin-4-yl-5-oxo-pentanoylamino)-pyridin-2-yl ester

Thionylchloride (237 microL, 3.27 mmol) was added to a stirred solution of 3,3-dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbamoyl]-butyric acid (0.63 g, 1.63 mmol) in dichloromethane (10 mL). After stirring for 10 minutes the solution was divided into 4 equal portions of acid chloride. To one portion was added morpholine (0.5 mL) and after stirring for 2 hours at room temperature the crude product was purified by flash column chromatography (SiO₂), yielding the title compound (117 mg, 63% yield).

¹H NMR (300MHz, CDCl₃): δ = 1.11 (s, 6H), 2.44 (s, 4H), 3.42 (br.s, 3H), 3.59 (m, 2H), 3.70 (m, 6H), 7.00 (br.s, 1H), 7.24 (m, 1H), 7.38 (d, 4H), 8.21 (dd, 1H), 8.45 (d, 1H), 10.76 (s, 1H); HPLC-MS (Method A): m/z = 455 (M+H)⁺; R_t = 2.87 min.

Example 500

Methyl-phenyl-carbamic acid 5-[4-(2-dimethylamino-ethylcarbamoyl)-3,3-dimethylbutyrylamino]-pyridin-2-yl ester

Thionylchloride (237 microL, 3.27 mmol) was added to a stirred solution of 3,3-dimethyl-4-[6-(methyl-phenyl-carbamoyloxy)-pyridin-3-ylcarbamoyl]-butyric acid (0.63 g, 1.63 mmol) in dichloromethane (10 mL). After stirring for 10 minutes the solution was divided into 4 equal portions of acid chloride. To one portion was added N,N-dimethylethylenediamine (0.5 mL) and after stirring for 2 hours at room temperature the crude product was purified by flash column chromatography (SiO₂), yielding the title compound (74 mg, 39% yield).

¹H NMR (300MHz, CDCl₃): δ = 1.10 (s, 6H), 2.22 (s, 2H), 2.27 (s, 6H), 2.44 (m, 4H), 3.37 (m, 2H), 3.43 (br.s, 3H), 6.60 (br.t, 1H), 7.00 (br.s, 1H), 7.25 (m, 1H), 7.38 (d, 4H), 8.21 (dd, 1H), 8.46 (d, 1H), 10.70 (s, 1H); HPLC-MS (Method A): m/z = 456 (M+H)⁺; R_t = 1.93 min.

Example 501 (General procedure 12)

4-[Methyl-(2-pyridin-4-yl-ethyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl ester

The title product was prepared from methyl-piperidine-4-yl-(2-pyridin-4-yl-ethyl)-amine and 4-(5-trifluoromethylpyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine was added, preparative HPLC (method C) (15%, colourless oil). HPLC-MS m/z = 501.4 (M+1), R_t: 2.04 min.

Example 502 (General procedure 12)

4-(Cyclopropyl-pyridin-4-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-

pyridin-2-yloxy)-phenyl ester

The title product was prepared from cyclopropyl-piperidine-4-yl-pyridin-4-ylmethyl-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine
5 was added, solvent: dimethylformamide. The crude reaction mixture was evaporated without addition of acetic acid, preparative HPLC (method C) (19%, yellow oil). HPLC-MS m/z = (M+1), Rt: 2.82 min.

Example 503 (General procedure 12)

10 4-[Cyclopropyl-(2-fluoro-benzyl)-amino]-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from cyclopropyl-(2-fluoro-benzyl)-piperidin-4-yl-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine was
15 added, solvent: dimethylformamide. The crude reaction mixture was evaporated without addition of acetic acid, preparative HPLC (method C) (60%, colourless oil). HPLC-MS m/z = (M+1), Rt: 3.02 min.

Example 504 (General procedure 12)

20 4-(Cyclopropyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from cyclopropyl-piperidin-4-yl-pyridin-3-ylmethyl-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine
25 was added, solvent: dimethylformamide. The crude reaction mixture was evaporated without addition of acetic acid, preparative HPLC (method C) (26%, light brown solid). HPLC-MS m/z = (M+1) 513.3, Rt: 2.64 min.

Example 505 (General procedure 12)

30 4-(Cyclopropylmethyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from cyclopropylmethyl-piperidin-4-yl-pyridin-3-ylmethyl-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine was added, solvent: dimethylformamide. The crude reaction mixture was evapo-
35 rated without addition of acetic acid, preparative HPLC (method C) (47%, off-white solid).

HPLC-MS m/z = (M+1) 513.3, Rt: 2.70 min.

Example 506 (General procedure 12)

4-(Cyclopropylmethyl-pyridin-3-ylmethyl-amino)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from cyclopropylmethyl-piperidin-4-yl-pyridin-4-ylmethyl-amine and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, 3 equivalent of diisopropylamine was added, solvent: dimethylformamide. The crude reaction mixture was evaporated without addition of acetic acid, preparative HPLC (method C) (22%, off-white solid. HPLC-MS m/z = (M+1) 513.3, Rt: 2.70 min.

Example 507 (General procedure 12)

4-(4-Hydroxy-piperidin-1-ylmethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 1-piperidin-4-ylmethyl-piperidine-4-ol (released from the correspondent hydrochloride by a standard procedure) and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (reaction performed in a mixture of dichloromethane and dimethylformamide). 1.7 M HCl in ethyl acetate was added to the pooled fractions containing the title product, and the fractions was evaporated to dryness (92%, white solid. HPLC-MS m/z = (M+1) 480.4, Rt: 2.38 min.

Example 508 (General procedure 11)

4-{3-[1-(2-Hydroxy-ethyl)-piperidin-4-yl]-propyl}-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 2-[4-(3-piperidin-4-yl-propyl)-piperidin-1-yl]-ethanol and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate. The reaction mixture was evaporated, diethyl ether (30 ml) was added and the title product isolated by filtration, washed with diethyl ether and dried to give (67%, white solid. HPLC-MS m/z = (M+1) 536.2, Rt: 3.39 min.

Example 509 (General procedure 12)

4-(2-Pyrrolidin-1-yl-ethyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester

The title product was prepared from 2-[4-(3-piperidin-4-yl-propyl)-piperidin-1-yl]-ethanol and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C). 1.7 M HCl in ethyl acetate was added to the pooled fractions containing the title product, and the fractions was evaporated to dryness (65%, white solid). HPLC-MS $m/z = (M+1)$ 464.1, Rt:

5 2.99 min.

Example 510 (General procedure 16)

Methyl-o-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester

10 The title compound was prepared from 1-hydroxy-4-iodopyrazole and methyl-o-tolylamine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (40%, oil).

HPLC-MS : $m/z = 380.1$ (M+23); $R_t = 4.05$ min.

15 **Example 511 (General procedure 16)**

Methyl-m-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and methyl-m-tolylamine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane)
20 (44%, oil).

HPLC-MS : $m/z = 380.1$ (M+23); $R_t = 4.13$ min.

Example 512 (General procedure 16)

Methyl-p-tolyl-carbamic acid 4-iodo-pyrazol-1-yl ester

25

The title compound was prepared from 1-hydroxy-4-iodopyrazole and methyl-p-tolylamine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (55%, oil).

HPLC-MS : $m/z = 380.1$ (M+23); $R_t = 4.13$ min.

30

Example 513 (General procedure 16)

(3-Chloro-phenyl)-methyl-carbamic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 3-chlorophenyl-methylamine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (54%, oil).
35

HPLC-MS : m/z = 399.9 (M+23); R_t = 4.28 min.

Example 514 (General procedure 16)

(3-Fluoro-phenyl)-methyl-carbamic acid 4-iodo-pyrazol-1-yl ester

5

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 3-fluorophenyl-methylamine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (51%, oil).

HPLC-MS : m/z = 384.0 (M+23); R_t = 4.00 min.

10

Example 515 (General procedure 16)

4-(3-Trifluoromethyl-pyridin-2-yl)-piperazine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

15

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 1-(3-trifluoromethylpyridin-2-yl)piperazine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (13%, oil).

HPLC-MS : m/z = 468.1 (M+1); R_t = 4.38 min.

Example 516 (General procedure 16)

20

2,6-Dimethyl-morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 2,6-dimethylmorpholin. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (11%, oil).

25

HPLC-MS : m/z = 374.0 (M+23); R_t = 3.24 min.

Example 517 (General procedure 16)

Thiomorpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester

30

The title compound was prepared from 1-hydroxy-4-iodopyrazole and thiomorpholin. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (27%, oil).

HPLC-MS : m/z = 340.0 (M+1); R_t = 3.22 min.

35

Example 518 (General procedure 16)

3,5-Dimethyl-morpholine-4-carboxylic acid 4-iodo-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-4-iodopyrazole and 3,5-dimethylmorpholin. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (15%, oil).

- 5 HPLC-MS : m/z = 352.0 (M+1); R_t = 3.17 min.

Example 519 (General procedure 16)

Piperidine-1-carboxylic acid 4-iodo-pyrazol-1-yl ester

- 10 The title compound was prepared from 1-hydroxy-4-iodopyrazole and piperidine. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (46%, oil).
HPLC-MS : m/z = 322.0 (M+1); R_t = 3.38 min.

Example 520 (General procedure 16)

- 15 Methyl-o-tolyl-carbamic acid 2-chloro-imidazol-1-yl ester

The title compound was prepared from 1-hydroxy-2-chloroimidazole, hydrochloride and methyl-o-tolylamine. The crude product was purified by preparative HPLC (Gilson) (4%, red oil).

- 20 HPLC-MS : m/z = 266.0 (M); R_t = 3.28 min.

Example 521 (General procedure 16)

(3-Fluoro-phenyl)-methyl-carbamic acid 2-chloro-imidazol-1-yl ester

- 25 The title compound was prepared from 1-hydroxy-2-chloroimidazole, hydrochloride and 3-fluorophenyl-methylamine. The crude product was purified by preparative HPLC (Gilson) (2%, oil).
HPLC-MS : m/z = 270.1 (M); R_t = 3.08 min.

30 **Example 522**

Methyl-phenyl-carbamic acid 4-iodo-phenyl ester

- 35 To a solution of 4-iodophenol (30 mmol) in CH_2Cl_2 (100 mL) was added N-methyl-N-phenylcarbonyl chloride (27 mmol) and diisopropylethylamine (60 mmol) at room temperature. The reaction mixture was stirred for 16 hours at rt, added CH_2Cl_2 (20 mL) and washed with aqueous citric acid (5%), aqueous Na_2CO_3 and brine. The organic phase was dried

(MgSO₄) and evaporated to give the crude product which was purified by FC (Quad flash 40 EtOAc-Heptane) to give 6.55 g (69%) of the title compound as light brown crystals.

¹H NMR (300MHz; CDCl₃): δ 3.38 (br s, 3H), 6.88 (d, 2H), 7.22-7.46 (m, 5H), 7.62 (d, 2H);

HPLC-MS : m/z = 354.0 (M+1); R_t = 4.54 min.

5

Example 523 (General procedure 20)

Methyl-phenyl-carbamic acid 4'-trifluoromethyl-biphenyl-4-yl ester

10 The title compound was prepared from methyl-phenyl-carbamic acid 4-iodo-phenyl ester and 4-trifluoromethylphenylboronic acid. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane 1:9) (42%, light brown crystals).

HPLC-MS : m/z = 372.1 (M+1); R_t = 5.19 min.

Example 524 (General procedure 20)

15 Methyl-phenyl-carbamic acid 4'-trifluoromethoxy-biphenyl-4-yl ester

The title compound was prepared from methyl-phenyl-carbamic acid 4-iodo-phenyl ester and 4-trifluoromethoxyphenylboronic acid. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane 1:9) (17%, brown oil).

20 HPLC-MS : m/z = 388.1 (M+1); R_t = 5.27 min.

Example 525 (General procedure 20)

Methyl-phenyl-carbamic acid 4-pyridin-3-yl-phenyl ester

25 The title compound was prepared from methyl-phenyl-carbamic acid 4-iodo-phenyl ester and pyridine-3-boronic acid. The crude product was purified by preparative HPLC (Gilson) (5%, brown oil).

¹H NMR (300MHz; CDCl₃): δ 3.46 (br s, 3H), 7.29-7.46 (m, 7H), 7.61 (d, 2H), 7.88 (dd, 1H), 8.41 (d, 1H), 8.78 (d, 1H); HPLC-MS : m/z = 305.1 (M+1); R_t = 2.99 min.

30

Example 526 (General procedure 20)

Methyl-phenyl-carbamic acid 4-(5-chloro-thiophen-2-yl)-phenyl ester

35 The title compound was prepared from methyl-phenyl-carbamic acid 4-iodo-phenyl ester and 5-chloro-2-thiopheneboronic acid. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane 1:9) (53%, pink crystals).

¹H NMR (300MHz; CDCl₃): δ 3.43 (br s, 3H), 6.87 (d, 1H), 7.00 (d, 1H), 7.13 (br d, 2H), 7.26-7.48 (m, 7H); HPLC-MS : m/z = 344.0 (M+1); R_t = 5.16 min.

Example 527 (General procedure 20)

5 Methyl-phenyl-carbamic acid 4'-benzylsulfamoyl-biphenyl-4-yl ester

The title compound was prepared from methyl-phenyl-carbamic acid 4-iodo-phenyl ester and 4-benzylsulfamoylbenzeneboronic acid. The crude product was purified by preparative HPLC (Gilson) (35%, pink crystals).

10 HPLC-MS : m/z = 473.0 (M+1); R_t = 4.80 min.

Example 528

Methyl-phenyl-carbamic acid 4-styryl-phenyl ester

15 Styrene (1.2 mmol), N-methyldicyclohexylamine (1.2 mmol), Pd₂(dba)₃ (0.03 mmol), Pd(P(*t*-Bu)₃)₂ (0.06 mmol) and methyl-phenyl-carbamic acid 4-iodo-phenyl ester (1.0 mmol) were added to a Schlenk tube under nitrogen. The Schlenk tube was evacuated and refilled with nitrogen five times. Then dioxane (2 mL) was added and the reaction mixture was stirred at 70 °C for 8 h. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) giving the title compound in 17% yield as colorless crystals.

20 HPLC-MS : m/z = 330.1 (M+1); R_t = 5.08 min.

Example 529

Methyl-phenyl-carbamic acid 4-phenylethynyl-phenyl ester

25 Phenylacetylene (1.2 mmol), diisopropylamine (1.2 mmol), CuI (0.03 mmol), Pd₂(dba)₃ (0.03 mmol), Pd(P(*t*-Bu)₃)₂ (0.06 mmol) and methyl-phenyl-carbamic acid 4-iodo-phenyl ester (1.0 mmol) were added to a Schlenk tube under nitrogen. The Schlenk tube was evacuated and refilled with nitrogen five times. Then dioxane (2 mL) was added and the reaction mixture

30 was stirred at 70 °C for 8 h. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) giving the title compound in 41% yield as brown oil.

HPLC-MS : m/z = 328.1 (M+1); R_t = 5.07 min.

Example 530

35 3-[4-(Methyl-phenyl-carbamoyloxy)-phenyl]-acrylic acid methyl ester

Methylacrylate (1.2 mmol), N-methyldicyclohexylamine (1.2 mmol), Pd₂(dba)₃ (0.03 mmol), Pd(P(*t*-Bu)₃)₂ (0.06 mmol) and methyl-phenyl-carbamic acid 4-iodo-phenyl ester (1.0 mmol) were added to a Schlenk tube under nitrogen. The Schlenk tube was evacuated and refilled with nitrogen five times. Then dioxane (2 mL) was added and the reaction mixture was stirred
5 at 70 °C for 8 h. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) giving the title compound in 70% yield as a yellow solid.
HPLC-MS : *m/z* = 312.1 (M+1); R_t = 4.19 min.

Example 531 (General procedure 8)

10 Methyl-phenyl-carbamic acid 5-phenylsulfanyl-pyrazol-1-yl ester

The title compound was prepared from 1-hydroxy-5-phenylsulfanylpirazole and N-methyl-N-phenylcarbamoyl chloride. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (80%, oil).

15 ¹H NMR (300MHz; CDCl₃): δ 3.36 (br s, 3H), 6.47 (d, 1H) 7.16-7.32 (m, 10H), 7.40 (d, 1H);
HPLC-MS : *m/z* = 326.0 (M+1); R_t = 4.42 min.

Example 532 (General procedure 21)

Methyl-phenyl-carbamic acid 4-(toluene-4-sulfonylamino)-phenyl ester

20

The title compound was prepared in 8% yield as a clear oil using toluenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : *m/z* = 397.1 (M+1); R_t = 4.13 min

25 **Example 533** (General procedure 21)

Methyl-phenyl-carbamic acid 4-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-phenyl ester

The title compound was prepared in 7% yield as an oil using 5-pyridin-2-yl-thiophene-2-sulfonyl chloride as the aryl sulfonyl chloride.

30 HPLC-MS : *m/z* = 466.1 (M+1); R_t = 4.23 min.

Example 534 (General procedure 21)

Methyl-phenyl-carbamic acid 4-(1-methyl-1H-imidazole-4-sulfonylamino)-phenyl ester

35 The title compound was prepared in 21% yield as crystals using 1-methyl-1H-imidazole-4-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 387.1 (M+1); R_t = 3.14 min.

Example 535 (General procedure 21)

Methyl-phenyl-carbamic acid 4-(2,5-dichloro-thiophene-3-sulfonylamino)-phenyl ester

5

The title compound was prepared in 2% yield as an oil using 2,5-dichloro-thiophene-3-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 458.6 (M+1); R_t = 4.38 min.

10 **Example 536** (General procedure 21)

Methyl-phenyl-carbamic acid 4-(5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonylamino)-phenyl ester

15

The title compound was prepared in 3% yield as an oil using 5-chloro-1,3-dimethyl-1H-pyrazole-4-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 435.1 (M+1); R_t = 3.74 min.

Example 537 (General procedure 21)

Methyl-phenyl-carbamic acid 4-(5-dimethylamino-naphthalene-1-sulfonylamino)-phenyl ester

20

The title compound was prepared in 14% yield as orange crystals using 5-dimethylamino-naphthalene-1-sulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 476.0 (M+1); R_t = 4.48 min.

25 **Example 538** (General procedure 21)

2-[4-(Methyl-phenyl-carbamoyloxy)-phenylsulfamoyl]-benzoic acid methyl ester

The title compound was prepared in 48% yield as a yellow oil using 2-chlorosulfonyl-benzoic acid methyl ester as the aryl sulfonyl chloride.

30

HPLC-MS : m/z = 441.1 (M+1); R_t = 4.19 min.

Example 539 (General procedure 21)

Methyl-phenyl-carbamic acid 4-(3,4-difluoro-benzenesulfonylamino)-phenyl ester

35

The title compound was prepared in 1% yield as a clear oil using 3,4-difluoro-benzenesulfonyl chloride as the aryl sulfonyl chloride.

HPLC-MS : m/z = 419.1 (M+1); R_t = 4.23 min.

Example 540 (General procedure 22 and 1)

Methyl-phenyl-carbamic acid 4-pyridin-2-ylmethyl-phenyl ester

5

4-Pyridin-2-ylmethyl-phenol was prepared following general procedure PVe3 using pyridine-2-carboxaldehyde. Subsequent carbamoylation using general procedure 1 (CH_2Cl_2 was used as solvent) produced the crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (86%, oil).

10 ^1H NMR (400MHz; CDCl_3): δ 3.40 (br s, 3H), 4.14 (s, 2H), 7.04-7.12 (m, 4H), 7.21-7.26 (m, 3H), 7.32-7.40 (m, 4H), 7.55 (t, 1H), 8.52 (d, 1H).

Example 541 (General procedure 22 and 1)

Methyl-phenyl-carbamic acid 4-pyridin-3-ylmethyl-phenyl ester

15

4-Pyridin-3-ylmethyl-phenol was prepared following general procedure PVe3 using pyridine-3-carboxaldehyde. Subsequent carbamoylation using general procedure 1 (CH_2Cl_2 was used as solvent) produced the crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (50%, oil).

20 ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 3.96 (s, 2H), 7.03-7.42 (m, 11H), 8.46 (d, 1H), 8.49 (d, 1H).

Example 542 (General procedure 22 and 1)

Methyl-phenyl-carbamic acid 4-(4-trifluoromethyl-benzyl)-phenyl ester

25

4-(4-Trifluoromethyl-benzyl)-phenol was prepared following general procedure PVe3 using 4-trifluoromethylbenzaldehyde. Subsequent carbamoylation using general procedure 1 (CH_2Cl_2 was used as solvent) produced the crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (92%, oil).

30 ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 4.00 (s, 2H), 7.06 (br s, 2H), 7.13 (d, 2H), 7.24-7.40 (m, 7H), 7.52 (d, 2H).

Example 543 (General procedure 22 and 1)

Methyl-phenyl-carbamic acid 4-thiophen-3-ylmethyl-phenyl ester

35

4-Thiophen-3-ylmethyl-phenol was prepared following general procedure PVe3 using thio-

phene-3-carboxaldehyde. Subsequent carbamoylation using general procedure 1 (CH_2Cl_2 was used as solvent) produced the crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (83%, oil).

^1H NMR (400MHz; CDCl_3): δ 3.42 (bs, 3H), 3.95 (s, 2H), 6.86-7.39 (m, 12H).

5

Example 544 (General procedure 22 and 1)

Methyl-phenyl-carbamic acid 4-thiophen-2-ylmethyl-phenyl ester

4-Thiophen-2-ylmethyl-phenol phenol was prepared following general procedure PVe3 using
10 thiophene-3-carboxaldehyde. Subsequent carbamoylation using general procedure 1 (CH_2Cl_2 was used as solvent) produced the crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane) (83%, oil).

^1H NMR (400MHz; CDCl_3): δ 3.40 (br s, 3H), 4.13 (s, 2H), 6.78 (dd, 1H), 6.90 (dd, 1H), 7.04 (br d, 2H), 7.13 (dd, 1H), 7.20-7.39 (m, 7H). HPLC-MS : m/z = 324.1 (M+1); R_t = 4.82 min.

15

Example 545

4-Hydroxy-piperidine-1-carboxylic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester

N-Boc protected tyramin (10 mmol), triethylamine (10 mmol) and 3-[4-(tert-Butyl-dimethyl-
20 silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium; iodide (10 mmol) in MeCN (25 mL) was stirred at room temperature for 16 hours. Acetonitrile was removed by evaporation and the crude product was purified by flash chromatography (Quad flash 40, EtOAc-heptane 1:2) providing 72% 4-tert-butylsilanyloxy-piperidine-1-carboxylic acid 4-(2-tert-butoxycarbonylamino-ethyl)-phenyl ester. This was deprotected by stirring with a 3.2 M solution of HCl in Et_2O (50 mL) for 3 h at rt and subsequently washed with ether to give 91% of 4-
25 (4-hydroxy-piperidine-1-carboxyloxy)-phenyl-ammonium; chloride as a solid. This compound was N-tosylated as described for methyl-phenyl-carbamic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester to give the title compound in 26% yield as yellow crystals after purification by flash chromatography (Quad flash 12, CH_2Cl_2 -MeOH 95:5).

30 HPLC-MS : m/z = 391.0 (M+1); R_t = 3.04 min.

Example 546

4-Hydroxy-piperidine-1-carboxylic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-phenyl

35

4-(4-Hydroxy-piperidine-1-carboxyloxy)-phenyl-ammonium; chloride (see above) was N-

sulfonylated as described for methyl-phenyl-carbamic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-phenyl ester to give the title compound in 59% yield as an oil after purification by preparative HPLC (Gilson).

HPLC-MS : m/z = 488.0 (M+1); R_t = 3.10 min.

5

Example 547

4-Hydroxy-piperidine-1-carboxylic acid 4-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-phenyl ester

- 10 N-Boc protected 4-aminophenol (10 mmol), triethylamine (10 mmol) and 3-[4-(tert-Butyl-dimethyl-silanyloxy)-piperidine-1-carbonyl]-1-methyl-3H-imidazol-1-ium; iodide (10 mmol) in MeCN (25 mL) was stirred at room temperature for 16 hours. Acetonitrile was removed by evaporation and the crude product was purified by flash chromatography (Quad flash 40, EtOAc-heptane 1:2) providing 64% 4-tert-butylsilanyloxy-piperidine-1-carboxylic acid 4-tert-
- 15 butoxycarbonylamino-phenyl ester. This was deprotected by stirring with a 3.2 M solution of HCl in Et₂O (50 mL) for 3 h at rt and subsequently washed with ether to give 91% of 4-(4-Hydroxy-piperidine-1-carboxyloxy)-phenyl-ammonium; chloride as a hygroscopic solid. This compound was N-sulfonylated as described for methyl-phenyl-carbamic acid 4-[2-(5-pyridin-2-yl-thiophene-2-sulfonylamino)-ethyl]-phenyl ester to give the title compound in 1% yield as
- 20 crystals after purification by preparative HPLC (Gilson).
- HPLC-MS : m/z = 482.8 (M+1); R_t = 1.01 min.

Example 548

Methyl-phenyl-carbamic acid 4-[2-(4-amino-benzenesulfonylamino)-ethyl]-phenyl ester

25

Methyl-phenyl-carbamic acid 4-[2-(4-nitro-benzenesulfonylamino)-ethyl]-phenyl ester, 5% palladium on carbon, and ethanol were stirred under hydrogen at 1 bar and rt for 16 h. Filtration and removal of ethanol produced the title compound in 94% yield as an oil.

HPLC-MS : m/z = 426.1 (M+1); R_t = 3.61 min.

30

Example 549

Methyl-phenyl-carbamic acid 4-{2-[(pyridine-3-carbonyl)-amino]-ethyl}-phenyl ester

- A solution of 3-pyridinecarboxylic acid (0.3 mmol), EDAC (0.36 mmol) and triethylamine (0.36 mmol) in CH₂Cl₂ (10 mL) was added N-hydroxybenzotriazole and N-methyl-N-phenyl-
- 35 carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol). The mixture was stir-

stirred 16 h at rt and purified by preparative HPLC (Gilson) to give the title compound in 15% yield as an oil.

HPLC-MS : m/z = 376.1 (M+1); R_t = 3.01 min.

5 Example 550

Methyl-phenyl-carbamic acid 4-[2-(2-dimethylamino-acetyl-amino)-ethyl]-phenyl ester

A solution of N,N-dimethylglycine, HCl (0.3 mmol), EDAC (0.36 mmol) and triethylamine (1.0 mmol) in CH_2Cl_2 (10 mL) was added N-hydroxybenzotriazole and N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol). The mixture was stirred 16 h at rt and purified by preparative HPLC (Gilson) to give the title compound in 66% yield as an oil.

HPLC-MS : m/z = 356.4 (M+1); R_t = 2.09 min.

15 Example 551 (General procedure 23)

Methyl-phenyl-carbamic acid 2-(toluene-4-sulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl ester

The title compound was prepared using methyl-phenyl-carbamic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester as the sulfonamide. The crude product was purified by flash chromatography (Quad flash 12, EtOAc-heptane 2:3) (71%, yellow oil).

HPLC-MS : m/z = 437.4 (M+1); R_t = 4.43 min.

Example 552

Methyl-phenyl-carbamic acid 4-[4-(2-pyrrolidin-1-yl-ethoxy)-benzyl]-phenyl ester

25

To a stirred solution of 1-[2-(4-bromo-phenoxy)-ethyl]-pyrrolidine (2.89 g, 10.7 mmol) in THF (30 mL) was added dropwise 1.6 M solution in hexanes n-BuLi (7 mL, 10.2 mmol) over a 5-min period at -78°C. The mixture was stirred at -78°C for 15 min before 4-trimethylsilyloxybenzaldehyde (2.08 g, 10.7 mmol) was added. The mixture was allowed to warm to rt during 20 min and quenched with water. Extraction with CH_2Cl_2 , drying (MgSO_4), filtration and evaporation provided the crude diarylmethanol which was dissolved in CH_2Cl_2 (30 mL) and stirred with triethylsilane (4 mL) and TFA (5 mL) for 16 h at rt. Evaporation gave 3.0 g (84%) 4-[4-(2-pyrrolidin-1-yl-ethoxy)-benzyl]-phenol. This was carbamoylated using general procedure 1 (CH_2Cl_2 was used as solvent) to give the title product after purification by preparative HPLC (Gilson) (35%, oil).

35

HPLC-MS : m/z = 431.5 (M+1); R_t = 2.93 min.

Example 553 (General procedure 24)

4-Hydroxy-piperidine-1-carboxylic acid 4-pyridin-2-ylmethyl-phenyl ester

- 5 The title compound was prepared as its hydrochloride using 4-pyridin-2-ylmethyl-phenol as the phenol.

¹H NMR (300MHz; D₂O): δ 1.50 (br s, 2H), 1.90 (d, 2H), 3.04-3.27 (m, 2H), 3.78-4.08 (m, 3H), 4.41 (s, 2H), 7.07 (d, 2H), 7.30 (d, 2H), 7.79-7.85 (m, 2H), 8.40 (dt, 1H), 8.55 (d, 1H).

10 **Example 554** (General procedure 24)

4-Hydroxy-piperidine-1-carboxylic acid 4-pyridin-3-ylmethyl-phenyl ester

The title compound was prepared as its hydrochloride using 4-pyridin-3-ylmethyl-phenol as the phenol.

- 15 ¹H NMR (300MHz; D₂O): δ 1.48 (br s, 2H), 1.90 (d, 2H), 3.03-3.26 (m, 2H), 3.80-4.05 (m, 3H), 4.18 (s, 2H), 7.03 (d, 2H), 7.25 (d, 2H), 7.40 (dd, 1H), 8.40 (d, 1H), 8.54-8.57 (m, 2H).

Example 555 (General procedure 24)

4-Hydroxy-piperidine-1-carboxylic acid 4-(4-trifluoromethyl-benzyl)-phenyl ester

20

The title compound was prepared using 4-(4-trifluoromethyl-benzyl)-phenol as the phenol.

¹H NMR (300MHz; D₂O): δ 1.43-1.55 (m, 2H), 1.76-1.88 (m, 2H), 3.10-3.29 (m, 2H), 3.79-3.94 (m, 5H), 6.97 (d, 2H), 7.08 (d, 2H), 7.20 (d, 1H), 7.45 (d, 2H).

25 **Example 556** (General procedure 23)

Methyl-phenyl-carbamic acid 2-[4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfonyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl ester

- 30 The title compound was prepared using methyl-phenyl-carbamic acid 4-{2-[4-(2-pyrrolidin-1-yl-ethoxy)-benzenesulfonylamino]-ethyl}-phenyl ester as the sulfonamide. The crude product was purified by preparative HPLC (Gilson) and isolated as its TFA salt (45%, foam).

HPLC-MS : m/z = 536.2 (M+1); R_t = 3.10 min.

Example 557

- 35 Methyl-phenyl-carbamic acid 4-{2-[(1-methyl-piperidine-4-carbonyl)-amino]-ethyl}-phenyl ester

A solution of 1-methylpiperidine-4-carboxylic acid (0.3 mmol), EDAC (0.36 mmol) and triethylamine (1.0 mmol) in CH₂Cl₂ (10 mL) was added N-hydroxybenzoetriaizole and N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol). The mixture
5 was stirred 16 h at rt and purified by preparative HPLC (Gilson) to give the title compound in 5% yield as an oil.

HPLC-MS : m/z = 396.4 (M+1); R_t = 2.03 min.

Example 558 (General procedure 23)

10 Methyl-phenyl-carbamic acid 2-(3,4-difluoro-benzenesulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl ester

The title compound was prepared using methyl-phenyl-carbamic acid 4-[2-(3,4-difluoro-benzenesulfonylamino)-ethyl]-phenyl ester as the sulfonamide. The crude product was purified
15 by preparative HPLC (Gilson) (24%, oil).

HPLC-MS : m/z = 481.0 (M+23); R_t = 4.73 min.

Example 559 (General procedure 23)

20 Methyl-phenyl-carbamic acid 1-methyl-2-(toluene-4-sulfonyl)-1,2,3,4-tetrahydro-isoquinolin-7-yl ester

The title compound was prepared using methyl-phenyl-carbamic acid 4-[2-(toluene-4-sulfonylamino)-ethyl]-phenyl ester as the sulfonamide and acetaldehyde in stead of formaldehyde. The crude product was purified by preparative HPLC (Gilson) (22%, brown oil).

25 HPLC-MS : m/z = 451.5 (M+1); R_t = 4.43 min.

Example 560 (General procedure 23)

Methyl-phenyl-carbamic acid 2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydro-isoquinolin

30

The title compound was prepared using methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonylamino]-ethyl}-phenyl ester as the sulfonamide. The crude product was purified by preparative HPLC (Gilson) and isolated as its TFA salt (79%, crystals).

35 HPLC-MS : m/z = 521.5 (M+1); R_t = 2.65 min.

Example 561 (General procedure 23)

Methyl-phenyl-carbamic acid 1-methyl-2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl ester

- 5 The title compound was prepared using methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-yl)-benzenesulfonylamino]-ethyl}-phenyl ester as the sulfonamide and acetaldehyde in stead of formaldehyde. The crude product was purified by preparative HPLC (Gilson) (12%, brown oil).

HPLC-MS : m/z = 535.4 (M+1); R_t = 2.69 min.

10

Example 562

3,3-Dimethyl-4-{2-[4-(methyl-phenyl-carbamoyloxy)-phenyl]-ethylcarbamoyl}-butyric acid

- A solution of 3,3-dimethylglutaric anhydride (0.3 mmol), diisopropylethylamine (0.30 mmol) and N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol) in CH_2Cl_2 (3 mL) was stirred 1 h at rt. The mixture was washed with water and brine, dried and evaporated to give the title compound in 95% yield as an oil.

HPLC-MS : m/z = 413.2 (M+1); R_t = 3.20 min.

20 **Example 563**

Methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-yl)-benzoylamino]-ethyl}-phenyl ester

- A solution of 4-(4-methylpiperazino)benzoic acid (0.3 mmol), EDAC (0.36 mmol) and triethylamine (1.0 mmol) in CH_2Cl_2 (10 mL) was added N-hydroxybenzoetriaizole and N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol). The mixture was stirred 16 h at rt and purified by preparative HPLC (Gilson) to give the title compound in 54% yield as its crystalline hydrochloride after treatment with HCl in diethyl ether.

HPLC-MS : m/z = 473.2 (M+1); R_t = 2.22 min.

30

Example 564

Methyl-phenyl-carbamic acid 4-{2-[4-(4-methyl-piperazin-1-ylmethyl)-benzoylamino]-ethyl}-phenyl ester

- 35 A solution of 4-(4-methylpiperazinyl)methyl benzoic acid (0.3 mmol), EDAC (0.36 mmol) and triethylamine (1.0 mmol) in CH_2Cl_2 (10 mL) was added N-hydroxybenzoetriaizole and N-

methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol). The mixture was stirred 16 h at rt and purified by preparative HPLC (Gilson) to give the title compound in 25% yield as its crystalline hydrochloride after treatment with HCl in diethyl ether.. HPLC-MS : m/z = 487.3 (M+1); R_t = 2.13 min.

5

Example 565

Methyl-phenyl-carbamic acid 4-[2-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-ethyl]-phenyl ester

A solution of 3,3-dimethylglutaric anhydride (0.3 mmol), diisopropylethylamine (0.30 mmol) and N-methyl-N-phenyl-carbamic acid 4-(2-amino-ethyl)phenyl ester as its TFA salt (0.3 mmol) in CH_2Cl_2 (3 mL) was stirred 1 h at rt. Thionylchloride (3 mmol) was added and the mixture was stirred for 2 h at rt. Addition of ethanol (5 mL) followed by evaporation to dryness gave a crude product which was purified by flash chromatography (Quad flash 12, EtOAc-heptane 1:1). This gave the title compound in 23% yield as crystals. Furthermore 3,3-dimethyl-4-{2-[4-(methyl-phenyl-carbamoyloxy)-phenyl]-ethylcarbamoyl}-butyric acid ethyl ester could be isolated in 27% yield as an oil (see characterization below)

15

^1H NMR (400MHz; CDCl_3): δ 1.04 (s, 6H), 2.47 (s, 4H), 2.79 (t, 2H), 3.42 (br s, 3H), 3.97 (t, 2H), 7.03 (br d, 2H), 7.21-7.28 (m, 3H), 7.33-7.41 (m, 4H).

HPLC-MS : m/z = 395.2 (M+1); R_t = 4.23 min.

20

Example 566

3,3-Dimethyl-4-{2-[4-(methyl-phenyl-carbamoyloxy)-phenyl]-ethylcarbamoyl}-butyric acid ethyl ester

For experimental details, see preparation of methyl-phenyl-carbamic acid 4-[2-(4,4-dimethyl-2,6-dioxo-piperidin-1-yl)-ethyl]-phenyl ester.

^1H NMR (400MHz; CDCl_3): δ 1.05 (s, 6H), 1.24 (t, 3H), 2.19 (s, 2H), 2.22 (s, 2H), 2.80 (t, 2H), 3.42 (br s, 3H), 3.50 (q, 2H), 4.1 (q, 2H), 6.53 (t, 1H), 7.03 (br d, 2H), 7.18 (d, 2H), 7.26 (t, 1H), 7.33-7.41 (m, 4H).

HPLC-MS : m/z = 441.2 (M+1); R_t = 4.11 min.

30

Example 567

Methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester

To a solution of 4-hydroxymethylphenol (10 mmol) and 4-diazabicyclo[2.2.2]octane (DABCO) (10 mmol) in CH_2Cl_2 (30 mL) was added N-methyl-N-phenylcarbamoyl chloride (10 mmol).

35

The reaction mixture was stirred for 16 hours at rt, added CH_2Cl_2 (20 mL) and washed with 1M aqueous HCl and brine. The organic phase was dried (MgSO_4) and evaporated to give the crude product which was purified by FC (Quad flash 40 EtOAc-Heptane 1:1) to give 2.18 g (85%) of the title compound as an oil.

- 5 ^1H NMR (400MHz; CDCl_3): δ 1.70 (br s, 1H), 3.40 (br s, 3H), 4.66 (s, 2H), 7.08 (br d, 2H), 7.25-7.42 (m, 7H); HPLC-MS : m/z = 258.1 (M+1); R_t = 2.99 min.

Example 568

Methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester

10

To a solution of 4-(2-Hydroxyethyl)phenol (10 mmol) and 4-diazabicyclo[2.2.2]octane (DABCO) (10 mmol) in CH_2Cl_2 (30 mL) was added N-methyl-N-phenylcarbonyl chloride (10 mmol). The reaction mixture was stirred for 16 hours at rt, added CH_2Cl_2 (20 mL), and washed with 1M aqueous HCl and brine. The organic phase was dried (MgSO_4) and evaporated to give the 2.69 g (99%) of the title compound as crystals.

15

^1H NMR (400MHz; CDCl_3): δ 1.52 (br s, 1H), 2.84 (t, 2H), 3.41 (br s, 3H), 3.81 (t, 2H), 7.04 (br d, 2H), 7.20 (d, 2H), 7.22-7.42 (m, 5H); HPLC-MS : m/z = 272.1 (M+1); R_t = 3.17 min.

Example 569 (General procedure 1)

20

Methyl-phenyl-carbamic acid 4-(4-dimethylamino-pyridin-2-ylmethyl)-phenyl ester

25

A solution of 2-(dimethylamino)ethanol (32 mmol) in hexane (120 mL) was cooled to -5°C and n-BuLi (64 mmol) was added. After 30 min 4-(dimethylamine=pyridine) (16 mmol) was added and the red-orange mixture was stirred for further 1 h. The solution was cooled to -78°C and 4-(trimethylsilyloxy)benzaldehyde (40 mmol) dissolved in hexane (80 mL) was added and the suspension was allowed to warm to rt over 20 min. The reaction mixture was quenched with water, and the aqueous phase was washed with CH_2Cl_2 . The aqueous phase was evaporated to dryness, added NaI (96 mmol) and dissolved in MeCN (160 mL). Addition of trimethylsilylchloride (96 mmol) and stirring at rt for 16h. The purple reaction mixture was evaporated to dryness and treated with an aqueous solution of Na_2SO_3 and pH was adjusted to 8. Extraction with CH_2Cl_2 gave after evaporation 560 mg (15%) slightly impure 4-(4-dimethylamino-pyridin-2-ylmethyl)phenol as yellow crystals. This was carbamoylated using general procedure 1 (CH_2Cl_2 was used as solvent) to give the title product as its hydrochloride after purification by preparative HPLC (Gilson) and treatment with HCl in Et_2O (48%, light yellow crystals).

35

HPLC-MS : m/z = 362.2 (M+1); R_t = 2.27 min.

Example 570 (General procedure 25)

Methyl-phenyl-carbamic acid 4-(4-imidazol-1-yl-phenoxy-methyl)-phenyl ester

5

The title compound was prepared in 53% yield as light yellow crystals using and 4-imidazol-1-yl-phenol. ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 5.08 (s, 2H), 7.03 (d, 2H), 7.12-7.42 (m, 15H), 7.77 (s, 1H); HPLC-MS : m/z = 400.1 (M+1); R_t = 2.62 min.

10 **Example 571** (General procedure 25)

Methyl-phenyl-carbamic acid 4-[4-(2-dimethylamino-ethyl)-phenoxy-methyl]-phenyl ester

15 The title compound was prepared in 52% yield as colorless crystals using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 4-(2-dimethylaminoethyl)-phenol. ^1H NMR (400MHz; CDCl_3): δ 2.85 (s, 6H), 2.98-3.02 (m, 2H), 3.17-3.21 (m, 2H), 3.42 (br s, 3H), 5.02 (s, 2H), 6.90 (d, 2H), 7.12 (d, 2H), 7.25-7.43 (m, 7H); HPLC-MS : m/z = 405.2 (M+1); R_t = 2.91 min.

Example 572 (General procedure 25)

20 Methyl-phenyl-carbamic acid 4-(pyrazol-1-yloxy-methyl)-phenyl ester

The title compound was prepared in 79% yield as an oil using and 1-hydroxypyrazole. ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 5.26 (s, 2H), 6.03 (t, 1H), 6.95 (dd, 1H), 7.11 (br s, 2H), 7.25-7.42 (m, 8H); HPLC-MS : m/z = 324.1 (M+1); R_t = 3.58 min.

25

Example 573 (General procedure 25)

Methyl-phenyl-carbamic acid 4-(imidazol-1-yloxy-methyl)-phenyl ester

30 The title compound was prepared as its TFA salt in 87% yield as a solid using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 1-hydroxyimidazole, hydrochloride. ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 5.21 (s, 2H), 6.95 (s, 1H), 7.16-7.45 (m, 10H), 8.28 (br s, 1H); HPLC-MS : m/z = 324.1 (M+1); R_t = 1.92 min.

Example 574 (General procedure 25)

35 Methyl-phenyl-carbamic acid 4-(2-oxo-2H-pyridin-1-ylmethyl)-phenyl ester

The title compound was prepared in 29% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 2-hydroxypyridine. In addition 50% of the isomeric methyl-phenyl-carbamic acid 4-[2-(pyridin-2-yloxy)-ethyl]-phenyl ester was isolated, see characterization below. ¹H NMR (400MHz; CDCl₃): δ 3.05 (t, 2H), 3.41 (br s, 3H), 4.14 (s, 2H), 6.08 (t, 1H), 6.66 (d, 1H), 6.90 (dd, 1H), 7.02 (br s, 2H), 7.10 (d, 2H), 7.25-7.42 (m, 6H); HPLC-MS : *m/z* = 349.2 (M+1); *R*_t = 3.04 min.

Example 575 (General procedure 25)

Methyl-phenyl-carbamic acid 4-[2-(pyridin-2-yloxy)-ethyl]-phenyl ester

The title compound was prepared in 50% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 2-hydroxypyridine. In addition 29% of the isomeric methyl-phenyl-carbamic acid 4-(2-oxo-2H-pyridin-1-ylmethyl)-phenyl ester was isolated, see characterization above. ¹H NMR (400MHz; CDCl₃): δ 3.08 (t, 2H), 3.41 (br s, 3H), 4.49 (s, 2H), 6.73 (d, 1H), 6.88 (dd, 1H), 7.05 (br d, 2H), 7.22-7.42 (m, 7H), 7.61 (dt, 1H), 8.18 (dd, 1H); HPLC-MS : *m/z* = 349.2 (M+1); *R*_t = 3.97 min.

Example 576 (General procedure 25)

Methyl-phenyl-carbamic acid 4-[2-(4-imidazol-1-yl-phenoxy)-ethyl]-phenyl ester

The title compound was prepared as its TFA salt in 62% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 4-imidazol-1-yl-phenol.

¹H NMR (400MHz; CDCl₃): δ 3.11 (t, 2H), 3.42 (br s, 3H), 4.22 (s, 2H), 7.00-7.10 (m, 5H), 7.25-7.42 (m, 9H), 7.52 (s, 1H), 8.81 (s, 1H); HPLC-MS : *m/z* = 414.2 (M+1); *R*_t = 2.73 min.

Example 577 (General procedure 25)

Methyl-phenyl-carbamic acid 4-[2-[4-(2-dimethylamino-ethyl)-phenoxy]-ethyl]-phenyl ester

The title compound was prepared as its TFA salt in 92% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 4-(2-dimethylaminoethyl)-phenol.

HPLC-MS : *m/z* = 419.2 (M+1); *R*_t = 2.77 min.

Example 578 (General procedure 25)

Methyl-phenyl-carbamic acid 4-[2-(pyrazol-1-yloxy)-ethyl]-phenyl ester

The title compound was prepared in 62% yield as an oil using methyl-phenyl-carbamic acid

4-(2-hydroxy-ethyl)-phenyl ester and 1-hydroxypyrazole.

^1H NMR (400MHz; CDCl_3): δ 3.02 (t, 2H), 3.42 (br s, 3H), 4.50 (t, 2H), 6.16 (t, 1H), 7.05 (br d, 2H), 7.20-7.41 (m, 9H); HPLC-MS : m/z = 338.2 (M+1); R_t = 3.74 min.

5 **Example 579** (General procedure 25)

Methyl-phenyl-carbamic acid 4-[2-(imidazol-1-yloxy)-ethyl]-phenyl ester

The title compound was prepared as its TFA salt in 79% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 1-hydroxyimidazole hydrochlorid.

10 ^1H NMR (400MHz; CDCl_3): δ 3.06 (t, 2H), 3.42 (br s, 3H), 4.52 (t, 2H), 7.08-7.42 (m, 11H), 8.53 (s, 1H); HPLC-MS : m/z = 338.2 (M+1); R_t = 2.18 min.

Example 580 (General procedure 1)

Methyl-phenyl-carbamic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester

15

To a stirred solution of 2-bromo-5-methylpyridine (3.45 g, 20 mmol) in THF (10 mL) was added dropwise 1.6 M solution in hexanes n-BuLi (12 mL, 19.2 mmol) over a 10-min period at -78°C . The mixture was stirred at -78°C for 2 min before 4-trimethylsilyloxybenzaldehyde (4.2 g, 21.6 mmol) dissolved in THF (10 mL) was added. The mixture was allowed to warm to -40°C and quenched with water. The pH of the aqueous phase was adjusted to 7 which causes precipitation of 4-[hydroxy-(5-methyl-pyridin-2-yl)-methyl]phenol. This was isolated by filtration to give 1.13 g (27%) of 4-[hydroxy-(5-methyl-pyridin-2-yl)-methyl]phenol as crystals. This intermediate was dissolved in CH_2Cl_2 (15 mL) and stirred with triethylsilane (4 mL) and TFA (5 mL) for 16 h at 50°C . Evaporation gave 99% of 4-(5-methyl-pyridin-2-ylmethyl)phenol as a hygroscopic solid. This was carbamoylated using general procedure 1 (CH_2Cl_2 was used as solvent and N-phenyl-N-methyl carbamoyl chlorid) to give 99% of the title product as its crystalline hydrochloride after purification by flash chromatography (Quad flash 12, EtOAc-heptane) and treatment with HCl in Et_2O .

20 ^1H NMR (400MHz; CDCl_3): δ 2.50 (s, 3H), 3.41 (br s, 3H), 4.57 (s, 2H), 7.10 (br s, 2H), 7.25-7.42 (m, 8H), 8.00 (dd, 1H), 8.46 (br s, 1H); HPLC-MS : m/z = 333.1 (M+1); R_t = 2.60 min.

30

Example 581 (General procedure 24)

4-Hydroxy-piperidine-1-carboxylic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester

35 The title compound was prepared as its hydrochloride using of 4-(5-methyl-pyridin-2-

ylmethyl)phenol as the phenol.

¹H NMR (300MHz; CDCl₃): δ 1.57-1.67 (m, 3H), 1.95 (d, 2H), 2.51 (s, 3H), 3.21-3.40 (m, 2H), 3.90-4.05 (m, 3H), 4.58 (s, 2H), 7.11 (d, 2H), 7.38 (d, 2H), 7.45 (d, 1H), 8.02 (d, 1H), 8.48 (br s, 1H).

5

Example 582 (General procedure 25)

Methyl-phenyl-carbamic acid 4-(4-oxo-4H-pyridin-1-ylmethyl)-phenyl ester

The title compound was prepared in 33% yield as colorless crystals using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 4-hydroxypyridine. ¹H NMR (400MHz; CDCl₃): δ 3.43 (br s, 3H), 4.92 (s, 2H), 6.43 (d, 2H), 7.17 (br s, 4H), 7.22-7.42 (m, 9H); HPLC-MS : m/z = 335.0 (M+1); R_t = 2.55 min.

10

Example 583 (General procedure 25)

15 Methyl-phenyl-carbamic acid 4-[2-(pyridin-3-yloxy)-ethyl]-phenyl ester

The title compound slightly contaminated with tributylphosphine oxide was prepared in 80% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 3-hydroxypyridine.

20 ¹H NMR (400MHz; CDCl₃): δ 3.11 (t, 2H), 3.42 (br s, 3H), 4.23 (t, 2H), 7.07 (br d, 2H), 7.24-7.41 (m, 9H), 8.29 (br s, 1H), 8.40 (br s, 1H); HPLC-MS : m/z = 349.2 (M+1); R_t = 2.87 min.

Example 584 (General procedure 25)

Methyl-phenyl-carbamic acid 4-(2-oxo-2H-pyridin-1-ylmethyl)-phenyl ester

25

The title compound was prepared in 66% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 2-hydroxypyridine. ¹H NMR (400MHz; CDCl₃): δ 3.42 (br s, 3H), 5.12 (s, 2H), 6.13 (t, 1H), 6.60 (d, 1H), 7.10 (br s, 2H), 7.22-7.41 (m, 9H); HPLC-MS : m/z = 335.2 (M+1); R_t = 2.97 min.

30

Example 585 (General procedure 25)

Methyl-phenyl-carbamic acid 4-(pyridin-3-yloxymethyl)-phenyl ester

The title compound was prepared in 43% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 3-hydroxypyridine. ¹H NMR (400MHz; CDCl₃): δ 3.44 (br

35

s, 3H), 5.12 (s, 2H), 7.17 (br s, 2H), 7.27-7.46 (m, 9H), 8.29 (br s, 1H), 8.49 (br s, 1H); HPLC-MS : m/z = 335.0 (M+1); R_t = 2.74 min.

Example 586 (General procedure 26)

5 Methyl-phenyl-carbamic acid 4-[2-(2,5-dioxo-pyrrolidin-1-yl)-ethyl]-phenyl ester

The title compound was prepared in 85% yield as crystals using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and succinimide. ^1H NMR (400MHz; CDCl_3): δ 2.65 (s, 4H), 2.86 (t, 2H), 3.41 (br s, 3H), 3.72 (t, 2H), 7.04 (br d, 2H), 7.19 (d, 2H), 7.26-7.42 (m, 5H);

10 HPLC-MS : m/z = 353.2 (M+1); R_t = 3.17 min.

Example 587 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(1,3-dioxo-1,3-dihydro-pyrrolo[3,4-]pyridin-2-yl)-ethyl]-phenyl ester

15

The title compound was prepared in 58% yield as crystals using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 3,4-pyridinedicarboximide. ^1H NMR (400MHz; CDCl_3): δ 2.98 (t, 2H), 3.40 (br s, 3H), 3.93 (t, 2H), 7.05 (br d, 2H), 7.20-7.42 (m, 7H), 7.73 (dd, 1H), 9.06 (d, 1H), 9.12 (d, 1H); HPLC-MS : m/z = 402.1 (M+1); R_t = 3.56 min.

20

Example 588 (General procedure 26)

Methyl-phenyl-carbamic acid 4-(1-methyl-1H-imidazol-2-ylsulfanylmethyl)-phenyl ester

The title compound was prepared as its TFA salt in 22% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 2-mercapto-1-methylimidazole. ^1H NMR (400MHz; CDCl_3): δ 3.34 (s, 3H), 3.40 (br s, 3H), 4.30 (s, 2H), 6.98 (br s, 2H), 7.08-7.11 (m, 3H), 7.27-7.42 (m, 5H), 7.48 (d, 1H); HPLC-MS : m/z = 354.1 (M+1); R_t = 2.12 min.

25

Example 589 (General procedure 26)

30 Methyl-phenyl-carbamic acid 4-tetrazol-1-ylmethyl-phenyl ester

The title compound was prepared in 6% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and tetrazole. ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 5.57 (s, 2H), 7.17 (br s, 2H), 7.26-7.42 (m, 7H), 8.52 (s, 1H); HPLC-MS : m/z = 332.0 (M+23); R_t = 3.24 min.

35

Example 590 (General procedure 26)

Methyl-phenyl-carbamic acid 4-(2,5-dioxo-pyrrolidin-1-ylmethyl)-phenyl ester

- 5 The title compound was prepared in 57% yield as beige crystals using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and succinimide. ^1H NMR (400MHz; CDCl_3): δ 2.68 (s, 4H), 3.41 (br s, 3H), 4.60 (s, 2H), 7.04 (br d, 2H), 7.23-7.41 (m, 7H); HPLC-MS : m/z = 339.1 (M+1); R_t = 3.40 min.

10 **Example 590** (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(2-thioxo-2H-pyridin-1-yl)-ethyl]-phenyl ester

- The title compound was prepared in 25% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 2-mercaptopyridine. HPLC-MS : m/z = 365.2 (M+1); R_t = 4.08 min.
- 15

Example 591 (General procedure 26)

Methyl-phenyl-carbamic acid 4-(1,3-dioxo-1,3-dihydro-pyrrolo[3,4]pyridin-2-ylmethyl)-phenyl ester

20

The title compound was prepared in 21% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 3,4-pyridinedicarboximide. ^1H NMR (400MHz; CDCl_3): δ 3.40 (br s, 3H), 4.82 (s, 2H), 7.06 (d, 2H), 7.22-7.44 (m, 8H), 7.73 (d, 1H), 9.04 (d, 1H), 9.12 (s, 1H); HPLC-MS : m/z = 388.0 (M+1); R_t = 3.80 min.

25

Example 592 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[1,2,4]triazol-1-ylmethyl-phenyl ester

- The title compound was prepared as its TFA salt in 27% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 1,2,4-triazole. ^1H NMR (400MHz; CDCl_3): δ 3.42 (br s, 3H), 5.34 (s, 2H), 7.15 (br d, 2H), 7.26-7.43 (m, 7H), 8.10 (s, 1H), 8.38 (s, 1H); HPLC-MS : m/z = 309.1 (M+1); R_t = 2.74 min.
- 30

Example 593 (General procedure 26)

- 35 Methyl-phenyl-carbamic acid 4-(2-thioxo-2H-pyridin-1-ylmethyl)-phenyl ester

The title compound was prepared in 14% yield as an oil using methyl-phenyl-carbamic acid 4-hydroxymethyl-phenyl ester and 2-mercaptopyridine. HPLC-MS : m/z = 351.1 (M+1); R_t = 3.95 min.

5

Example 594 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-phenyl ester

10 The title compound was prepared as its TFA salt in 37% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 2-mercapto-1-methylimidazole. NMR (400MHz; $CDCl_3$): δ 3.01 (t, 2H), 3.40 (br s, 3H), 3.47 (s, 1H), 3.64 (t, 2H), 6.92 (br d, 2H), 6.98 (s, 1H), 7.08 (d, 2H), 7.26-7.43 (m, 6H); HPLC-MS : m/z = 368.2 (M+1); R_t = 2.30 min.

Example 595 (General procedure 26)

15 Methyl-phenyl-carbamic acid 4-(2-tetrazol-1-yl-ethyl)-phenyl ester

The title compound was prepared in 10% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and tetrazole. NMR (400MHz; $CDCl_3$): δ 3.19 (t, 2H), 3.40 (br s, 3H), 3.47 (s, 1H), 4.61 (t, 2H), 6.98-7.07 (m, 4H), 7.25-7.42 (m, 5H), 8.26 (s, 1H);
20 HPLC-MS : m/z = 324.1 (M+1); R_t = 3.36 min.

Example 596 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(pyrimidin-2-yloxy)-ethyl]-phenyl ester

25 The title compound was prepared in 24% yield as light yellow crystals using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 2-hydroxypyrimidine. NMR (400MHz; $CDCl_3$): δ 3.11 (t, 2H), 3.42 (br s, 3H), 4.54 (t, 2H), 6.93 (t, 1H), 7.05 (d, 2H), 7.25-7.42 (m, 7H), 8.51 (d, 2H); HPLC-MS : m/z = 350.2 (M+1); R_t = 2.86 min.

Example 597 (General procedure 26)

30 Methyl-phenyl-carbamic acid 4-[2-(pyridin-4-ylsulfanyl)-ethyl]-phenyl ester

The title compound was prepared as its TFA salt in 5% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 4-mercaptopyridine. NMR (400MHz;
35 $CDCl_3$): δ 3.07 (t, 2H), 3.36 (t, 2H), 3.44 (br s, 3H), 7.06 (br d, 2H), 7.20 (d, 2H), 7.25-7.43

(m, 7H), 8.51 (d, 2H); HPLC-MS : m/z = 365.2 (M+1); R_t = 2.53 min.

Example 598 (General procedure 12)

4-(3-Amino-phenyl)-piperidine-1-carboxylic acid 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl ester,

5

The title product was prepared from 4-(3-aminophenyl)piperidine (released from the correspondent hydrochloride by a standard procedure) and 4-(5-trifluoromethyl-pyridin-2-yloxy)-phenyl chloroformate, preparative HPLC (method C) (reaction performed in a mixture of dichloromethane and dimethylformamide, 5:3). 1.7 M HCl in ethyl acetate was added to the pooled fractions containing the title product, and the fractions was evaporated to dryness (7%, light yellow solid). HPLC-MS m/z = (M+1) 458.0, R_t : 3.09 min.

10

Example 599 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(1-pyridin-3-yl-1H-imidazol-2-ylsulfanyl)-ethyl]-phenyl ester

15

The title compound was prepared as its TFA salt in 5% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and 3-(2thio-1H-imidazol-1-yl)pyridine. NMR (400MHz; $CDCl_3$): δ 2.95 (t, 2H), 3.42 (br s, 2H), 3.56 (t, 3H), 6.98 (br d, 2H), 7.05 (d, 2H), 7.15 (d, 1H), 7.23-7.43 (m, 5H), 7.48 (d, 1H), 7.53 (dd, 1H), 7.70 (ddd, 1H), 8.65 (d, 1H), 8.77 (dd, 1H); HPLC-MS : m/z = 431.2 (M+1); R_t = 2.79 min.

20

Example 600 (General procedure 26)

Methyl-phenyl-carbamic acid 4-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethyl]-phenyl ester

25

The title compound was prepared in 39% yield as an oil using methyl-phenyl-carbamic acid 4-(2-hydroxy-ethyl)-phenyl ester and isoindole-1,3-dione. NMR (400MHz; $CDCl_3$): δ 2.98 (t, 2H), 3.41 (br s, 2H), 3.89 (t, 3H), 7.04 (br d, 2H), 7.20-7.40 (m, 7H), 7.68-7.71 (m, 2H), 7.80-7.83 (m, 2H); HPLC-MS : m/z = 401.1 (M+1); R_t = 4.41 min.

30

Example 601

4-Phenyl-piperidine-1-carboxylic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester

To a solution of 4-(5-methyl-pyridin-2-ylmethyl)phenol (0.8 mmol) prepared as described above and ethyldiisopropylamine (1.5 mmol) in CH_2Cl_2 (5 mL) at -30 °C was added trichloromethyl chloroformate (1.0 mmol). The solution was stirred at -30 °C for 10 min and at reflux

35

temperature for 2 h. The solution was evaporated to dryness and redissolved in CH₂Cl₂ (5 mL) and cooled to 0 °C before addition of 4-phenylpiperidine (1.5 mmol). The solution was stirred at room temperature for 16 h evaporated to give the crude product which was purified by FC (Quad flash 40 CH₂Cl₂:Et₂O:Heptane:Et₃N 1:1:2:0.25 ->1:1:1:0.25) to give the title

5 compound in 28% yield as an oil.

¹H NMR (400MHz; CDCl₃): δ 1.70-1.80 (m, 2H), 1.91 (br d, 2H), 2.29 (s, 3H), 2.72 (tt, 1H), 2.94 (br t, 1H), 3.08 (br t, 1H), 4.10 (s, 2H), 4.42 (br s, 1H), 7.00 (d, 1H), 7.06 (d, 2H), 7.21-7.26 (m, 5H), 7.32 (d, 2H), 7.38 (dd, 1H), 8.37 (d, 1H); HPLC-MS : *m/z* = 387.2 (M+1); R_t = 2.95 min.

10

Example 602

4-(4-Methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid 4-(5-methyl-pyridin-2-ylmethyl)-phenyl ester

15 To a solution of 4-(5-methyl-pyridin-2-ylmethyl)phenol (0.8 mmol) prepared as described above and ethyldiisopropylamine (1.5 mmol) in CH₂Cl₂ (5 mL) at -30 °C was added trichloromethyl chloroformate (1.0 mmol). The solution was stirred at -30 °C for 10 min and at reflux temperature for 2 h. The solution was evaporated to dryness and redissolved in CH₂Cl₂ (5 mL) and cooled to 0 °C before addition of 4-(4-methoxyphenyl)-1,2,3,6-tetrahydro-pyridine
20 (1.5 mmol). The solution was stirred at room temperature for 16 h evaporated to give the crude product which was purified by FC (Quad flash 40 CH₂Cl₂:Et₂O:Heptane:Et₃N 1:1:2:0.25 ->1:1:1:0.25) to give the title compound in 10% yield as colorless crystals.

¹H NMR (400MHz; CDCl₃): δ 2.30 (s, 3H), 2.60 (br s, 2H), 3.75-3.88 (m, 5H), 4.12 (s, 2H), 4.20 (br s, 1H), 4.30 (br s, 1H), 5.98 (br s, 1H), 6.88 (d, 2H), 7.00 (d, 1H), 7.07 (d, 2H), 7.24
25 (d, 2H), 7.34 (d, 2H), 7.39 (dd, 1H), 8.38 (d, 1H); HPLC-MS : *m/z* = 415.3 (M+1); R_t = 2.95 min.

PHARMACOLOGICAL METHODS

Compounds of formula I may be evaluated in vitro for their efficacy and potency to inhibit HSL, and such evaluation may be performed as described below.

5

ASSAYS

Hormone-sensitive lipase (HSL)

- 10 Materials. The Hormone-sensitive lipase was provided by Dr. Cecilia Holm, from Lund University Sweden or produced and purified by Novo Nordisk (NN) using the reagents and protocols used by Dr. Holm. The substrates used are: ^3H -labeled triolein (TO) from Amersham, Buckinghamshire, U.K. cat No. TRA191; 5-20 Ci/mmol dissolved in toluene, triolein (Sigma, Cat. No. T-1740), fluorochrome-labeled triacylglyceride (*cis*-octadec-9-enoic acid 2-[12-(7-nitrobenzo[1,2,5]oxadiazol-4-ylamino)dodecanoyloxy]-1-*cis*-octadec-9-enoyloxymethyl-ethyl ester) prepared by Novo Nordisk (NN) by conventional methods, and 1,3-(di[^3H]-stearin), 2-(PEG-Biotin)glycerol prepared in collaboration with Amersham Pharmacia Biotech, UK and described in WO 01/073442. Phosphatidyl choline (PC) and phosphatidyl inositol (PI) are from Sigma (St Luis MO cat. Nos. P-3556 and P-5954 respectively). All other reagents are of commercial grade and obtained from various commercial sources.
- 15
- 20

Methods.

3180.1: Assay for determination of inhibitor IC_{50} values.

- A lipid emulsion with ^3H -Triolein and phospholipid is used as substrate with a standard concentration of highly purified HSL. BSA is added as product receptor. The substrate is prepared as follows:
- 30 $30\ \mu\text{l}$ PC:PI (20 mg/ml solution of PC:PI 3:1 prepared in chloroform) + $128\ \mu\text{l}$ cold TO + $15\ \mu\text{l}$ ^3H -TO are mixed and then evaporated under a gentle stream of N_2 followed by 20-30 minutes in a Speedvac to ensure the absence of residual solvent.
- Compound and HSL are incubated for 30 min at $25\ ^\circ\text{C}$ before addition of substrate. Reaction is stopped after 30 min at $25\ ^\circ\text{C}$ by adding a mixture of methanol, chloroform and heptane at high pH. Formed product is separated from substrate by phase separation.
- Standard concentrations of compound are $100\ \mu\text{M}$, $20\ \mu\text{M}$, $4\ \mu\text{M}$, $0.8\ \mu\text{M}$, $0.16\ \mu\text{M}$ and $0.032\ \mu\text{M}$ (sample concentrations).
- 35 Results are given as IC_{50} values after 4PL fit of obtained activity data.

3180.2: Assay for determination of percent inhibition by compound at 10 μ M concentration. A lipid emulsion with 3 H-Triolein and phospholipid is used as substrate with a standard concentration of highly purified HSL. BSA is added as product receptor. The substrate is prepared as follows:

- 5 30 μ l PC:PI (20 mg/ml solution of PC:PI 3:1 prepared in chloroform) + 128 μ l cold TO + 15 μ l 3 H-TO are mixed and then evaporated under a gentle stream of N_2 followed by 20-30 minutes in a Speedvac to ensure the absence of residual solvent.

Compound and HSL are incubated for 30 min at 25 $^{\circ}$ C before addition of substrate. Reaction is stopped after 30 min at 25 $^{\circ}$ C by adding a mixture of methanol, chloroform and heptane at high pH. Formed product is separated from substrate by phase separation.

Results are given as percent activity relative to an un-inhibited sample (no compound).

3190.1: Assay for determination of percent inhibition of hormone sensitive lipase by compound at 10 μ M sample concentration.

A lipid emulsion with fluorochrome-labeled triacylglyceride and phospholipid is used as substrate with a standard concentration of highly purified HSL (12 μ g/mL initial concentration corresponding to 600ng/mL final concentration). BSA is added as product receptor. The transfer of the fluorochrome from the lipid phase to the water (BSA) phase changes the fluorescent properties of the fluorochrome. The changes can be monitored on a fluorimeter with an excitation wavelength of 450nm and an emission wavelength of 545nm.

Compound and HSL (20 μ L compound, 10 μ L enzyme and 70 μ L PED-BSA buffer) is pre-incubated for 30min at 25 $^{\circ}$ C before addition of substrate (100 μ L). Amount of formed product is measured after 120min incubation at 37 $^{\circ}$ C.

Results are given as percent activity relative to a non-inhibited sample (no compound).

3190.2: Assay for determination of IC_{50} value for the inhibition of hormone sensitive lipase by compound. Standard concentrations of compound are 100 μ M and 5-fold dilutions (initial concentration corresponding to 10 μ M final concentration and 5-fold).

A lipid emulsion with fluorochrome-labeled triacylglyceride and phospholipid is used as substrate with a standard concentration of highly purified HSL (12 μ g/mL initial concentration corresponding to 600ng/mL final concentration). BSA is added as product receptor. The transfer of the fluorochrome from the lipid phase to the water (BSA) phase changes the fluorescent

properties of the fluorochrome. The changes can be monitored on a fluorimeter with an excitation wavelength of 450nm and an emission wavelength of 545nm.

Compound and HSL (20µL compound, 10µL enzyme and 70µL PED-BSA buffer) is pre-incubated for 30min at 25°C before addition of substrate (100µL). Amount of formed product is measured after 120min incubation at 37°C.

Results are given as IC₅₀ values after 4PL fit of obtained activity data.

2848.2: This high-volume screening assay uses para-nitrophenyl butyrate (p-NPB) as substrate for HSL. HSL cleaves p-NPB and the reaction is monitored as an increase in the concentration of para-nitrophenol (p-NP). p-NP can be monitored as an increase in UV-absorbance at 405 nm. The reaction is carried out at room-temp. for 20 min. The action is not stopped, but instead UV-abs is measured at a fixed time (20 min.) Due to autohydrolysis of the substrate the reaction is read at t = 0 min. too and t = 20 min. and the increase in Abs is calculated as the difference between the two readings. When a compound that inhibits HSL is present, it results in a relative decrease in UV-absorbance.

$$\% \text{Eff} (\% \text{Inhibition}) = (S - S0\text{Eff}) / (S_{\text{maxEff}} - S0\text{Eff}) \times 100$$

Where S = signal in UV-abs., S0Eff = assaybuffer alone, SmaxEff = Assaybuffer with the lipase inhibitor.

2898.2: This method is an enzyme assay based upon SPA (scintillation proximity assay) particles. The substrate, 1,3-(di-[³H]-stearin), 2-(PEG-Biotin)-glycerol, is marked with ³H in both fatty acid moieties in the tri-glyceride. The third moiety of the tri-glyceride is a PEG linked Biotin. The substrate binds through Biotin to streptavidin in the SPA particles and the proximity between the radioactive tritium in the stearic acid moiety and the SPA particle results in emission of light from the SPA particles. The assay is an on-bead assay where HSL degrades the substrate, where after ³H-stearin acid is released from the bead. The amount of light emitted is proportional to the amount of substrate bound to the receptor. When a compound that inhibits the activity of HSL is present it results in a decrease in degradation of the substrate and thus an increase in the amount of light emitted and a concomitant increase in %Eff.

$$\% \text{Eff } (\% \text{Inhibition}) = (S - S0\text{Eff}) / (S_{\text{maxEff}} - S0\text{Eff}) \times 100$$

Where S = signal in dpm., S0Eff = no inhibitor added, SmaxEff = with maximum concentration of inhibitor.

5

RESULTS

10

With these methods the following results were obtained for the compounds of the examples.


Compound according to example #	Test 3190.1 HSL_FL % ACTIVITY All	Test 3180.2 HSL % ACTIVITY All	Test 2898.2 HSL-2 % Inhibition All	Test 2848.2 HSL % Inhibition All
120		40		
122		14		
123		22		
121		24		
118		24		
119		10		
47	42			
46	43			
11	17			
87	48			
81	36			
48	15			
61	23			
6	49			
85	88			

90	24			
54	52			
62	88			
59	16			
24	30			
20	51			
92		62		
73	39			
109				15.9
106				23.8
108				37.2
114				42.4
126				33
110				18.8
111				36.2
107				59.9
105				26.8
104				19.8
127				58.7
128				33.8
103		2	101.4	
115		85	73.6	
113			18.3	
129				27
112			86	
116				23.5
124			75.1	
117		2		
130				30.1

131				21.6
132				19.6
133			15.8	
144	25			

CLAIMS

1. Use of a compound of the general formula I



wherein R¹ is selected from hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and


X is O or S; and

L is a group such that -C(=X)-L is a hydrolysable group; or

- 5 a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs

10 for inhibition of the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters.

2. Use of a compound of the general formula I



15

wherein R¹ is selected from hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

- 20 R² is selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo,
- 25
- 30

halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

10 X is O or S; and

L is a group such that -C(=X)-L is a hydrolysable group; or

a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs

for the preparation of a medicament for the treatment of any disorder where it is desirable to

20 modulate the plasma level of free fatty acids, glycerol, LDL-cholesterol, HDL-cholesterol, insulin and/or glucose; and/or

modulate intracellular triacylglycerol and cholesterol ester stores, intracellular level of fatty acids, fatty acid esters such as diacylglycerols, phosphatidic acids, long chain acyl-CoA's as well as citrate or malonyl-CoA; and/or

increase insulin sensitivity in adipose tissue, skeletal muscle, liver or pancreatic β cells; and/or

30 modulate insulin secretion from pancreatic β cells.

3. Use of a compound of the general formula I according to claim 1 for the preparation of a pharmaceutical medicament.

4. Use of a compound of the general formula (I) according to claim 1 for the preparation of a medicament for the treatment of insulin resistance, diabetes type 1, diabetes type 2, metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

5

5. The use according to any one of claims 1-4, wherein X is O.


6. The use according to any one of claims 1-5, wherein the group L comprises an O, via which L is bound to the C in formula (I).

10


7. The use according to any one of claims 1-5, wherein the group L comprises a N, via which L is bound to the C in formula (I).


15

8. The use according to any one of claims 1-5, wherein the group L is selected from the group consisting of



379





wherein Y is O or S; and

- 5 R^{a1} , R^{a2} , R^{a3} , R^{a4} , R^{a5} and R^{a6} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, or C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino,

- cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy.
9. The use according to claim 8, wherein at least one of R^{a1}, R^{a2}, R^{a3}, R^{a4}, R^{a5} and R^{a6} is F.
10. The use according to any one of claims 1-6, wherein the group L is an optionally substituted -O-phenyl, via which L is bound to the C in formula (I).
11. The use according to any one of claims 1-10, wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperazine, said piperazine being bound to the C in formula (I).
12. The use according to any one of claims 1-10, wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperidine, said piperidine being bound to the C in formula (I).
13. The use according to any one of claims 1-10, wherein R¹ is selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro.
14. The use according to any one of claims 1-10, wherein R¹ is methyl.
15. The use according to any one of claims 1-10, wherein R² is phenyl.

16. The use according to any one of claims 1-10, wherein R^2 is a heteroaryl.

17. The use according to any one of claims 1-10, wherein R^1 is methyl and R^2 is phenyl.

5

18. The use according to any one of claims 1-6, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 and R^2 are covalently bound to each other so that the group R^1 -N- R^2 forms a piperazine, said piperazine being bound to the C in formula (I).

10

19. The use according to any one of claims 1-6, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 and R^2 are covalently bound to each other so that the group R^1 -N- R^2 forms a piperidine, said piperidine being bound to the C in formula (I).

15

20. The use according to any one of claims 1-6, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R^1 is methyl and R^2 is phenyl.

20

21. The use according to any one of claims 1-20, wherein pK_a of the group L is between 4 and 12, between 6 and 12, between 7 and 12, between 8 and 12, preferably between 8.5 to 11.5, and most preferable between 9.0 to 11.0.

25

22. The use according to any one of claims 1-21, wherein administration of said compound is by the oral route.

23. The use according to any one of claims 1-21, wherein administration of said compound is by the nasal, transdermal, pulmonal, or parenteral route.


30

24. A method of treating any disorder where it is desirable to inhibit the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters, wherein said method comprises the use according to any one of claims 1-23.

25. A method of treating any disorder where it is desirable to modulate the plasma level of free fatty acids or to modulate the handling, storage and oxidation of intracellular fatty acid and cholesterol, wherein said method comprises the use according to any one of claims 1-23.

26. The method according to any one of claims 24-25, wherein said disorder is selected from the group consisting of insulin resistance, diabetes type 1, diabetes type 2, metabolic syndrome X, impaired glucose tolerance, hyperglycemia, dyslipidemia, obesity, abnormalities of lipoprotein metabolism and any combination thereof.

27. A pharmaceutical composition comprising a compound of formula I



wherein R^1 is selected from hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{3-10} -cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro; and

R^2 is selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl.

₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy; and

5

wherein R² is optionally covalently bound to R¹ by an ether, thioether, C-C or C-N bond, to form a ring system with the N-atom to which R¹ and R² are bound; and

X is O or S; and

10

L is a group such that -C(=X)-L is a hydrolysable group; or

a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs

15

28. The pharmaceutical composition according to claim 27, wherein X is O.

29. The pharmaceutical composition according to any one of claims 27-28, wherein the group L comprises an O, via which L is bound to the C in formula (I).


20

30. The pharmaceutical composition according to any one of claims 27-28, wherein the group L comprises a N, via which L is bound to the C in formula (I).


31. The pharmaceutical composition according to any one of claims 27-29, wherein the group L is selected from the group consisting of

25


385



386



387



wherein Y is O or S; and

- 5 R^{a1} , R^{a2} , R^{a3} , R^{a4} , R^{a5} and R^{a6} are independently selected from hydrogen, hydroxy, sulfanyl, sulfo, halogen, amino, cyano, nitro, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl, or C_{3-10} -cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C_{1-6} -alkyl, C_{2-6} -alkenyl, aryl, heteroaryl, C_{3-8} -heterocyclyl and C_{3-10} -cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino,

- cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl, wherein each of hydroxy, sulfanyl, sulfo, amino, C₁₋₆-alkyl, C₂₋₆-alkenyl, aryl, heteroaryl, C₃₋₈-heterocyclyl and C₃₋₁₀-cycloalkyl is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, sulfo, oxo, halogen, amino, cyano, nitro, C₁₋₆-alkyl, C₂₋₆-alkenyl, perhalomethyl and perhalomethoxy.
32. The pharmaceutical composition according to claim 31, wherein at least one of R^{a1}, R^{a2}, R^{a3}, R^{a4}, R^{a5} and R^{a6} is F.
33. The pharmaceutical composition according to any one of claims 27-29, wherein the group L is an optionally substituted -O-phenyl, via which L is bound to the C in formula (I).
34. The pharmaceutical composition according to any one of claims 27-33, wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperazine, said piperazine being bound to the C in formula (I).
35. The pharmaceutical composition according to any one of claims 27-33, wherein R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperidine, said piperidine being bound to the C in formula (I).
36. The pharmaceutical composition according to any one of claims 27-33, wherein R¹ is selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl and C₃₋₁₀-cycloalkyl, each of which is optionally substituted with one or more substituents independently selected from hydroxy, sulfanyl, oxo, halogen, amino, cyano and nitro.
37. The pharmaceutical composition according to any one of claims 27-33, wherein R¹ is methyl.

38. The pharmaceutical composition according to any one of claims 27-33, wherein R² is phenyl.

5 39. The pharmaceutical composition according to any one of claims 27-33, wherein R² is a heteroaryl.

40. The pharmaceutical composition according to any one of claims 27-33, wherein R¹ is methyl and R² is phenyl.

10

41. The pharmaceutical composition according to any one of claims 27-29, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperazine, said piperazine being bound to the C in formula (I).

15

42. The pharmaceutical composition according to any one of claims 27-29, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R¹ and R² are covalently bound to each other so that the group R¹-N-R² forms a piperidine, said piperidine being bound to the C in formula (I).

20

43. The pharmaceutical composition according to any one of claims 27-29, wherein the group L is an optionally substituted -O-phenyl via which L is bound to the C in formula (I), and R¹ is methyl and R² is phenyl.

25

44. The pharmaceutical composition according to any one of claims 27-43, wherein pK_a of the group L is between 4 and 12, between 6 and 12, between 7 and 12, between 8 and 12, preferably between 8.5 to 11.5, and most preferable between 9.0 to 11.0.

30

45. The pharmaceutical composition according to any one of claims 27-44, which is for oral administration.

46. The pharmaceutical composition according to any one of claims 27-44, which is for administration by the nasal, transdermal, pulmonal, or parenteral route.

35

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 2003/051842 A3

(51) International Patent Classification⁷: **A61K 31/325**,
31/4196, 31/4192, 31/415, 31/352, 31/4035, 31/40,
31/404, 31/423, 31/42, 31/428, 31/44, 31/495, 31/505,
31/41

DK-2880 Bagsværd (DK). JACOBSEN, Poul; Morænevej
2A, DK-3500 Slangerup (DK).

(21) International Application Number:
PCT/DK2002/000853

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC,
VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
13 December 2002 (13.12.2002)

(25) Filing Language: English

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(26) Publication Language: English

(30) Priority Data:
PA 2001 01879 14 December 2001 (14.12.2001) DK
PA 2002 00645 30 April 2002 (30.04.2002) DK
PA 2002 01000 27 June 2002 (27.06.2002) DK
PA 2002 01562 11 October 2002 (11.10.2002) DK

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé,
DK-2880 Bagsværd (DK).

(88) Date of publication of the international search report:
3 June 2004

(72) Inventors: EBDROP, Søren; Nymarksvej 11, DK-4000
Roskilde (DK). HANSEN, Holger, Claus; An-
nexgårdsparken 43, DK-3500 Værløse (DK). VEDSØ,
Per; Dalgas Have 4, 2th, DK-2000 Frederiksberg (DK).
CORNELIS DE JONG, Johannes; Møllemarken 16, 1th,

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS DECREASING ACTIVITY OF HORMONE-SENSITIVE LIPASE

(57) Abstract: Carbamates and related compounds of formula R1R2NCO-Y or R1R2NCS-Y wherein R1 and R2 may form a ring, are inhibitors of hormone sensitive lipase and are useful for the treatment of any disorder where it is desirable to modulate the plasma level of free fatty acids, glycerol, LDL-cholesterol, HDL-cholesterol, insulin and/or glucose; and or modulate intracellular triacylglycerol and cholesterol stores, intracellular level of fatty acids, fatty acid esters such as diacylglycerols, phosphatidic acids, long chain acyl-CoA's as well as citrate or malonyl-CoA; and or increase insulin sensitivity; and or modulate insulin secretion from pancreatic beta cells. More particularly said disorders are selected from the group consisting of insulin resistance, diabetes type 1, hyperglycemia, dislipidemia, obesity, abnormalities of lipoprotein metabolism and any combinations thereof.



WO 2003/051842 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/325 A61K31/4196 A61K31/4192 A61K31/415 A61K31/352
A61K31/4035 A61K31/40 A61K31/404 A61K31/423 A61K31/42
A61K31/428 A61K31/44 A61K31/495 A61K31/505 A61K31/41

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BIOSIS, EMBASE, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/69810 A (NOVO NORDISK AS ; AGOURON PHARMA (US)) 23 November 2000 (2000-11-23) abstract; examples 398-421 -----	1-6, 8-10, 13, 20-29, 31, 33, 36, 44-46
X	WO 02/20483 A (HOFFMANN LA ROCHE) 14 March 2002 (2002-03-14) abstract examples 1.3, 1.11-1.14, 3.5, 3.6, 3.11, 3.12, 4.5-4.8, 4. 13, 4.14, claims 24, 25, 31, 33, 35; examples 4.19-4.22, 6, 7 ----- -/-	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 December 2003

Date of mailing of the international search report

31 03. 2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

A. Jakobs

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 02/06189 A (HOFFMANN LA ROCHE) 24 January 2002 (2002-01-24)</p> <p>abstract page 9, line 26 - page 10, line 10 page 17, lines 13-19 page 19, lines 19-26 page 21, line 27 - page 22, line 2 page 23, line 3 - page 24, line 4 page 35, line 26 - page 37, line 11; examples 4.1-4.17 claims 1-20,24; examples 9.1-9.4</p>	<p>1-6, 8-10,13, 14, 21-29, 31-33, 36,37, 39,44-46</p>
X	<p>WO 99/62870 A (ANDERSSON KJELL ;ASTRA AB (SE); LINDSTEDT ALSTERMARK EVA LOTTE (SE) 9 December 1999 (1999-12-09)</p> <p>abstract page 20, line 25 - page 21, line 15</p>	<p>1-6,8,9, 21-29, 31,44-46</p>
X	<p>WO 02/38153 A (BESENCON OLIVIER ;OLSSON ROLF (SE); BIOVITRUM AB (SE); OEHMAN JOHA) 16 May 2002 (2002-05-16)</p> <p>abstract page 5, lines 1-26 claims 1-9; examples 6-13,17-26,28-39; tables 2,3</p>	<p>1-6, 8-10, 21-29, 31-33, 44-46</p>
X	<p>WO 01/021584 A (GENENTECH INC ;JACKSON DAVID Y (US); SAILES FREDERICK C (US); SUTH) 29 March 2001 (2001-03-29)</p> <p>abstract page 19, line 23 - page 20, line 2; claims 1-24; examples 1-9,11-42,44-57,111-128; tables 2,3</p>	<p>1-6,8, 10, 13-16, 21-29, 31,33, 36-39, 44-46</p>
X	<p>WO 00/63208 A (BOEHRINGER INGELHEIM INT ;NOVO NORDISK AS (DK)) 26 October 2000 (2000-10-26)</p> <p>abstract page 23, line 22 - page 28, line 4 page 55, line 18 - page 56, line 26; claims 1-47</p>	<p>1</p>

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 199709 Derwent Publications Ltd., London, GB; Class B03, AN 1997-100136 XP002230558 & WO 97/01533 A (GREEN CROSS CORP) 16 January 1997 (1997-01-16) abstract</p>	<p>1-6,8, 13, 21-29, 31,36, 44-46</p>
X	<p>----- EP 1 201 649 A (PFIZER PROD INC) 2 May 2002 (2002-05-02) abstract page 17, lines 31-51 page 24, line 37 - page 25, line 40; claims 1-13; examples 1-87</p>	<p>1</p>
X	<p>----- US 5 837 727 A (BAUER CARL-AXEL WILHELM EDVARD ET AL) 17 November 1998 (1998-11-17)</p> <p>abstract; claims 1-7</p>	<p>1-6,8, 10,13, 14, 20-29, 31,33, 36,37, 44-46</p>
X	<p>----- US 5 463 062 A (HEMMERLE HORST ET AL) 31 October 1995 (1995-10-31)</p> <p>abstract column 5, lines 26-47; claims 1-11; examples 102,103</p>	<p>1-6,8, 13, 21-29, 31,44-46</p>
X	<p>----- WO 98/53814 A (HAGMANN WILLIAM K ;MUMFORD RICHARD A (US); KEVIN NANCY J (US); MAC) 3 December 1998 (1998-12-03)</p> <p>abstract page 20, line 21 - page 21, line 10; claims 17-20; examples 272,319-324</p> <p>----- -/--</p>	<p>1-6,8, 10,13, 14, 20-29, 31,33, 36,37, 44-46</p>

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/54759 A (TULARIK INC) 21 September 2000 (2000-09-21) abstract page 2, lines 13-29 page 3, line 25 - page 4, line 22 page 13, lines 1-7; claims 1,9,27,28,40-45; examples 7-7.6 -----	1-6,8, 10,13, 14, 21-29, 31,33, 36,37, 44-46
X	WO 01/021602 A (DEVASTHALE PRATIK ;SQUIBB BRISTOL MYERS CO (US); CHEN SEAN (US); J) 29 March 2001 (2001-03-29) page 1, line 1 - page 16, line 27 -----	1-6,8, 10,13, 20-29, 31,33, 44-46
X,P	WO 02/12210 A (BERNARDON JEAN MICHEL ;CLARY LAURENCE (FR); GALDERMA RES & DEV (FR) 14 February 2002 (2002-02-14) abstract; examples 11,19-21,27,61,70,73,74 page 8, lines 37,38 page 9, lines 4-6,12-14 page 14, line 6 - page 15, line 19 -----	1-6,8, 10,13, 15, 21-29, 31,33, 36,38, 44-46
E	WO 03/040114 A (DEVASTHALE PRATIK ;SQUIBB BRISTOL MYERS CO (US); JEON YOON T (US)) 15 May 2003 (2003-05-15) abstract page 6, line 1 - page 7, line 17 -----	1-6,8, 10,13, 21-29, 31,33, 36,44-46
X	WO 92/07838 A (BEECHAM GROUP PLC) 14 May 1992 (1992-05-14) abstract page 1, line 15 - page 3, line 7; examples 1-5,11,12 ----- -/--	1-6,8, 10, 13-15, 21-29, 31,33, 36-38, 44-46

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00853

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 01/087843 A (CORNELIS DE JONG JOHANNES ;JACOBSEN POUL (DK); NOVO NORDISK AS (DK) 22 November 2001 (2001-11-22)</p> <p>abstract page 3, line 1 - page 18, line 9 page 25, line 32 - page 26, line 9; claims 1-22; examples 1-50; table 1</p> <p>-----</p>	<p>1-6, 8-10, 21-29, 31-33, 39,44-46</p>
X	<p>US 5 869 484 A (MARCHESINI DONATA ET AL) 9 February 1999 (1999-02-09)</p> <p>abstract; examples 1-28</p> <p>-----</p>	<p>27-29, 31,33, 36,44-46</p>
X	<p>WO 01/047923 A (ASTRAZENECA AB ;BARE THOMAS MICHAEL (US); BROWN DEAN GORDON (US);) 5 July 2001 (2001-07-05)</p> <p>abstract page 20, lines 22-25; table 1</p> <p>-----</p>	<p>27-29, 31,33, 36-38, 40,43-46</p>
X	<p>US 4 643 991 A (TSUJI KIYOSHI ET AL) 17 February 1987 (1987-02-17)</p> <p>abstract; examples 6-11,16,17,19-21,23-31,33-38,40-42,44-46,48-50 examples 52-63,65-67,69-83,85-87,93,97-99</p> <p>-----</p>	<p>27-29, 31-33, 36,37, 44-46</p>
X	<p>WO 95/29155 A (PHARMACIA AB ;FEX TOMAS (SE); STAMVIK ANDERS (SE); CARLSSON JAN IN) 2 November 1995 (1995-11-02)</p> <p>abstract page 14, lines 21-25; examples 4,5 page 27, line 31 - page 28, line 20 page 57, lines 24-30 page 59, lines 14-16,21-23 page 61, lines 1-3 page 66, lines 19-25 page 68, lines 1-3 page 71, lines 1-7 page 72, lines 19-21,26-29 page 74, lines 5-7; claims 10-12</p> <p>-----</p>	<p>27-29, 31,33, 36,37, 44-46</p>
X	<p>BE 665 273 A (FARBENFAB BAYER AG) 1 October 1965 (1965-10-01)</p> <p>page 6, paragraph 3 - page 12, paragraph 4</p> <p>-----</p> <p>-/--</p>	<p>27-29, 31,33, 36-38, 40,43-46</p>

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00853

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	<p>WO 02/42257 A (CASTRO PALOMINO JULIO ;EBBINGHAUS KINTSCHER ULRICH (DE); SCHUHMACH) 30 May 2002 (2002-05-30)</p> <p>abstract page 57, line 20 - page 58 page 82; examples XXXVI,XV examples 148-151,157-176,204,205,231,232,283,284</p>	<p>27-29, 31,33, 36,38, 44-46</p>
X	<p>EP 0 587 311 A (SANKYO CO) 16 March 1994 (1994-03-16) examples 1-12,15,16,19,22,24,26,28,29,34,36,37,42,4 9,51-53,57,59,65,66,68,70,71,73,74,76,77,8 8-94, abstract; claims 19,20</p>	<p>1</p>
X	<p>SCHMIDT F ET AL: "Glucuronide prodrugs of hydroxy compounds for antibody directed enzyme prodrug therapy (adept): a phenol nitrogen mustard carbamate" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 7, no. 8, 22 April 1997 (1997-04-22), pages 1071-1076, XP004136187 ISSN: 0960-894X abstract</p>	<p>1</p>
A	<p>WO 92/06075 A (WELLCOME FOUND) 16 April 1992 (1992-04-16) abstract; example 15</p>	<p>1</p>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 02/00853

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1,4-26 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-6,8-10,13-17,20-29,31-33,36-40,44-46 (all partially) 43

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-6,8-10,13-17,20-29,31-33,36-40,44-46 (all partially)
43

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the first, 27th, 28th, 29th, 30th, 46th 47th and 48th group of claim 8 (-Y-Ra1, Ra1 in -Y-Ra1 having any value given in claim 8 (and in particular substituted 0-phenyl) except heteroaryl and heterocyclyl).

2. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 2, 13 and 14 of claim 8 (1,2,4-triazolyl group).

3. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the 3rd and 4th group of claim 8 (1,2,3-benzotriazolyl group).

4. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 5 or 24 of claim 8 (pyrazolyl group).

5. claims: 1-9,13-17,20-29,31-32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 6 or 17 of claim 8 (imidazolyl group).

6. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 7, 8, 10 or 12 of claim 8 (Y-N=CRa1-H, Y-N=CRa1-Cl, Y-NCRa2CH=CHRa1, Y-N=CRa1Ra2).

7. claims: 1-6,8,9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 9 of claim 8 (indol-1-yl group).

8. claims: 1-6,8,9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 11 of claim 8 (3H-benzo(d)-1,2,3-triazin-4-one-3-yl group).

9. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 15 and 16 of claim 8 (1,2,3-triazolyl group).

10. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 18 and 19 of claim 8 (pyrrolyl group).

11. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 20, 21, 22, 23 of claim 8 (tetrazolyl group).

12. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 25, 26 of claim 8 (tetrahydro-1,2,3,4-isoquinolin-7-yl)(benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

13. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 31-34 of claim 8 (benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

14. claims: 1-9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 35-37 of claim 8 (benzisoxazol-1-yl, benzisothiazol-1-yl, 3H-isoindol-1-yl group).

15. claims: 1-6,8,9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 38, 39 or 40 of claim 8 (pyridinyl group) and the benzocondensed derivatives thereof like quinolin-2-yl of e.g example 244.

16. claims: 1-6,8,9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group consisting of the group 41, 43, 45 of claim 8 (pyrimidin-2-yl, pyrimidin-3-yl or pyrimidin-4-yl).

17. claims: 1-6,8,9,13-17,20-29,31,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is selected from the group 42 and 44 of claim 8 (pyridazin-yl group).

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

18. claims: 1-6,8,9,13-17,20-29,31-,32,36-40,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is not covalently bound to R2 and wherein L is heteroaryl or heterocyclyl not provided for in inventions 2-17, e.g. chromenyl as of examples 12, 40-49.

19. claims: 1-6,8-10,12,13,19,21-28,31-33,35,36,44-46 (all partially) 42

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the first, 27th, 28th, 29th, 30th, 46th 47th and 48th group of claim 8 (-Y-Ra1, Ra1 in -Y-Ra1 having any value given in claim 8 (and in particular substituted O-phenyl) except heteroaryl and heterocyclyl).

20. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 2, 13 and 14 of claim 8 (1,2,4-triazolyl group).

21. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the 3rd and 4th group of claim 8 (1,2,3-benzotriazolyl group).

22. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 5 or 24 of claim 8 (pyrazolyl group).

23. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 6 or 17 of claim 8 (imidazolyl group).

24. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 7, 8, 10 or 12 of claim 8 (Y-N=CRa1-H, Y-N=CRa1-Cl, Y-NCRa2CH=CHRa1, Y-N=CRa1Ra2).

25. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 9 of claim 8 (indol-1-yl group).

26. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 11 of claim 8 (3H-benzo(d)-1,2,3-triazin-4-one-3-yl group).

27. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 15 and 16 of claim 8 (1,2,3-triazolyl group).

28. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 18 and 19 of claim 8 (pyrrolyl group).

29. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 20, 21, 22, 23 of claim 8 (tetrazolyl group).

30. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 25, 26 of claim 8 (tetrahydro-1,2,3,4-isoquinolin-7-yl)(benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

31. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 31-34 of claim 8 (benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

INTERNATIONAL SEARCH REPORT

International Application No. ~~DP07~~ 00853 /

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

32. claims: 1-9,12,13,19,21-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 35-37 of claim 8 (benzisoxazol-1-yl, benzisothiazol-1-yl, 3H-isoindol-1-yl group).

33. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 38, 39 or 40 of claim 8 (pyridinyl group) and the benzocondensed derivatives thereof.

34. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group consisting of the group 41, 43, 45 of claim 8 (pyrimidin-2-yl, pyrimidin-3-yl or pyrimidin-4-yl).

35. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is selected from the group 42 and 44 of claim 8 (pyridazin-yl group).

36. claims: 1-6,8,9,12,13,19,21-28,30-32,35,36,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by an ether, thioether or C-C bond to form a N-containing ring system, (e.g. morpholine, thiomorpholine, piperidine) and wherein L is heteroaryl or heterocyclyl not provided for in inventions 20-35.

37. claims: 1-6,8-11,18,20-28,31-34,45,46 (all partially) 41

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the first, 27th, 28th, 29th, 30th, 46th 47th and 48th group of claim 8 (-Y-Ra1, Ra1 in -Y-Ra1 having any value given in claim 8 (and in particular substituted O-phenyl) except heteroaryl and heterocyclyl).

38. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 2, 13 and 14 of claim 8 (1,2,4-triazolyl group).

39. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the 3rd and 4th group of claim 8 (1,2,3-benzotriazolyl group).

40. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 5 or 24 of claim 8 (pyrazolyl group).

41. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 6 or 17 of claim 8 (imidazolyl group).

42. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 7, 8, 10 or 12 of claim 8 (Y-N=CRa1-H, Y-N=CRa1-Cl, Y-NCRa2CH=CHRa1, Y-N=CRa1Ra2).

43. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 9 of claim 8 (indol-1-yl group).

44. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 11 of claim 8 (3H-benzo(d)-1,2,3-triazin-4-one-3-yl group).

45. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 15 and 16 of claim 8 (1,2,3-triazolyl group).

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

46. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 18 and 19 of claim 8 (pyrrolyl group).

47. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 20, 21, 22, 23 of claim 8 (tetrazolyl group).

48. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 25, 26 of claim 8 (tetrahydro- 1,2,3,4-isoquinolin-7-yl)(benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

49. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 31-34 of claim 8 (benzimidazol-2-yl, benzothiazol-2-yl-, benzoxazol-2-yl, 3H-indol-2-yl group).

50. claims: 1-9,11,18,21-32,34,36,39,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 35-37 of claim 8 (benzisoxazol-1-yl, benzisothiazol-1-yl, 3H-isoindol-1-yl group).

51. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 38, 39 or 40 of claim 8 (pyridinyl group) and the benzocondensed derivatives thereof as far .

52. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group consisting of the group 41, 43, 45 of claim 8 (pyrimidin-2-yl, pyrimidin-3-yl or pyrimidin-4-yl).

53. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is selected from the group 42 and 44 of claim 8 (pyridazin-yl group).

54. claims: 1-6,8-9,11,18,21-28,30,32,34,36,39,44-46 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Pharmaceutical composition comprising a compound of formula I and uses according to claims 1-26 wherein R1 is covalently bound to R2 by a C-N bond to form a N-containing ring system, (e.g. piperazine, 1,4-diazepane, 2,5-diazabicyclo[2.2.1]heptane) and wherein L is heteroaryl or heterocyclyl not provided for in inventions 38-53.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: -

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible.

Present claims 1-46 relate to an extremely large number of possible compounds/uses. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a small proportion of the compounds/uses claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Present claims 1-26 relate to a compound/use defined

by reference to the following parameters/expressions:

a hydrolysable group, wherein pKa of the group L is as specified in claim 21, inhibition of the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters, and disorders where it is desirable to inhibit the lipolytic activity of hormone-sensitive lipase against triacylglycerols, diacylglycerols, cholesterol acyl esters or steroid acyl esters; any disorder where it is desirable to modulate the plasma level of free fatty acids, glycerol, LDL-cholesterol, HDL-cholesterol, insulin and/or glucose and/or to modulate intracellular triacylglycerol, cholesterol ester stores, intracellular levels of fatty acids, fatty acid esters such as diacylglycerols, phosphatidic acids, long chain acyl-CoA's, as well as citrate or malonyl-CoA; and or increase insulin sensitivity in adipose tissue, skeletal muscle, liver or pancreatic beta cells, and or modulate insulin secretion from pancreatic beta cells, abnormalities of lipoprotein metabolism.

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible.

Consequently, the search has been restricted to pharmaceutical

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

compositions comprising the individual compounds of the description as far as relating to invention 1, and their use in relation to the treatment of diabetes type 1, diabetes type 2, metabolic syndrome X, dyslipidemia, hyperglycemia and obesity.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0069810 A	23-11-2000	AU 4537900 A	05-12-2000
		BR 0010651 A	19-03-2002
		CA 2373892 A1	23-11-2000
		CN 1356977 T	03-07-2002
		CZ 20013767 A3	17-04-2002
		WO 0069810 A1	23-11-2000
		EP 1183229 A1	06-03-2002
		HU 0201033 A2	29-07-2002
		JP 2002544254 T	24-12-2002
		NO 20015607 A	17-01-2002
		PL 352367 A1	25-08-2003
		US 2003220350 A1	27-11-2003
		US 6503949 B1	07-01-2003
		ZA 200108560 A	13-06-2002
WO 0220483 A	14-03-2002	AU 8591201 A	22-03-2002
		BR 0113752 A	29-07-2003
		CA 2419588 A1	14-03-2002
		CN 1468220 T	14-01-2004
		WO 0220483 A1	14-03-2002
		EP 1317432 A1	11-06-2003
		US 2002068753 A1	06-06-2002
WO 0206189 A	24-01-2002	AU 8964201 A	30-01-2002
		BR 0112609 A	01-07-2003
		CA 2415551 A1	24-01-2002
		CN 1443179 T	17-09-2003
		WO 0206189 A2	24-01-2002
		EP 1303503 A2	23-04-2003
		JP 2004504280 T	12-02-2004
		US 2002038025 A1	28-03-2002
WO 9962870 A	09-12-1999	AT 261429 T	15-03-2004
		AT 251130 T	15-10-2003
		AU 4667099 A	20-12-1999
		AU 752262 B2	12-09-2002
		AU 4667299 A	20-12-1999
		BR 9910913 A	06-03-2001
		BR 9910921 A	06-03-2001
		CA 2334107 A1	09-12-1999
		CA 2334374 A1	09-12-1999
		CN 1312795 T	12-09-2001
		CN 1311769 T	05-09-2001
		DE 69911770 D1	06-11-2003
		DK 1084102 T3	02-02-2004
		EE 200000717 A	15-08-2001
		EE 200000725 A	17-06-2002
		EP 1084101 A1	21-03-2001
		EP 1084102 A1	21-03-2001
		HU 0103226 A2	28-01-2002
		ID 28164 A	10-05-2001
		ID 29457 A	30-08-2001
		JP 2002516898 T	11-06-2002
		JP 2002516899 T	11-06-2002
		NO 20006114 A	02-02-2001
		NO 20006116 A	02-02-2001
		NZ 508453 A	30-06-2003
		PL 344681 A1	19-11-2001

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9962870 A		PL 344682 A1	19-11-2001
		WO 9962870 A1	09-12-1999
		WO 9962871 A1	09-12-1999
		SI 1084102 T1	29-02-2004
		SK 17672000 A3	06-08-2001
		SK 17692000 A3	10-05-2001
		TR 200003543 T2	20-04-2001
		TR 200003583 T2	21-05-2001
		TW 446694 B	21-07-2001
		TW 548263 B	21-08-2003
		US 6630600 B1	07-10-2003
		US 6362360 B1	26-03-2002
		ZA 200006771 A	20-05-2002
		ZA 200006773 A	20-02-2002
WO 0238153 A	16-05-2002	AU 1450502 A	21-05-2002
		WO 0238153 A1	16-05-2002
		US 2002198189 A1	26-12-2002
WO 0121584 A	29-03-2001	AU 7613800 A	24-04-2001
		CA 2385882 A1	29-03-2001
		EP 1214292 A1	19-06-2002
		JP 2003509488 T	11-03-2003
		WO 0121584 A1	29-03-2001
		US 6469047 B1	22-10-2002
WO 0063208 A	26-10-2000	AU 3956900 A	02-11-2000
		WO 0063208 A1	26-10-2000
		EP 1173438 A1	23-01-2002
		JP 2002542245 T	10-12-2002
WO 9701533 A	16-01-1997	WO 9701533 A1	16-01-1997
		JP 9071562 A	18-03-1997
EP 1201649 A	02-05-2002	BR 0104831 A	20-08-2002
		CA 2360308 A1	28-04-2002
		EP 1201649 A1	02-05-2002
		JP 2002193911 A	10-07-2002
		US 2002107235 A1	08-08-2002
US 5837727 A	17-11-1998	AT 138570 T	15-06-1996
		AU 653149 B2	22-09-1994
		AU 7569191 A	10-10-1991
		CA 2077762 A1	17-09-1991
		CS 9100681 A2	15-10-1991
		DE 69119896 D1	04-07-1996
		DK 521967 T3	09-09-1996
		EG 19645 A	30-09-1995
		EP 0521967 A1	13-01-1993
		ES 2087291 T3	16-07-1996
		GR 3020700 T3	31-10-1996
		HK 52197 A	02-05-1997
		HU 61461 A2	28-01-1993
		HU 211998 B	29-01-1996
		IE 910862 A1	25-09-1991
		IL 97509 A	31-12-1995
		JP 5504971 T	29-07-1993
		JP 3081640 B2	28-08-2000

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5837727 A		LT 1735 A ,B	25-07-1995
		LV 10917 A ,B	20-12-1995
		NZ 237279 A	26-08-1993
		PT 97052 A ,B	31-10-1991
		WO 9113615 A1	19-09-1991
		SG 64895 A1	25-01-2000
		SK 278815 B6	04-03-1998
		RU 2093153 C1	20-10-1997
		US 6034123 A	07-03-2000
		ZA 9101452 A	27-11-1991
US 5463062 A	31-10-1995	AT 137735 T	15-05-1996
		AU 662073 B2	17-08-1995
		AU 4616993 A	17-03-1994
		CA 2105709 A1	10-03-1994
		CN 1087622 A ,B	08-06-1994
		CZ 9301866 A3	16-03-1994
		DE 59302501 D1	13-06-1996
		DK 587088 T3	23-09-1996
		EP 0587088 A1	16-03-1994
		ES 2087625 T3	16-07-1996
		FI 933903 A	10-03-1994
		GR 3019957 T3	31-08-1996
		HK 1006708 A1	12-03-1999
		HU 65693 A2	28-07-1994
		IL 106936 A	12-03-1999
		JP 6211736 A	02-08-1994
		KR 275604 B1	15-12-2000
		NO 933200 A ,B,	10-03-1994
		NZ 248596 A	21-12-1995
		PH 30250 A	05-02-1997
		PL 300327 A1	21-03-1994
		RU 2126378 C1	20-02-1999
		SG 44819 A1	19-12-1997
		TW 399041 B	21-07-2000
		ZA 9306611 A	29-03-1994
WO 9853814 A	03-12-1998	CA 2291778 A1	03-12-1998
		EP 1001764 A1	24-05-2000
		JP 2002512625 T	23-04-2002
		WO 9853814 A1	03-12-1998
WO 0054759 A	21-09-2000	AU 3627300 A	04-10-2000
		CA 2367595 A1	21-09-2000
		EP 1161233 A2	12-12-2001
		JP 2002539155 T	19-11-2002
		WO 0054759 A2	21-09-2000
		US 6316503 B1	13-11-2001
WO 0121602 A	29-03-2001	AU 7593500 A	24-04-2001
		BR 0014189 A	20-08-2002
		CA 2388452 A1	29-03-2001
		CN 1374955 T	16-10-2002
		EP 1218361 A1	03-07-2002
		HU 0204416 A2	28-04-2003
		JP 2003509503 T	11-03-2003
		NO 20021408 A	14-05-2002
		TR 200200732 T2	21-10-2002

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0121602 A		WO 0121602 A1	29-03-2001
		US 2003069275 A1	10-04-2003
		US 2003096846 A1	22-05-2003
		US 2003087935 A1	08-05-2003
		US 6414002 B1	02-07-2002
		ZA 200200937 A	02-05-2003
WO 0212210 A	14-02-2002	FR 2812876 A1	15-02-2002
		AU 8598101 A	18-02-2002
		BR 0113251 A	24-06-2003
		CA 2418895 A1	14-02-2002
		CN 1468226 T	14-01-2004
		EP 1309575 A1	14-05-2003
		WO 0212210 A1	14-02-2002
		JP 2004505958 T	26-02-2004
		US 2004039038 A1	26-02-2004
WO 03040114 A	15-05-2003	WO 03040114 A1	15-05-2003
		US 2003130306 A1	10-07-2003
WO 9207838 A	14-05-1992	AU 8734791 A	26-05-1992
		EP 0555251 A1	18-08-1993
		WO 9207838 A1	14-05-1992
		JP 6502145 T	10-03-1994
WO 0187843 A	22-11-2001	AU 6007601 A	26-11-2001
		WO 0187843 A1	22-11-2001
		US 2002082254 A1	27-06-2002
US 5869484 A	09-02-1999	IT 1271679 B	04-06-1997
		AU 3078395 A	16-02-1996
		EP 0772608 A1	14-05-1997
		WO 9602524 A1	01-02-1996
WO 0147923 A	05-07-2001	AU 2420001 A	09-07-2001
		AU 2420101 A	09-07-2001
		AU 2420201 A	09-07-2001
		AU 2420301 A	09-07-2001
		AU 2566001 A	09-07-2001
		AU 2566101 A	09-07-2001
		AU 2566201 A	09-07-2001
		BG 106831 A	31-03-2003
		BG 106832 A	31-03-2003
		BR 0016646 A	08-10-2002
		BR 0016654 A	03-09-2002
		CA 2394561 A1	05-07-2001
		CA 2394888 A1	05-07-2001
		CN 1411459 T	16-04-2003
		CN 1434714 T	06-08-2003
		CZ 20022177 A3	13-11-2002
		EE 200200348 A	15-08-2003
		EE 200200350 A	15-10-2003
		EP 1244660 A1	02-10-2002
		EP 1244453 A1	02-10-2002
		EP 1244661 A1	02-10-2002
		EP 1244662 A1	02-10-2002
		EP 1248621 A1	16-10-2002
		EP 1244663 A1	02-10-2002

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0147923 A		EP 1244664 A1	02-10-2002
		HU 0300043 A2	28-05-2003
		HU 0300313 A2	28-06-2003
		JP 2003518499 T	10-06-2003
		JP 2003518500 T	10-06-2003
		JP 2003519145 T	17-06-2003
		JP 2003519146 T	17-06-2003
		JP 2003519147 T	17-06-2003
		JP 2003519148 T	17-06-2003
		JP 2003519149 T	17-06-2003
		NO 20022984 A	20-08-2002
		NO 20022990 A	20-08-2002
		NZ 519390 A	25-07-2003
		WO 0147523 A1	05-07-2001
		WO 0147923 A1	05-07-2001
		WO 0147924 A1	05-07-2001
		WO 0147524 A1	05-07-2001
		WO 0147925 A1	05-07-2001
		WO 0147926 A1	05-07-2001
		WO 0147927 A1	05-07-2001
		SK 8792002 A3	02-05-2003
		SK 8802002 A3	03-06-2003
		US 2003162783 A1	28-08-2003
		US 2003153571 A1	14-08-2003
US 4643991 A	17-02-1987	NONE	
WO 9529155 A	02-11-1995	AU 2354595 A	16-11-1995
		WO 9529155 A1	02-11-1995
BE 665273 A	01-10-1965	NONE	
WO 0242257 A	30-05-2002	DE 10058461 A1	19-09-2002
		AU 2483902 A	03-06-2002
		BR 0115611 A	06-01-2004
		CA 2429328 A1	30-05-2002
		WO 0242257 A1	30-05-2002
		EP 1339670 A1	03-09-2003
EP 0587311 A	16-03-1994	AU 4443893 A	10-02-1994
		CA 2103536 A1	08-02-1994
		CN 1091432 A	31-08-1994
		CZ 9301619 A3	16-02-1994
		EP 0587311 A1	16-03-1994
		FI 933472 A	08-02-1994
		HU 64774 A2	28-02-1994
		IL 106600 A	30-09-1997
		JP 6100533 A	12-04-1994
		KR 177838 B1	01-04-1999
		MX 9304837 A1	31-05-1994
		NO 932813 A	08-02-1994
		NZ 248332 A	27-01-1995
		RU 2106357 C1	10-03-1998
		US 5629406 A	13-05-1997
		ZA 9305697 A	10-03-1994
WO 9206075 A	16-04-1992	AU 8541591 A	28-04-1992
		EP 0550568 A1	14-07-1993

INTERNATIONAL SEARCH REPORT

International Application No
PCT/DK 02/00853

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9206075	A	WO 9206075 A1	16-04-1992
		JP 6501252 T	10-02-1994
		US 5422373 A	06-06-1995
